

NATIONAL SENIOR CERTIFICATE EXAMINATION NOVEMBER 2017

MATHEMATICS: PAPER I

MARKING GUIDELINES

Time: 3 hours 150 marks

These marking guidelines are prepared for use by examiners and sub-examiners, all of whom are required to attend a standardisation meeting to ensure that the guidelines are consistently interpreted and applied in the marking of candidates' scripts.

The IEB will not enter into any discussions or correspondence about any marking guidelines. It is acknowledged that there may be different views about some matters of emphasis or detail in the guidelines. It is also recognised that, without the benefit of attendance at a standardisation meeting, there may be different interpretations of the application of the marking guidelines.

IEB Copyright © 2017 PLEASE TURN OVER

SECTION A

QUESTION 1

(a)
$$(x-1)^2 = 2(1-x)$$

$$(x-1)^2 = -2(x-1)$$

$$(x-1)^2 + 2(x-1) = 0$$

$$(x-1)(x-1+2) = 0$$


$$(x-1)(x+1) = 0$$

$$x = 1 x = -1$$

(2)
$$5^{-x}.5^{x-2} = \frac{25^{2x}}{5}$$
$$5^{-x}.5^{x-2} = \frac{5^{4x}}{5^{1}}$$
$$5^{-x+x-2} = 5^{4x-1}$$
$$-2 = 4x - 1$$
$$x = -\frac{1}{4}$$

(b)
$$(x+1)^2 < 9$$

 $\therefore x^2 + 2x + 1 < 9$
 $\therefore x^2 + 2x - 8 < 0$
 $\therefore (x+4)(x-2) < 0$
Critical Values: -4; 2



Solution: $\{x: -4 < x < 2\}$

Alternative

$$(x+1)^2 < 9$$

 $\therefore -3 < x+1 < 3$
 $\therefore -4 < x < 2$


(c)
$$(x-2)(x+4) = 0$$

 $x^2 + 2x - 8 = 0$
 $b = 2$ and $c = -8$

Alternative

$$(x+1)^2 < 9$$

 $x^2 + 2x - 8 < 0$
Sketch: $y = x^2 + 2x - 8$
 $x - \text{int} : x = -4; x = 2$



Solution: $\{x: -4 < x < 2\}$

(d) (1)
$$x-2 = \frac{-4}{x-2} - 4 \text{ let } x-2 = y$$

 $y = -\frac{4}{y} - 4 \text{ LCD: } y$
 $y^2 = -4 - 4y$
 $\therefore y^2 + 4y + 4 = 0$

(2)
$$(y+2)^2 = 0$$

 $\therefore y = -2$

Roots are real and equal.

Alternative

$$y^{2} + 4y + 4 = 0$$

$$\therefore \Delta = 4^{2} - 4(1)(4)$$

$$\therefore \Delta = 0$$

.. Roots are real and equal.

Alternative

$$x-2 = \frac{-4}{x-2} - 4$$


$$\therefore (x-2)^2 = -4 - 4(x-2)$$

$$\therefore x^2 - 4x + 4 = -4 - 4x + 8$$

$$\therefore x^2 = 0$$

.. Roots are real and equal.

(a) (1)



(2)
$$E = \{HTT, THT, TTH\}$$

∴ P(2 tails and 1 head) =
$$\frac{3}{8}$$

(b) (1)
$$P(A \cap B) = 0$$


- (2) (i) You cannot pick a R2 and a R5 coin at the same time.
 - (ii) P(either a R5 or a R2)

$$= P(A \text{ or } B)$$

$$= 0.36 + 0.47$$

$$= 0.83$$

(c) (1)



(2) P(exactly one machine is stamping R5 coins)

$$=0,3+0,2$$

$$= 0,5$$

- (a) $480\ 163 \div 0{,}502 = R956\ 500$
- (b) $R956 500 \times 5\% = R47 825$
- (c) Cost of machinery including import charges = R956 500 + R47 825 = R1 004 325 $A = P \left(1+i\right)^{n}$ $1004 325 = 225 450 \left(1+\frac{9,5}{100}\right)^{n}$ $\frac{1004 325}{225 450} = \left(\frac{219}{200}\right)^{n}$ $\log_{\left(\frac{219}{200}\right)} \left(\frac{1004 325}{225 450}\right) = n$ n = 16,46171594 $\therefore n \approx 16,46$ $\therefore \text{ approx. 17 years}$
- (d) (1) Loan required: R1 004 325 R225 450 = R778 875

$$P = x \left[\frac{1 - (1+i)^{-n}}{i} \right]$$

$$778 875 = x \left[\frac{1 - \left(1 + \frac{12}{1200}\right)^{(-4\times12)}}{\frac{12}{1200}} \right]$$

$$x = R20 510,76607$$

$$\therefore x = R20 510,77$$

(2) Outstanding Balance = A - F

$$A = 778 875 \left(1 + \frac{12}{1200} \right)^{24}$$

$$A = 988 964,5744$$

$$A \approx 988 964,57$$

$$F = 20510,76607 \left[\frac{\left(1 + \frac{12}{1200}\right)^{24} - 1}{\frac{12}{1200}} \right]$$

F = 553 246,4277 $F \approx 553 246,43$

Outstanding balance = $988\ 964,5744 - 553\ 246,4277$ = $R435\ 718,1467 \approx R435\ 718,15$


NB: If A and F are rounded to the nearest cent, consider Outstanding balance = 988 964,57 – 553 246,43 = R435 718,14

Alternative

Outstanding balance = 20 510,76607
$$\frac{1 - \left(1 + \frac{12}{1200}\right)^{-24}}{\frac{12}{1200}}$$

= R435 718,1466

≈ R435 718,15



Constant second difference

(2)
$$T_n = an^2 + bn + c$$

 $T_1 = a + b + c = 1$
 $T_2 = 4a + 2b + c = 3$
 $T_3 = 9a + 3b + c = 6$
 $\therefore 3a + b = 2 \text{ and } 5a + b = 3$
Substitute $b = 2 - 3a \text{ into } 5a + b = 3$
 $\therefore 5a + (2 - 3a) = 3$
 $2a = 1$
 $a = \frac{1}{2}$
 $b = \frac{1}{2}$ and $c = 0$
 $\therefore T_n = \frac{1}{2}n^2 + \frac{1}{2}n$

(b)
$$T_3 = 52 \text{ cm}$$

 $T_7 = 78 \text{ cm}$
 $T_3 = a + 2d = 52$
 $T_7 = a + 6d = 78$
 $4d = 26$ $\therefore d = 6\frac{1}{2}$
 $\therefore a = 39 \text{ cm}$
 $T_{43} = 39 + 42\left(6\frac{1}{2}\right)$
 $T_{43} = 312 \text{ cm}$

(a)
$$f(x) = x^{2} - 6x + 9$$

$$f'(x) = \lim_{h \to 0} \frac{(x+h)^{2} - 6(x+h) + 9 - (x^{2} - 6x + 9)}{h}$$

$$f'(x) = \lim_{h \to 0} \frac{x^{2} + 2xh + h^{2} - 6x - 6h + 9 - x^{2} + 6x - 9}{h}$$

$$f'(x) = \lim_{h \to 0} \frac{2xh + h^{2} - 6h}{h}$$

$$f'(x) = \lim_{h \to 0} \frac{h(2x + h - 6)}{h}$$

$$f'(x) = 2x - 6$$


(2)
$$f'(-3) = 2(-3) - 6 = -12$$

(b)
$$y = \pi x^{-1} + 3x^{\frac{1}{3}}$$

$$\frac{dy}{dx} = -\pi x^{-2} + x^{-\frac{2}{3}}$$

SECTION B

- (a) (1) Domain = $x \in \mathbb{R}$; $x \neq 3$
 - (2) Range = $y \in \mathbb{R}$; $y \neq -3$
 - (3) (i) 5 units
 - (ii) 5 units
- (b) $y = a.b^{x} \text{ substitute } \left(0; \frac{1}{4}\right)$ $\frac{1}{4} = a.b^{0}$ $a = \frac{1}{4}$ $y = \frac{1}{4}b^{x} \text{ substitute } \left(2; \frac{9}{4}\right)$ $\frac{9}{4} = \frac{1}{4}b^{2}$ $b^{2} = 9$



- f: shape intercept domain
- f⁻¹ shape intercept range

- (3) Range = $\left[\frac{1}{4};\infty\right)$
- (4) $f(x) = \frac{1}{4} \cdot 3^{x}$ For f^{-1} : $x = \frac{1}{4} \cdot 3^{y}$; $y \ge 0$ $4x = 3^{y}$ $y = \log_{3}(4x)$ for $x \ge \frac{1}{4}$
- (5) See graph in Question 6 (b) (2) above.

(a)
$$f(x) = x^2 + 6x + (3)^2 + 5 - (3)^2$$

 $f(x) = (x+3)^2 - 4$
 $\therefore \text{T.P.}(-3; -4)$

- (b) (1) $x^2 + 6x + 5 = -x 5$ $x^2 + 7x + 10 = 0$ x = -2 or x = -5 A(-5;0) and B(-2;-3)
 - (2) Horizontal shift: $\therefore -5 < t < -2$
- (c) (1) Length MN = $(-x-5)-(x^2+6x+5)$ Length MN = $-x-5-x^2-6x-5$ Length MN = $-x^2-7x-10$

For max. length: Let $D_x = 0$ -2x-7=0 $x=-\frac{7}{2}$

∴ Max. length MN = $-\left(-\frac{7}{2}\right)^2 - 7\left(-\frac{7}{2}\right) - 10$ ∴ Max. length MN = $\frac{9}{4}$ units i.e. 2,25 units

(2) Vertical shift: $\therefore k > \frac{9}{4}$

(a) (1)
$$\frac{3}{2}$$
; $-\frac{9}{2}$; $\frac{27}{2}$; ...

 $\therefore r = -3$ and series is geometric however, series is not convergent since $r < -1$.

 $\therefore x \neq -\frac{3}{2}$

(2)
$$\frac{x-3}{x+3} = \frac{12-x}{x-3}$$
$$(x-3)^2 = (12-x)(x+3)$$
$$x^2 - 6x + 9 = 12x + 36 - x^2 - 3x$$
$$2x^2 - 15x - 27 = 0$$
$$x = 9 \text{ or } x \neq -\frac{3}{2}$$

(b)
$$S_4 = 7\frac{1}{2}$$
; $S_5 = 15\frac{1}{2}$ and $S_6 = 31\frac{1}{2}$
 $T_5 = S_5 - S_4$
 $T_5 = 8$

$$T_6 = S_6 - S_5$$

 $T_6 = 16$

$$T_5 = ar^4 = 8$$

 $T_6 = ar^5 = 16$

$$\frac{T_6}{T_5} = r = 2$$

$$\therefore a = \frac{1}{2}$$

$$S_n = \frac{\frac{1}{2}(2^n - 1)}{2 - 1}$$

$$= 2^{n - 1} - \frac{1}{2}$$

(a)
$$f(x) = -x^3 + bx^2 + cx - 3$$
$$f(1) = -(1)^3 + b(1)^2 + c(1) - 3 = 4$$
$$b + c = 8$$

$$f'(x) = -3x^{2} + 2bx + c$$

$$f''(x) = -6x + 2b$$

$$f''\left(\frac{1}{2}\right) = -6\left(\frac{1}{2}\right) + 2b = 1$$

$$b = 2$$

$$\therefore c = 6$$

(b) For concave up:
$$f''(x) > 0$$

 $-6x + 4 > 0$
 $x < \frac{2}{3}$

$$\frac{340}{x} - \frac{340}{x+2} = 3 \quad \text{LCD: } x(x+2)$$

$$340(x+2) - 340x = 3x(x+2)$$

$$3x^2 + 6x - 680 = 0$$

$$x = 14,09 \quad \text{or} \quad x \neq -16,09$$
Therefore Time = $\frac{340}{14.09} \approx 24,13 \text{ seconds}$

Alternative

Let original time taken be represented by y.

$$\therefore xy = 340 \text{ ... eq. 1}$$

$$(x+2)(y-3) = 340 \text{ ... eq. 2}$$
From eq. 1 $y = \frac{340}{x}$


$$\therefore (x+2)\left(\frac{340}{x} - 3\right) = 340$$

$$\therefore 3x^2 + 6x - 680 = 0$$

$$\therefore x = 14,09 \text{ or } x \neq -16,09$$

Therefore Time =
$$\frac{340}{14,09} \approx 24,13$$
 seconds

- (a) (1) When x = -2 and x = 0
 - (2)



- (b) $y = \frac{1}{5}x^3 + \frac{3}{4}x + 3$
 - $\frac{dy}{dx} = \frac{3}{15}x^2 + \frac{3}{4}$ substitute x = 0
 - $\frac{dy}{dx} = \frac{3}{4}$
 - Equation of tangent: $y = \frac{3}{4}x + c$ where c = 3
 - Equation of tangent: $y = \frac{3}{4}x + 3$
 - For point of intersection between tangent
 - and line BC, substitute x = 2 into $y = \frac{3}{4}x + 3$

$$\therefore y = 4\frac{1}{2} \therefore Pt\left(2; 4\frac{1}{2}\right)$$

Area of Busi's region = $\frac{1}{2} \left(5 + 3\frac{1}{2} \right) \times 2$

$$=8\frac{1}{2}$$
 units²

Area of Khanya's region = $\frac{1}{2} \left(3 + 4\frac{1}{2} \right) \times 2$

$$=7\frac{1}{2}$$
 units²

Therefore Busi's region is larger.