INLIGTINGSBLAD

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$x = \frac{b}{2a}$$

$$y = \frac{4ac - b^2}{4a}$$

$$a^{x} = b \Leftrightarrow x = \log_{a} b$$
, $a > 0$, $a \ne 1$ en $b > 0$

$$A = P(1 + ni)$$

$$A = P(1 - ni)$$

$$A = P(1+i)^n$$

$$A = P(1+ni)$$
 $A = P(1-ni)$ $A = P(1+i)^n$ $A = P(1-i)^n$

$$i_{\text{eff}} = \left(1 + \frac{i^m}{m}\right)^m - 1$$

$$f'(x) = \lim_{h \to 0} \frac{f(x+h) - f(x)}{h}$$

$$\int \frac{1}{x} dx = \ln(x) + C, \quad x > 0$$

$$\int x^n dx = \frac{x^{n+1}}{n+1} + C \quad , \quad n \neq -1$$

$$\int a^X dx = \frac{a^X}{\ln a} + C \quad , \quad a > 0$$

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

$$M\left(\frac{x_1+x_2}{2};\frac{y_1+y_2}{2}\right)$$

$$y = mx + c$$

$$y-y_1=m(x-x_1)$$

$$m = \frac{y_2 - y_1}{x_2 - x_1}$$

$$m = \tan \theta$$

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$

In ∆ABC:

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc.\cos A$$

oppervlakte \triangle ABC = $\frac{1}{2}ab$.sin C

$$\sin^2 \theta + \cos^2 \theta = 1$$

$$1 + \tan^2 \theta = \sec^2 \theta$$

$$\cot^2 \theta + 1 = \csc^2 \theta$$

 $\pi rad = 180^{\circ}$

Hoeksnelheid = $\omega = 2\pi n = 360^{\circ}n$ waar n = rotasiefrekwensie

Omtreksnelheid = $v = \pi Dn$ waar D = diameter en n = rotasiefrekwensie

 $s = r\theta$ waar $r = \text{radius en } \theta = \text{middelpunthoek in radiale}$

Oppervlakte van sektor = $\frac{rs}{2} = \frac{r^2\theta}{2}$ waar r = radius, s = booglengte en $\theta = \text{middelpunthoek in radiale}$

 $4h^2 - 4dh + x^2 = 0$ waar h = hoogte van segment, d = diameter van sirkel enx = lengte van koord

$$\mathsf{A}_\mathsf{T} = a \bigg(\frac{o_{_1} + o_{_n}}{2} + o_{_2} + o_{_3} + o_{_4} + \ldots + o_{_{n-1}} \bigg) \qquad \text{waar} \qquad a = \mathsf{gelyke} \; \mathsf{dele},$$

$$o_i = i^\mathsf{de} \; \mathsf{ordinaat} \; \mathsf{en}$$

$$n = \mathsf{getal} \; \mathsf{ordinate}$$

OF

$$A_T = a(m_1 + m_2 + m_3 + ... + m_n)$$
 waar $a = gelyke dele, m_1 = \frac{o_1 + o_2}{2}$ en $n = getal ordinate$