

Obesity Medications: Evidence-Based Management

Authors: Venkata Sushma Chamarthi; Sharon F. Daley

Affiliations: 1 Valley Children's Healthcare; 2 Cape Cod Hospital, Hyannis, MA

Last Update: September 2, 2025

Introduction

Obesity has emerged as one of the most significant public health challenges of the 21st century, affecting over 42% of United States adults and contributing to substantial morbidity, mortality, and healthcare costs. [1] The recognition of obesity as a chronic disease requiring medical management has fundamentally shifted treatment paradigms from short-term interventions to comprehensive, long-term treatment approaches. [2] The therapeutic landscape for obesity has been revolutionized by breakthrough medications that achieve weight loss previously only attainable through bariatric surgery. The introduction of glucagon-like peptide-1 receptor agonists and dual incretin receptor agonists has established new standards of care, with some agents producing weight loss of 15% to 25%, compared to 5% to 10% with traditional therapies. [2][3] Additional benefits, including reductions in cardiovascular risk and improvements in obesity-related comorbidities, have accompanied these advances.

Pharmacotherapy is increasingly recognized as a crucial component of evidence-based approaches to comprehensive obesity care. [4] The 2025 updates reflect new comparative effectiveness data, real-world outcomes, and strategies to address persistent health disparities in

medication access. [5] This comprehensive review synthesizes current evidence and provides practical guidance for implementing contemporary obesity pharmacotherapy in clinical practice.

Function

Neurohormonal Dysregulation in Obesity

Obesity fundamentally involves the disruption of complex neurohormonal pathways that regulate energy homeostasis. The hypothalamic-pituitary axis becomes dysregulated, with altered sensitivity to key hormones including leptin, ghrelin, and insulin. [1] Leptin resistance develops as adipose tissue expands, thereby diminishing the hormone's ability to signal satiety to hypothalamic centers that regulate appetite. Concurrently, ghrelin levels remain elevated, promoting persistent hunger signals and food-seeking behaviors.

Palatable foods induce hyperactivation of the mesolimbic dopamine reward pathways, creating neurobiological reinforcement patterns that drive compulsive eating behaviors. Genetic polymorphisms in genes such as MC4R, FTO, and POMC predispose individuals to disrupted satiety mechanisms, increasing their susceptibility to weight gain. These neurohormonal disruptions provide the scientific rationale for pharmacological interventions that target appetite regulation, enhance satiety, and modulate the reward pathway.

Metabolic Dysfunction and Therapeutic Opportunities

The metabolic dysfunction characteristic of obesity creates a pathophysiological environment that perpetuates weight gain and complicates weight loss efforts. Insulin resistance and compensatory hyperinsulinemia develop as adipose tissue expands, promoting further fat storage through enhanced lipogenesis and reduced lipolysis. This metabolic dysregulation makes weight loss increasingly difficult through dietary interventions alone.

Incretin hormone responses become progressively blunted in obesity, with reduced secretion of glucagon-like peptide-1 (GLP-1) and glucose-dependent insulintropic polypeptide (GIP) contributing to impaired glucose homeostasis and diminished meal-related satiety signals. Glucose metabolism becomes dysregulated with the development of peripheral insulin resistance, hepatic glucose overproduction, and ultimately, beta-cell dysfunction. These metabolic alterations provide multiple targets for pharmacological intervention through agents that enhance incretin responses, improve insulin sensitivity, and restore normal glucose and lipid metabolism.

Current Food and Drug Administration-Approved Medications

Tier 1: Superior Efficacy Agents

Agents that have demonstrated the ability to help achieve a 15% to 25% weight loss are categorized as tier 1 pharmacologic therapies.

Tirzepatide

Tirzepatide (brand name Zepbound) represents the most significant advancement in obesity pharmacotherapy, as it is the first dual GLP-1/ GIP receptor agonist approved for chronic weight management. [6] This innovative mechanism targets 2 complementary incretin pathways, resulting in unprecedented weight loss efficacy that surpasses that of all other obesity medications. In the SURMOUNT-1 trial, tirzepatide achieved a dose-dependent weight loss of 15.0%, 19.5%, and 20.9% with doses of 5 mg, 10 mg, and 15 mg, respectively. [6]

The SURMOUNT-5 head-to-head comparison demonstrated tirzepatide's superiority over semaglutide, with 20.2% versus 13.7% weight loss at 72 weeks. [3] Beyond weight loss, tirzepatide showed superior glycemic control, with 91% of participants achieving clinically meaningful weight loss of 5% or more. The SURMOUNT-OSA trial is

ongoing, evaluating the effects of tirzepatide on moderate to severe obstructive sleep apnea. Preliminary data indicate an 18.1% weight loss, accompanied by improvements in sleep apnea parameters.

Semaglutide

Semaglutide (trade name Wegovy) is a once-weekly GLP-1 receptor agonist that has revolutionized obesity treatment by combining substantial weight loss and cardiovascular protection. [7] The STEP clinical trial program demonstrated consistent weight loss of 10.9% to 14.9% across diverse patient populations, with 83.5% of participants achieving clinically meaningful weight loss of 5% or more. [7]

The landmark SELECT cardiovascular outcomes trial established semaglutide as the first obesity medication to demonstrate a reduction in major adverse cardiovascular events, with a 20% reduction in the primary composite endpoint. [7] This cardiovascular benefit occurred independently of weight loss, suggesting direct cardiovascular protective effects. Based on SELECT trial data, the cardiovascular benefits of semaglutide are being evaluated for potential expansion of its Food and Drug Administration (FDA) indication beyond weight management.

In August 2025, the FDA granted accelerated approval to Wegovy (semaglutide injection 2.4 mg) for treating noncirrhotic metabolic dysfunction-associated steatohepatitis (MASH) with moderate to advanced liver fibrosis in adults. This approval represents semaglutide's fourth FDA indication and the second approved therapy for MASH, following resmetirom, previously approved in March 2024. Wegovy's approval utilized the accelerated pathway based on surrogate endpoints, with continued approval contingent upon confirmatory trial results demonstrating long-term clinical benefit.

The approval of Wegovy/semaglutide was supported by interim 72-week results from the Phase 3 ESSENCE trial, which enrolled 800 patients randomized 2:1 to receive semaglutide 2.4 mg or placebo, along with lifestyle modifications. The primary endpoints demonstrated that 63% of patients treated with semaglutide achieved MASH resolution without worsening fibrosis, compared to 34% with placebo ($P < 0.001$). Additionally, 37% of patients treated with semaglutide showed improvement in fibrosis without worsening of MASH, compared to 22%

of patients treated with placebo ($P < 0.001$). Patients experienced a mean weight loss of 10.5% compared to 2.0% with the placebo. The safety profile remained consistent with established data on semaglutide, with gastrointestinal adverse events being the most common. The trial will continue for 240 weeks to assess whether histological improvements translate into a reduction in liver-related morbidity and mortality.

Tier 2: Established Agents

Agents that have demonstrated the ability to help achieve a 5% to 10% weight loss are categorized as tier 2, established pharmacologic therapies.

Liraglutide

Liraglutide (Saxenda) is a daily GLP-1 receptor agonist that pioneered incretin-based obesity therapy. Please refer to StatPearls' companion resource, "Pharmacologic Therapy for Obesity," for additional information. While offering more modest weight loss compared to newer weekly agents, liraglutide maintains clinical relevance through its established safety profile, pediatric indication for adolescents 12 years and older, and flexibility for dose adjustments. The SCALE clinical trial program demonstrated a 5.4% to 8.0% weight loss at 56 weeks, with 63.2% of participants achieving a weight loss of 5% or greater.

Phentermine-Topiramate Extended-Release

The combination therapy phentermine-topiramate extended-release (Qsymia) leverages complementary mechanisms through phentermine's sympathomimetic appetite suppression and topiramate's complex neurological effects on appetite and metabolism. Please refer to StatPearls' companion resource, "Pharmacologic Therapy for Obesity," for additional information. The CONQUER and SEQUEL trials demonstrated 6.6% to 8.6% weight loss at 56 weeks, with 62% to 70% of participants achieving 5% weight loss or more. Long-term safety data from the SEQUEL study confirmed sustained efficacy for up to 2 years, with an acceptable tolerability profile.

Naltrexone-Bupropion Extended Release

Naltrexone-bupropion extended-release (Contrave) targets the neurological aspects of obesity by affecting reward pathways and appetite regulation. The synergistic interaction of naltrexone's opioid receptor blockade and bupropion's monoamine reuptake inhibition modulates food cravings and hedonic eating behaviors. The COR clinical trial program demonstrated a 5% to 6% weight loss at 56 weeks, with 42% to 50% of participants achieving a weight loss of 5% or greater.

Orlistat

Orlistat (Xenical/Alli) remains the only FDA-approved obesity medication available over-the-counter (OTC), providing patients with accessibility without requiring a prescription. Orlistat's unique mechanism of lipase inhibition provides local gastrointestinal effects without systemic absorption, offering a favorable safety profile for patients with multiple comorbidities. Clinical trials demonstrate an additional weight loss of 2 to 3 kg beyond diet and exercise, with 37% of participants achieving a weight loss of 5% or more.

Specialized Medications

Setmelanotide

Setmelanotide (IMCIVREE) represents precision medicine in obesity treatment, specifically targeting rare genetic forms of obesity through melanocortin-4 receptor agonism. This highly specialized medication addresses the underlying pathophysiology of monogenic obesity disorders by restoring hypothalamic appetite regulation pathways disrupted by genetic mutations in the proopiomelanocortin pathway.

Issues of Concern

Comparative Effectiveness Analysis

Head-to-Head Trial Evidence

The SURMOUNT-5 trial provided the first direct head-to-head comparison between the 2 most effective obesity medications currently available. [3] This 72-week, randomized, double-blind study compared tirzepatide 15 mg weekly versus semaglutide 2.4 mg weekly in 751 adults with obesity or overweight with comorbidities. Tirzepatide demonstrated superior efficacy, resulting in 20.2% weight loss compared to 13.7% with semaglutide, representing a clinically meaningful 6.5 percentage point difference.

Beyond mean weight loss, tirzepatide showed superior performance across multiple efficacy endpoints. In the tirzepatide group, 91% of participants achieved a weight loss of 5% or more, compared to 77% in the semaglutide group. Additionally, 57% of participants achieved a weight loss of 20% or more, compared to 28% of those taking semaglutide. These results established tirzepatide as the most effective obesity medication currently available and influenced treatment algorithms favoring dual incretin receptor agonists when clinically appropriate.

Real-World Effectiveness Studies

Real-world evidence studies consistently demonstrate 20% to 30% lower effectiveness than clinical trial results, reflecting the challenges of medication adherence, lifestyle factors, and patient selection in routine clinical practice. [8] An extensive retrospective cohort study of over 40,000 patients found that only 36% of semaglutide users remained on therapy at 1 year, with discontinuation primarily due to gastrointestinal adverse effects and cost barriers. [9]

Persistence rates vary significantly by medication class, with weekly formulations (eg, semaglutide and tirzepatide) showing superior adherence compared to daily injections (eg, liraglutide) or oral medications that require multiple daily doses. Real-world weight loss outcomes with GLP-1 agonists average 8% to 12%, compared to 12% to 20% in clinical trials, highlighting the importance of comprehensive

support systems and realistic expectations for achieving sustained weight loss.

Pediatric Obesity Pharmacotherapy

The treatment of pediatric obesity has been revolutionized by the approval of several adult obesity medications for adolescent use, reflecting the growing recognition that childhood obesity often requires pharmacological intervention alongside lifestyle modifications. [10] Current American Academy of Pediatrics guidelines recommend considering pharmacotherapy for adolescents 12 years or older with obesity (BMI >95th percentile) who have not achieved meaningful weight loss through lifestyle interventions alone.

Note: The 1999 FDA approval date refers to prescription-strength Xenical (120 mg). Over-the-counter Alli (60 mg) was approved later in 2007, but Alli is not FDA-approved for use in pediatric patients.

Key Pediatric Considerations

The following factors should be considered in pediatric patients:

- All medications are adjuncts to lifestyle interventions.
- Growth and development monitoring is required.
- Higher discontinuation rates are observed in children compared to adults, primarily due to gastrointestinal adverse effects.
- Limited long-term safety data should be taken into consideration.
- The American Academy of Pediatrics 2023 guidelines recommend GLP-1 agonists as preferred agents.

The STEP TEENS trial demonstrated the superior efficacy of semaglutide in adolescents, with a 16.1% reduction in BMI compared to a 0.6% increase in those receiving the placebo. [10] Notably, 77% of adolescents achieved a BMI reduction of 5% or more, establishing semaglutide as the preferred first-line pharmacotherapy for adolescent obesity when lifestyle interventions are insufficient.

Emerging Therapies and Pipeline Development

Phase 3 Medications

The obesity medication pipeline represents one of the most robust therapeutic development programs in modern medicine, with over 157 clinical-stage assets spanning more than 60 distinct mechanisms of action. Several breakthrough therapies are expected to receive FDA approval by 2026, potentially offering even greater efficacy than current standard-of-care medications.

Retatrutide

Retatrutide (manufactured by Eli Lilly) represents the next evolution in incretin-based therapy as a triple GLP-1/GIP/glucagon receptor agonist currently in phase 3 trials. Phase 2 data demonstrated an unprecedented 24.2% weight loss at 48 weeks with the highest dose, surpassing the weight loss achieved with all currently available medications. The triple mechanism targets complementary pathways in glucose homeostasis, appetite regulation, and energy expenditure, providing synergistic effects that translate into superior clinical outcomes.

CagriSema

CagriSema (manufactured by Novo Nordisk) combines semaglutide with cagrilintide, a novel amylin analog, targeting both incretin and amylin receptor pathways. Phase 3 data showed 20.4% weight loss with potentially improved gastrointestinal tolerability compared to semaglutide alone. The combination approach leverages the complementary mechanisms of GLP-1 and amylin receptor activation, potentially offering enhanced efficacy with reduced adverse effects.

Oral Semaglutide

Oral semaglutide 50 mg represents a significant advancement in patient convenience, offering the efficacy of injectable semaglutide in an oral formulation. Phase 3 data demonstrated a 17.4% weight loss, approaching the effectiveness of injectable formulations while

eliminating the need for injections. This development could significantly enhance access to and adherence with GLP-1-based therapy.

Survodutide

Survodutide (manufactured by Boehringer Ingelheim) is a novel dual GLP-1/glucagon receptor agonist that achieved 18.7% weight loss in phase 2 trials. Adding glucagon receptor agonism to GLP-1 activity enhances metabolic rate and energy expenditure, providing a complementary mechanism to appetite suppression, thereby achieving superior weight loss outcomes. Survodutide has demonstrated direct hepatic effects, including enhanced fat mobilization, lipolysis, increased energy expenditure, and a reduction in hepatic steatosis and fibrotic processes, in clinical trials.

Novel Mechanisms in Development

Amylin Receptor Agonists

Amylin receptor agonists represent a promising class of medications that complement incretin-based therapies by targeting distinct but related pathways involved in appetite regulation and glucose homeostasis. Petrelintide (ZP8396) from Zealand Pharma and Roche has shown promising early-phase data with a clean safety profile and potential for combination therapy with GLP-1 agonists. Eloralintide from Eli Lilly demonstrated 11.3% weight loss when added to tirzepatide therapy, suggesting potential for enhanced efficacy through combination approaches.

Monthly Formulations

Monthly formulations address the critical issue of medication adherence through extended dosing intervals. MariTide from Amgen utilizes a novel peptide-antibody fusion technology, providing monthly dosing convenience while maintaining therapeutic efficacy. Early clinical data suggest a potential for substantial weight loss and improved patient compliance compared to weekly injection regimens.

Muscle Preservation Agents

Muscle preservation agents represent an emerging therapeutic approach addressing the challenge of lean muscle mass loss during weight reduction. Bimagrumab, a myostatin inhibitor, has shown promise in preserving muscle mass during caloric restriction, potentially improving metabolic outcomes and long-term weight maintenance.

Off-Label Medications and Alternative Approaches

Several medications have demonstrated efficacy for weight management, although they do not have specific FDA approval for the treatment of obesity. These agents may provide valuable alternatives for patients who cannot tolerate or access approved obesity medications; however, their use requires careful consideration of the risk-benefit profiles and comprehensive informed consent.

Short-Acting Sympathomimetic Amines

Short-acting sympathomimetic amines represent a potential option in obesity management, though their use demands careful clinical consideration. These agents stimulate the central nervous system to suppress appetite and increase energy expenditure, facilitating modest short-term weight loss. Their role remains limited, as data on long-term outcomes are insufficient to establish sustained efficacy.

Several common contraindications restrict the safe use of short-acting sympathomimetic amines. Patients with cardiovascular disease face significant risks due to increased heart rate and blood pressure. Hyperthyroidism and glaucoma also preclude their use, as sympathomimetic stimulation can worsen these conditions. Additional contraindications include agitated states, where central nervous system activation may intensify symptoms, and concurrent therapy with monoamine oxidase inhibitors, which creates the potential for severe drug interactions.

Close monitoring remains essential throughout treatment. Clinicians

must regularly assess blood pressure, heart rate, weight changes, mood stability, and the potential for abuse or misuse. When prescribed and closely supervised appropriately, these medications can help individuals achieve a weight loss of 3% to 8%. However, the benefits must be weighed against safety concerns and the limited evidence of long-term effectiveness.

Diabetes Medications with Weight Loss Benefits

Several diabetes medications provide weight loss benefits, making them valuable options for patients with concurrent diabetes and obesity or as off-label options when approved obesity medications are contraindicated or inaccessible.

Dosing Adjustments and Special Populations

Renal Impairment Considerations

In patients with severe renal impairment, certain obesity medications may require dose adjustments or may be contraindicated.

Hepatic Impairment Considerations

In patients with severe hepatic impairment, certain obesity medications may require dose adjustments or may be contraindicated.

Clinical Significance

Pharmacologic Management Approach for Obesity

Appropriate patient selection is essential for optimal outcomes with obesity pharmacotherapy. Determining which patients would best benefit from pharmacological treatment requires a systematic patient evaluation

that incorporates medical history, screening for contraindications, and setting realistic expectations. Evidence-based guidelines emphasize the importance of comprehensive assessment, including calculation of BMI, evaluation of comorbidities, previous attempts at weight loss, and psychosocial factors that may influence treatment success. [4]

Candidate Selection Criteria

Patients with the following characteristics indicate appropriate candidates for pharmacologic treatment of obesity:

- BMI >30 kg/m² or BMI ≥ 27 kg/m² with weight-related comorbidities
- Failure to achieve meaningful weight loss through lifestyle interventions alone
- Absence of contraindications to specific medications
- Realistic expectations and commitment to long-term treatment
- Adequate social support and access to ongoing medical care

Contraindication Screening

Pharmacologic treatment of obesity is contraindicated in individuals with the following factors:

- Personal or family history of medullary thyroid carcinoma or multiple endocrine neoplasia type 2 syndrome (specifically GLP-1 agonists)
- Pregnancy or breastfeeding (most obesity medications are contraindicated)
- Active eating disorders or substance abuse history
- Unstable cardiovascular disease or uncontrolled hypertension
- Severe psychiatric illness without adequate treatment

Medication Selection

Current evidence supports initiating treatment with the most effective medication unless specific contraindications exist. The superior efficacy of tirzepatide and semaglutide compared to other approved medications

justifies their use as first-line therapies when clinically appropriate and accessible. [6][3] Tirzepatide offers the highest efficacy, achieving an average weight loss of 20.2% through its dual mechanism of action, which involves the incretin effect. Semaglutide offers substantial weight reduction, along with additional proven cardiovascular benefits and a well-established safety profile. Alternative agents may be considered when GLP-1 receptor agonists are not feasible or unavailable.

Treatment Escalation

Treatment escalation strategies guide decision-making when initial therapy does not achieve expected outcomes. Clinicians should evaluate patient response at 12 weeks and discontinue therapy if the patient has not achieved a weight loss of at least 5%. Before switching medications, dose escalation may be attempted to enhance efficacy. If adverse effects remain intolerable, transitioning to a therapy with a different mechanism becomes appropriate. Combination pharmacotherapy may also be considered for suitable candidates, particularly when monotherapy fails to achieve adequate weight reduction.

Pharmacotherapy Monitoring Protocols

A comprehensive monitoring schedule supports the safe and effective use of pharmacologic therapy for obesity. Initial baseline clinical monitoring includes weight and BMI measurement, assessment of blood pressure and heart rate, laboratory testing, lifestyle counseling, and evaluation of comorbidities. Weight, BMI, blood pressure, and heart rate should be reassessed at months 1, 3, and 6. Ongoing blood pressure and heart rate monitoring should be performed at every visit, and every 3 months for weight and BMI assessment.

Baseline laboratory evaluation should include HbA1c (for patients with diabetes), lipid panel, liver and renal function tests, thyroid function when indicated, and lipase for those receiving GLP-1 receptor agonists. These tests should be repeated at 3 and 6 months, and every 6 months thereafter. Clinicians should inquire about adverse effects and medication adherence, and provide lifestyle counseling at the initial visit

and all subsequent appointments. Comorbidities should also be assessed at the initial visit, and reassessment is recommended at 3 months, 6 months, and then every 6 months thereafter.

Monitoring for GLP-1 Agonists and Sympathomimetics

Specific monitoring strategies vary by medication class. For GLP-1 receptor agonists, baseline evaluation includes assessment of lipase, renal function, and thyroid history, with ongoing monitoring of weight response, gastrointestinal symptoms, and warning signs of pancreatitis or gallbladder disease. Semaglutide requires additional monitoring for diabetic retinopathy progression.

For sympathomimetic agents, baseline evaluation includes cardiovascular risk assessment, blood pressure, and heart rate, with continued monitoring of blood pressure, heart rate, mood changes, and sleep patterns. Qsymia requires monthly pregnancy testing along with evaluation for abuse potential. This structured approach enhances patient safety, optimizes therapeutic outcomes, and supports adherence to long-term treatment.

Adverse Effect Management

Effective management of medication-related adverse effects is crucial for optimizing treatment adherence and long-term success in obesity pharmacotherapy. Healthcare practitioners must implement proactive monitoring strategies and evidence-based interventions to minimize adverse effects while maintaining therapeutic efficacy.

General Principles for Adverse Effect Management

General principles for adverse effect management emphasize the use of proactive, collaborative, and systematic strategies to ensure patient safety and optimize treatment outcomes. Open communication plays a central role, with clinicians encouraging patients to report adverse effects promptly and providing clear guidance on which symptoms require urgent medical attention versus those that can be managed

through supportive care and scheduled follow-up. Ongoing risk-benefit assessment remains essential throughout therapy. Clinicians must evaluate the severity and functional impact of adverse effects related to weight loss efficacy, recognizing that some patients may opt to continue treatment when the clinical benefits outweigh the tolerable side effects. Complex or severe reactions often necessitate an interprofessional approach, involving specialists, eg, cardiologists, psychiatrists, gastroenterologists, or endocrinologists, to provide targeted expertise.

Accurate documentation further strengthens clinical management. Detailed records of adverse effects, interventions, and patient responses help guide future care and support continuity across healthcare practitioners. Serious adverse events should be reported to the FDA's MedWatch program, with contributions to postmarketing surveillance databases encouraged to expand collective safety knowledge and advance best practices in obesity pharmacotherapy.

Gastrointestinal Adverse Effects

Gastrointestinal adverse events represent the most common cause of treatment discontinuation with GLP-1 receptor agonists, affecting 25% to 79% of patients depending on the specific agent and dose. These effects typically occur during dose escalation phases and often improve with time and the implementation of appropriate management strategies.

Management strategies for gastrointestinal adverse effects include:

- **Titration optimization:** Implement the slowest recommended titration schedule to allow physiological adaptation to delayed gastric emptying. Consider extending titration periods beyond standard protocols for patients experiencing significant gastrointestinal symptoms, as slower dose escalation often improves long-term tolerability without compromising efficacy.
- **Dietary modifications:** Counsel patients to consume smaller, more frequent meals (4 to 6 times daily) with reduced fat content (<30% of calories from fat). High-fat meals exacerbate delayed gastric emptying and increase nausea severity. Avoid carbonated beverages, spicy foods, and large meal portions, as these can trigger symptoms.
- **Dose management:** If severe symptoms persist beyond 2 to 3

weeks at the current dose, consider temporarily reducing the dose by 1 titration step. Maintain the reduced dose for 2 to 4 weeks before attempting re-escalation. Some patients may require maintenance at sub-maximal doses to balance efficacy with tolerability.

- Pharmacological interventions: For persistent nausea unresponsive to dietary modifications, prescribe antiemetics (eg, ondansetron 4-8 mg as needed or metoclopramide 10 mg before meals). Proton pump inhibitors may benefit patients experiencing symptoms of acid reflux or gastroesophageal reflux disease (GERD).
- Hydration and electrolyte management: Ensure adequate hydration (8 to 10 glasses of water daily) and monitor electrolyte balance, particularly in patients experiencing vomiting. Consider oral rehydration solutions for patients with significant fluid losses and monitor for signs of dehydration, including dizziness, fatigue, and decreased urine output.

Cardiovascular Adverse Effects

Sympathomimetic agents, including phentermine and phentermine-topiramate combinations, necessitate vigilant cardiovascular monitoring due to their potential to increase heart rate, blood pressure, and cardiac workload. These effects are dose-dependent and may be more pronounced in patients with underlying cardiovascular disease or other risk factors.

Management strategies for cardiovascular adverse effects include:

- Baseline and ongoing monitoring: Establish baseline blood pressure and heart rate measurements before treatment initiation. Monitor vital signs at every visit during the first 3 months, then monthly thereafter. Consider 24-hour ambulatory blood pressure monitoring for patients with borderline hypertension or white-coat syndrome.
- Hypertension management: Implement dose reduction or treatment discontinuation if systolic blood pressure consistently exceeds 140 mmHg or diastolic pressure exceeds 90 mmHg, despite optimal antihypertensive therapy. Consider adding or

adjusting antihypertensive medications in consultation with cardiology when appropriate.

- **Cardiac symptom evaluation:** Conduct immediate cardiac evaluation, including electrocardiogram, and consider stress testing or echocardiography if patients develop chest pain, palpitations, shortness of breath, or exercise intolerance. If symptoms develop that are concerning, discontinue treatment and consult a cardiologist for further evaluation and clearance.
- **Drug interaction surveillance:** Monitor carefully for interactions with antihypertensive medications, particularly beta-blockers, angiotensin-converting enzyme inhibitors, and diuretics. Assess for potential interactions with psychiatric medications, particularly monoamine oxidase inhibitors, tricyclic antidepressants, and other sympathomimetic agents that may compound cardiovascular effects.

Neuropsychiatric Adverse Effects

Topiramate-containing combinations, particularly Qsymia, can produce significant neuropsychiatric adverse effects that may impact cognitive function, mood stability, and overall quality of life. These effects are often dose-dependent and may be reversible with a reduction in dose or discontinuation of the medication.

Management strategies for neuropsychiatric adverse effects include:

- **Cognitive function assessment:** Conduct systematic evaluation of cognitive function, including attention, memory, word-finding ability, and processing speed, at baseline and every 3 months. When clinically indicated, use standardized tools, eg, the Montreal Cognitive Assessment or Mini-Mental State Examination. Monitor for reports of difficulty concentrating, confusion, or "brain fog."
- **Mood and psychiatric monitoring:** Implement regular screening for depression using validated tools such as the Patient Health Questionnaire-9, and assess for anxiety, irritability, and suicidal ideation at each visit. Establish collaborative care with mental health professionals for patients with pre-existing psychiatric conditions or those developing new symptoms during treatment.
- **Dose optimization strategies:** If cognitive impairment or mood

changes develop, consider reducing the dose to the lowest effective level. Some patients may benefit from alternate-day dosing or extended-release formulations to minimize peak concentration effects. Discontinuation may be necessary if symptoms significantly impact daily functioning or quality of life.

- Patient and family education: Provide comprehensive education to patients and family members about potential neuropsychiatric effects, emphasizing the importance of reporting subtle changes in thinking, mood, or behavior. Establish clear protocols for emergency contact if suicidal thoughts or severe mood changes develop.
- Metabolic monitoring: Monitor for metabolic acidosis by periodically measuring serum bicarbonate levels, particularly in patients with risk factors such as kidney disease, severe diarrhea, or concurrent use of carbonic anhydrase inhibitors. Assess for symptoms, including fatigue, anorexia, and hyperventilation, that may indicate acidosis development.

Other Issues

Health Equity Considerations

Significant racial and socioeconomic disparities exist in obesity medication prescribing, with 85% of prescriptions going to White patients despite higher obesity prevalence in minority populations. [5] These disparities reflect complex interactions among healthcare access, clinician bias, insurance coverage, and cultural factors that the healthcare system must address to achieve equitable care for obesity.

Strategies for improving health equity include:

- Cultural competency training for clinicians
- Community health worker programs in underserved areas
- Multilingual patient education materials
- Clinician bias recognition and mitigation training
- Targeted outreach to minority-serving healthcare organizations

Insurance Coverage and Access Barriers

The high cost of obesity medications (\$1000-\$1500 monthly) creates substantial barriers to access, with significant variation in insurance coverage across different plans and patient populations. Medicare generally does not cover obesity medications as standalone therapy; however, coverage may be available when the drugs are prescribed for diabetes or other approved indications. Medicaid coverage varies significantly by state, and ongoing policy efforts aim to expand coverage.

Prior Authorization Approval

Insurance providers often require prior authorizations for obesity medications to obtain patient coverage. Implementing the following strategies may increase the likelihood of obtaining approval for these medications:

- Document comprehensive lifestyle intervention attempts.
- Emphasize weight-related comorbidities and potential cost savings.
- Include BMI progression over time, demonstrating medical necessity.
- Highlight cardiovascular risk factors and potential for risk reduction.
- Provide peer-reviewed literature supporting specific medication selection.

Appeals Process Optimization

Clinicians may appeal denials of obesity medications by the insurance provider on behalf of the patient. The following strategies can optimize this process:

- Escalate denials with additional clinical documentation.
- Include specialist (endocrinologist, obesity medicine) recommendations.
- Document quality of life impacts and functional improvements.
- Consider an independent medical review when available.
- Utilize patient assistance programs as bridge therapy.

Cost-Effectiveness Considerations

Economic analyses demonstrate favorable cost-effectiveness ratios for obesity medications, considering long-term healthcare cost savings from comorbidities and improvements in cardiovascular risk reduction. Semaglutide's cardiovascular benefits offer robust economic justification, with estimated cost savings based on modeling from the SELECT trial and health economic analyses.

The economic benefits of these medications include:

- Diabetes prevention and remission, reducing long-term costs
- Cardiovascular event reduction, decreasing acute care utilization
- Sleep apnea improvement, reducing the need for continuous positive airway pressure (CPAP), and surgical costs
- Joint preservation, reducing orthopedic interventions
- Mental health improvements, reducing psychiatric care needs

Future Directions and Research Priorities

The future of obesity pharmacotherapy lies in personalized treatment selection based on individual patient characteristics, genetic factors, and predictive biomarkers. Research into the pharmacogenomics of obesity medications may enable precision dosing and selection to optimize efficacy while minimizing adverse effects.

Emerging precision medicine strategies include:

- Genetic testing for monogenic obesity forms (setmelanotide candidates)
- Biomarker development for treatment response prediction
- Phenotype-based treatment algorithms
- Personalized dosing based on pharmacokinetic factors
- Gut microbiome analysis for medication selection (future research area)

Combination Therapy Development

The success of existing combination therapies (phentermine-topiramate,

naltrexone-bupropion) provides a roadmap for developing more effective combinations targeting complementary pathways. Future combinations may include GLP-1 agonists with amylin analogs, muscle preservation agents, or novel mechanisms to achieve superior efficacy profiles.

Combination therapeutic approaches that have shown promise include:

- GLP-1 and amylin agonists (CagriSema model)
- Incretin and muscle preservation agents
- Appetite suppression and metabolic enhancement
- Weight loss and muscle preservation combinations
- Targeted genetic therapy and pharmacotherapy
- Triple agonist therapies targeting GLP-1/GIP/glucagon pathways (currently in phase 2-3 trials)

Technological Integration

Digital health technologies offer opportunities to enhance medication adherence, optimize dosing, and provide personalized support for patients receiving obesity pharmacotherapy. Integration of continuous glucose monitoring, activity tracking, and telemedicine platforms may improve outcomes and reduce healthcare costs.

Digital health innovations include:

- Smartphone apps for medication reminders and adverse effect tracking
- Continuous glucose monitoring for GLP-1 agonist optimization
- Artificial intelligence-powered treatment algorithms for personalized care
- Telemedicine platforms for rural and underserved populations
- Wearable devices for real-time metabolic monitoring

Clinical Takeaways

The landscape of obesity pharmacotherapy has been fundamentally transformed by breakthrough medications that achieve unprecedented weight loss efficacy, comparable to the outcomes of bariatric surgery. Tirzepatide and semaglutide represent paradigm-shifting therapies that

offer not only substantial weight loss but also significant improvements in cardiovascular health and obesity-related comorbidities. [6][3] The robust pipeline of emerging therapies promises even greater efficacy, with retatrutide demonstrating potential for 24% weight loss and novel combination approaches targeting multiple pathways simultaneously. These advances position pharmacotherapy as a cornerstone of comprehensive obesity management, necessitating that healthcare systems adapt their treatment algorithms, monitoring protocols, and access strategies accordingly.

However, realizing the full potential of these therapeutic advances requires addressing persistent barriers, including high medication costs, limited insurance coverage, and significant health disparities in access to care. [3] Clinicians must advocate for equitable access while developing systematic approaches to patient selection, monitoring, and long-term management. The future of obesity medicine lies in precision approaches combining optimal pharmacotherapy with comprehensive lifestyle interventions, advanced monitoring technologies, and personalized treatment algorithms.

As the field evolves rapidly, staying current with emerging evidence and focusing on health equity will be essential for optimizing patient outcomes and effectively addressing the global obesity epidemic. Clinicians should approach obesity pharmacotherapy as chronic disease management requiring a long-term commitment, comprehensive monitoring, and individualized treatment approaches. The exceptional efficacy of current and emerging therapies offers unprecedented opportunities to transform patient outcomes, provided these powerful tools are implemented thoughtfully and equitably across all patient populations.

Additional clinical factors that clinicians should bear in mind when employing obesity medications include:

- Tirzepatide and semaglutide represent first-line therapies for most patients, offering superior efficacy and established safety profiles.
- Early response assessment at 12 weeks guides treatment continuation and optimization strategies.
- Comprehensive monitoring protocols ensure safety while optimizing therapeutic outcomes.

- Health equity considerations must be integrated into all aspects of obesity pharmacotherapy.
- Pipeline therapies offer promise for even greater efficacy and improved patient convenience.
- Cost-effectiveness supporting insurance coverage requires emphasis on long-term health benefits and improvements in comorbidity.

Enhancing Healthcare Team Outcomes

Obesity is recognized as a chronic disease affecting a significant portion of United States adults and driving substantial morbidity, mortality, and healthcare costs. Advances in pharmacotherapy, particularly with GLP-1 receptor agonists and dual incretin receptor agonists, have transformed the treatment landscape, achieving weight loss outcomes once reserved for bariatric surgery. Current treatment guidelines emphasize pharmacotherapy as a cornerstone of long-term management, incorporating evidence-based strategies to address metabolic dysfunction, reduce cardiovascular risk, and improve comorbidities. Emerging therapies, pediatric treatment considerations, and efforts to reduce disparities in access highlight the importance of a comprehensive, patient-centered approach.

Delivering optimal care for obesity requires collaboration across multiple disciplines. Physicians and advanced practitioners guide diagnosis, patient selection, and long-term monitoring, while nurses play key roles in counseling, adherence support, and managing adverse effects. Pharmacists ensure safe prescribing, monitor for interactions, and assist with access barriers. Effective interprofessional communication and coordinated care strategies strengthen patient engagement, improve safety, and enhance outcomes. By aligning expertise, healthcare teams can deliver equitable, personalized obesity management that promotes sustainable weight loss and long-term health.

References

1. Chamarthi VS, Daley SF. StatPearls [Internet]. StatPearls Publishing; Treasure Island (FL): Aug 29, 2025. Comprehensive Assessment and Diagnosis of Metabolic and Biomechanical Complications in Obesity. [PubMed: 29083734]
2. Rodrigues Silva Sombra L, Anastasopoulou C. StatPearls [Internet]. StatPearls Publishing; Treasure Island (FL): Feb 12, 2024. Pharmacologic Therapy for Obesity (Archived) [PubMed: 32965940]
3. Aronne LJ, Horn DB, le Roux CW, Ho W, Falcon BL, Gomez Valderas E, Das S, Lee CJ, Glass LC, Senyucel C, Dunn JP; SURMOUNT-5 Trial Investigators. Tirzepatide as Compared with Semaglutide for the Treatment of Obesity. *N Engl J Med*. 2025 Jul 03;393(1):26-36. [PubMed: 40353578]
4. Gudzone KA, Wickham EP, Schmidt SL, Stanford FC. Physicians certified by the American Board of Obesity Medicine provide evidence-based care. *Clin Obes*. 2021 Feb;11(1):e12407. [PMC free article: PMC9999726] [PubMed: 33280270]
5. Isoyama Y, Nose-Ogura S, Ijitsu MJ, Kruse JGS, Nagai N, Kayaba M, Ogata H, Mangalam M, Kiyono K. Age- and height-dependent bias of underweight and overweight assessment standards for children and adolescents. *Front Public Health*. 2024;12:1379897. [PMC free article: PMC11076850] [PubMed: 38721543]
6. Jamshed H, Steger FL, Bryan DR, Richman JS, Warriner AH, Hanick CJ, Martin CK, Salvy SJ, Peterson CM. Effectiveness of Early Time-Restricted Eating for Weight Loss, Fat Loss, and Cardiometabolic Health in Adults With Obesity: A Randomized Clinical Trial. *JAMA Intern Med*. 2022 Sep 01;182(9):953-962. [PMC free article: PMC9361187] [PubMed: 35939311]
7. Wilding JPH, Batterham RL, Calanna S, Davies M, Van Gaal LF, Lingvay I, McGowan BM, Rosenstock J, Tran MTD, Wadden TA, Wharton S, Yokote K, Zeuthen N, Kushner RF; STEP 1 Study Group. Once-Weekly Semaglutide in Adults with Overweight or Obesity. *N Engl J Med*. 2021 Mar 18;384(11):989-1002. [PubMed: 33567185]

8. Chakhtoura M, Haber R, Ghezzawi M, Rhayem C, Tcheroyan R, Mantzoros CS. Pharmacotherapy of obesity: an update on the available medications and drugs under investigation. *EClinicalMedicine*. 2023 Apr;58:101882. [PMC free article: PMC10041469] [PubMed: 36992862]
9. Ghush W, De la Rosa A, Sacoto D, Cifuentes L, Campos A, Feris F, Hurtado MD, Acosta A. Weight Loss Outcomes Associated With Semaglutide Treatment for Patients With Overweight or Obesity. *JAMA Netw Open*. 2022 Sep 01;5(9):e2231982. [PMC free article: PMC9486455] [PubMed: 36121652]
10. Wadden TA, Bailey TS, Billings LK, Davies M, Frias JP, Koroleva A, Lingvay I, O'Neil PM, Rubino DM, Skovgaard D, Wallenstein SOR, Garvey WT; STEP 3 Investigators. Effect of Subcutaneous Semaglutide vs Placebo as an Adjunct to Intensive Behavioral Therapy on Body Weight in Adults With Overweight or Obesity: The STEP 3 Randomized Clinical Trial. *JAMA*. 2021 Apr 13;325(14):1403-1413. [PMC free article: PMC7905697] [PubMed: 33625476]