

Coccidiosis:



Coccidiosis is a common parasitic disease in poultry, caused by protozoan parasites of the genus *Eimeria*. These parasites invade the intestinal lining of chickens, leading to inflammation and damage to the intestinal tissues. The disease is particularly prevalent in young birds, though it can affect hens of any age, especially in overcrowded or unsanitary conditions.

Symptoms of Coccidiosis:

- Diarrhea (often with blood)
- Weight loss
- Decreased feed and water consumption
- Ruffled feathers
- Lethargy
- Pale comb and wattles
- Droopiness
- Reduced egg production (in laying hens)

Transmission:

Coccidiosis is primarily spread through the ingestion of infected fecal material. The oocysts (eggs of the parasite) are shed in the droppings of infected birds and can survive for a long time in the environment, particularly in moist and warm conditions.

Prevention of Coccidiosis:

1. Good Hygiene Practices:

- Regularly clean and disinfect the poultry house and equipment.
- Remove wet and soiled bedding frequently to reduce the risk of infection.
- Provide fresh and clean water daily.

2. Proper Management of Litter:

- Maintain dry litter conditions, as the oocysts thrive in moist environments.
- Regularly turn and replace the litter to keep it dry and clean.

3. Vaccination:

- Vaccines are available for some species of *Eimeria*. Vaccinating young chicks can help build immunity against coccidiosis.

4. Avoid Overcrowding:

- Ensure adequate space for each bird to reduce stress and lower the risk of spreading the disease.

Salmonella:



Salmonella is a bacterial infection that affects chickens and other poultry, caused by different strains of *Salmonella* bacteria. These bacteria can be present in the intestines of birds and are a significant concern due to their potential to cause foodborne illness in humans through contaminated eggs or meat.

Symptoms of Salmonella in Hens:

- Diarrhea (often watery and yellowish)
- Loss of appetite
- Lethargy
- Dehydration
- Drop in egg production
- Pale comb and wattles
- Weight loss
- Death (in severe cases)

Transmission:

- Salmonella is primarily spread through the fecal-oral route.
- Contaminated feed or water
- Infected equipment or housing
- Direct contact with infected birds
- Wild birds, rodents, or insects that carry the bacteria
- Vertical transmission (from hen to egg), leading to infected chicks hatching from contaminated eggs

Prevention of Salmonella in Hens:

1. Good Hygiene and Sanitation Practices:

- Regularly clean and disinfect poultry houses, feeders, drinkers, and all equipment.
- Ensure that the housing environment is dry and free of fecal contamination.

2. Safe Feed and Water:

- Provide clean, uncontaminated feed and water. Store feed in a dry, rodent-proof environment.
- Avoid feeding birds with raw or undercooked kitchen scraps that may harbor bacteria.


3. Vaccination:

- Vaccines are available for some strains of Salmonella, and vaccinating hens can reduce the incidence of the disease and the risk of contamination in eggs.

4. Proper Egg Handling:

- Collect eggs frequently and store them in a clean, cool environment.

New Castle Disease:



Newcastle Disease is a highly contagious viral disease affecting chickens and other poultry, caused by the Newcastle Disease Virus (NDV). The virus belongs to the family *Paramyxoviridae* and can cause severe economic losses in poultry due to its high mortality rate. The severity of the disease can vary depending on the strain of the virus, the species of bird affected, and the bird's immunity.

Symptoms of Newcastle Disease:

- **Respiratory Symptoms:** Coughing, Sneezing, Nasal discharge, Labored breathing, Swelling around the eyes
- **Digestive Symptoms:** Diarrhea (greenish in color), Reduced appetite, Weight loss.
- **Nervous Symptoms:** Tremors, Twisted neck (torticollis), Paralysis, Loss of balance, Circling or falling over.
- **General Symptoms:** Drop in egg production (soft-shelled or misshapen eggs), Sudden death, Lethargy

Transmission:

- Newcastle Disease is primarily spread through direct contact with infected birds or their secretions, including respiratory droplets, feces, and eggs.
- The virus can also be transmitted indirectly through contaminated feed, water, equipment, clothing, and footwear.

Prevention of Newcastle Disease:

1. Vaccination:

- Vaccination is the most effective way to prevent Newcastle Disease. Vaccines are available in different forms (live, killed, and recombinant), and their use depends on the local disease situation and the age of the birds.
- Vaccinate birds according to a schedule recommended by a veterinarian, which usually involves multiple doses throughout the birds' lives.

2. Strict Biosecurity Measures:

- Implement stringent biosecurity protocols to prevent the introduction and spread of the virus.
- Limit access to the poultry farm, and restrict visitors or require them to follow strict hygiene procedures.

3. Proper Hygiene and Sanitation:

- Regularly clean and disinfect poultry houses, equipment, feeders, and drinkers.
- Dispose of dead birds, litter, and waste properly to prevent the spread of the virus.

4. Health Monitoring and Reporting:

- Regularly monitor the health of the flock and report any signs of Newcastle Disease to a veterinarian or local animal health authority.

Healthy:

- Your hen is in great health! She is showing no signs of illness or distress and is thriving in her environment.
- Her feathers are smooth and shiny, her appetite is good, and she's active and alert.
- Continue to maintain proper hygiene, provide a balanced diet, and ensure she has a clean, safe living space.
- Regularly monitor her health and keep up with vaccinations to ensure she stays happy and healthy.



Tomato__Tomato_mosaic_virus:



Description:

The American Bollworm is a moth in its adult stage, but the larvae (caterpillars) are the primary concern for crops. These larvae are highly destructive, feeding on the cotton bolls, flowers, and leaves.

Life Cycle:

The life cycle of the American Bollworm includes egg, larval, pupal, and adult stages. The larvae are the most damaging stage, as they bore into cotton bolls and consume the seeds and fibers, leading to significant yield loss.

Damage:

The larvae bore into cotton bolls, which can lead to reduced cotton quality and yield. The damaged bolls may also become more susceptible to fungal infections.

Symptoms:

- The larva of American bollworm feeds on all parts of plants including leaves, squares, flowers and small bolls.
- In the early stage, larva feeds on the tender leaves.
- Whereas in the later stage, they feed on the internal content of squares or bolls by thrusting their head inside while leaving the rest of the body outside
- The presence of bore holes with frass material excreted by larva can be seen on the infested bolls.
- As a result of damage, the affected squares and bolls fall off from the plants.

Prevention and Control Measures:

1. Cultural Practices:

- **Crop Rotation:** Avoid planting cotton after another host crop (like tomatoes or corn) to reduce the pest's population.
- **Intercropping:** Planting non-host crops alongside cotton can reduce the pest's ability to spread.

2 Monitoring and Early Detection:

- **Pheromone Traps:** Use pheromone traps to monitor adult moth activity and predict potential infestations.
- **Scouting:** Regularly inspect fields for the presence of eggs and larvae to catch infestations early.

3. Biological Control:

- **Natural Predators:** Encourage the presence of natural predators like lady beetles, lacewings, and parasitic wasps, which can help control the Bollworm population.



Squash_Powdery_mildew:

Description:

Armyworms are caterpillars in the larval stage of moths. They are named for their behavior of moving in large groups ("armies") to consume crops. They are highly destructive, feeding on leaves, stems, and fruits.

Life Cycle:

The life cycle includes egg, larval, pupal, and adult stages. The larval stage is the most damaging, as the caterpillars feed aggressively on crops.

Damage:

Armyworms defoliate plants, leading to reduced photosynthesis and crop yield. In severe infestations, they can completely strip a field of vegetation, resulting in total crop loss.

Symptoms:

- Larvae cut the seedlings in large scale.
- chewing/leaf scalloping along the leaf margins.

Prevention and Control Measures:

1. Cultural Practices:

- **Early Planting:** Plant crops early in the season to avoid peak Armyworm infestations, as younger crops are more vulnerable.
- **Crop Rotation:** Rotate crops with non-host plants to break the life cycle of the pest.

2. Biological Control:

- **Natural Predators and Parasitoids:** Encourage beneficial insects like parasitic wasps, lady beetles, and birds that prey on Armyworms.
- **Bacillus thuringiensis (Bt):** The use of Bt, either through genetically modified crops or as a spray, can effectively control Armyworm larvae.

3. Chemical Control:

- **Insecticides:** Use insecticides such as pyrethroids, organophosphates, and carbamates for controlling large infestations. However, chemical control should be used judiciously to avoid resistance and harm to non-target organisms.
- **Insect Growth Regulators (IGRs):** These chemicals interfere with the molting process of larvae, reducing their ability to mature and cause damage.

4. Monitoring and Early Detection:

- **Pheromone Traps:** Deploy pheromone traps to monitor adult moth activity. This can help in predicting infestations and timing control measures effectively.

Apple_Apple_scab:



Description:

Bacterial Blight in rice is a serious disease caused by the bacterium *Xanthomonas oryzae* pv. *oryzae* (Xoo). It primarily affects the leaves of rice plants, leading to significant yield losses. The disease is most prevalent in warm, humid climates, especially in areas with heavy rainfall and high temperatures.

Life Cycle:

- The bacterium infects the rice plant primarily through wounds or natural openings like stomata (tiny pores on the leaf surface).
- It can be spread by rain splash, wind, irrigation water, contaminated tools, and by contact with infected plants.

Damage:

- **Leaf Blight:** The disease causes blighting of leaves, which can result in a significant reduction in photosynthetic activity, leading to poor plant growth and reduced grain filling.
- **Yield Loss:** Severe infections can cause yield losses of up to 50% or more, depending on the stage of infection and environmental conditions.

Symptoms:

- Green water-soaked layer along the cut portion or leaf tip of leaves as early symptom.
- Leaves wilt and roll up and become grayish green to yellow.
- Water-soaked to yellowish stripes on leaf blades or starting at leaf tips with a wavy margin.

Prevention:

1. Use of Resistant Varieties:

- Planting resistant rice varieties is one of the most effective ways to prevent Bacterial Blight. Varieties like IR64, IR24, and other locally adapted strains that are resistant to Xoo should be used.

2. Cultural Practices:

- **Field Sanitation:** Remove and destroy infected plant residues and weeds after harvest to reduce sources of infection.
- **Proper Irrigation Management:** Ensure proper drainage and avoid excessive irrigation to minimize water stagnation, which can facilitate the spread of the bacterium.

3. Biological Control:

- **Antagonistic Bacteria:** Introducing beneficial bacteria such as *Pseudomonas fluorescens* can help suppress the growth of the pathogen through competitive exclusion or production of antibacterial compounds.

Apple_Cedar_apple_rust:



- Rust diseases are caused by fungi from the order Pucciniales. They are characterized by the formation of rust-colored pustules on plant tissues. These pustules contain spores that can spread the disease to other plants.
- Stem Rust (*Puccinia graminis*): Forms reddish-brown pustules on stems, leaves, and sometimes ears.
- Leaf Rust (*Puccinia triticina*): Produces orange-red pustules on the upper surfaces of leaves.

Lifecycle:

- **Teliospores:** These are the overwintering spores that survive harsh conditions. They are usually produced on the infected plant's stem or leaves.
- **Germination:** When conditions become favorable (e.g., warmth and moisture), teliospores germinate and produce basidiospores.
- **Infection:** The appressorium penetrates the plant tissues, establishing an infection. The fungus develops intercellular hyphae, spreading within the plant.

Damages:

- **Reduced Photosynthesis:** Pustules on leaves interfere with photosynthesis by reducing the effective leaf area. This leads to lower growth and yield.
- **Weakened Plants:** Rust infections weaken plants, making them more susceptible to other stresses and diseases.
- **Yield Loss:** Infected plants can have reduced grain or fruit production. For example, in wheat, rust can lead to shriveled grains and lower overall yield.

Symptoms:

- Common rust produces rust-colored to dark brown, elongated pustules on both leaf surfaces.
- The pustules contain rust spores (urediniospores) that are cinnamon brown in color.
- Pustules darken as they age. Leaves, as well as sheaths, can be infected.

Prevention of Common Rust:

- **Resistant Varieties:** Use crop varieties that are resistant or tolerant to rust diseases. This is often the most effective method for managing rust.
- **Crop Rotation:** Rotate crops to reduce the build-up of rust pathogens in the soil. Avoid planting the same crop or related crops in the same field consecutively.
- **Fungicide Application:** Apply appropriate fungicides as a preventive or curative measure, based on the rust type and crop. Follow guidelines for application timing and rates.
- **Early Detection and Monitoring:** Regularly inspect crops for early signs of rust and take action promptly if symptoms are observed.
- **Clean Equipment and Seed:** Ensure that equipment and seed are clean and free from fungal spores to prevent spreading the disease.

Tomato_Tomato_Yellow_Leaf_Curl_Virus:



Description:

Gray Leaf Spot is a fungal disease affecting crops such as corn and some grasses. It is caused by the fungus *Cercospora zeae-maydis* in corn. The disease manifests as irregularly shaped lesions on the leaves, initially appearing as small, dark gray to brown spots with a yellowish halo.

As the disease progresses, these spots can coalesce, leading to large, grayish areas of dead tissue. The lesions are often characterized by a distinctive dark center with a lighter gray edge, and they can cause premature leaf senescence.

Lifecycle:

- Gray Leaf Spot has a relatively straightforward lifecycle. The fungus overwinters in infected crop residues and produces conidia, which are the asexual spores responsible for spreading the disease.
- During the growing season, conidia are dispersed by wind and rain. Upon landing on susceptible plants, these spores germinate and penetrate the leaf tissue.

Damages:

- Gray Leaf Spot primarily impacts corn by causing significant reductions in photosynthetic efficiency.
- Infected leaves can become severely damaged, leading to a decrease in the overall plant health and vigor.
- This results in lower grain yield and reduced quality of the harvested corn.

Symptoms:

- Initially, lesions of gray leaf spot begin as a small dot with a yellow halo.
- Lesions will elongate over time running parallel to the veins becoming pale brown to gray and rectangular in shape with blunt ends.
- Leaves appear grayish in color due to the presence of fungal spores.

Prevention of Gray Leaf Spot:

- **Resistant Varieties:** Planting hybrids or varieties of crops that are resistant or tolerant to Gray Leaf Spot is one of the most effective preventive measures.
- **Crop Rotation:** Rotate corn with non-host crops (such as soybeans or small grains) to break the disease cycle. This helps reduce the amount of fungal inoculum in the field since the pathogen cannot survive without its host.
- **Residue Management:** Properly manage crop residues by tilling or removing infected plant material after harvest. This reduces the overwintering sites for the fungus, lowering the chances of infection in the next growing season.
- **Fungicide Application:** Apply fungicides, especially in environments conducive to disease development (warm and humid conditions). Fungicides can be used preventively .

Peach_Bacterial_spot:



Description:

Leaf Curl is a common fungal disease that primarily affects stone fruit trees, such as peaches, nectarines, and almonds, but it can also occur in other plants. The disease is caused by the fungus *Taphrina deformans*, which infects the leaves, buds, and sometimes the fruit of the tree.

Lifecycle:

- The lifecycle of the Leaf Curl fungus begins in the fall when it overwinters as spores (ascospores) on the surface of the tree bark and in buds.
- In early spring, as buds begin to swell and break dormancy, the fungus infects the newly emerging leaves.
- Moist, cool conditions (temperatures between 50-70°F) are ideal for infection, with rain or dew helping to activate the spores and spread them to susceptible tissue.

Damages:

- Leaf Curl can cause extensive damage to affected trees.
- This, in turn, can lead to reduced fruit production and poor fruit quality. In addition, the loss of leaves and weakening of the tree can make it more susceptible to other diseases and environmental stresses.

Symptoms:

- Plants are stunted or dwarfed
- Plants are stunted or dwarfed
- Leaflets are rolled upwards and inwards
- Leaves are often bent downwards, stiff, thicker than normal, have a leathery texture, show interveinal chlorosis and are wrinkled
- Affected plants tend to be distributed randomly or in patches

Prevention of Leaf Curl:

- Practice crop rotation by avoiding planting susceptible crops in the same area consecutively. For example, rotating tomato/chilli crops with non-host crops like legumes, onion and brassicas.
- Early planting can help establish crops before the peak disease period.
- Remove and destroy any plants showing symptoms of leaf curl disease.
- Remove weeds in and around the field.
- Grow barrier crops around the field to deter whiteflies. For example, maize or sorghum in chilli field, sesbania in tobacco field.
- Install 6 – 8 yellow sticky traps for per acre field to control the whitefly population.

Pepper,_bell_Bacterial_spot:



- Sugar Cane mosaic virus (SCMV) poses a significant threat to sugar cane crops worldwide, necessitating effective management strategies to mitigate its impact. One such approach involves the use of resistant cultivars, which can help reduce the spread of the virus and minimize yield losses.
- SCMV is a plant virus that primarily affects sugar cane plants, causing characteristic symptoms such as mosaic patterns on leaves, stunted growth, and reduced sugar content in infected plants. These symptoms can vary depending on factors such as the strain of the virus and the age of the plant.


Symptoms:

- The mosaic disease symptoms include distinct color variations on infected sugarcane plants, with chlorotic areas ranging from green to yellow.
- These patterns become more pronounced as the disease progresses.
- Young plants infected with SCMV exhibit a mosaic pattern of light green to yellow patches, while mature plants may show partial reddening or necrosis.
- Infected sugarcane plants suffer a decrease in photosynthetic efficiency, which directly impacts the yield due to the disruptive effects of the virus.
- The typical mosaic symptoms of sugarcane mosaic disease include green streaks on younger leaves and mosaic patterns, ranging from faint and irregular to pronounced streaking and patching.

Prevention of Sugar Cane Mosaic Virus:

- To prevent the spread of the Sugarcane Mosaic Virus (SCMV), it is crucial to implement proper sanitation measures and control strategies
- SCMV primarily spreads through infected plant sap and mechanical means such as lawnmowers and equipment.
- Therefore, to minimize transmission, it's essential to thoroughly clean all equipment used in sugarcane fields, ensuring that clippings and sap are removed
- Furthermore, avoiding the planting of virus-host crops near sugarcane fields can help reduce the risk of SCMV spread.
- Furthermore, considering the diversity of strains and the potential for co-infection, it's imperative to adopt integrated management approaches, including resistance breeding in susceptible crops.

Pepper,_bell_Bacterial_spot:



Blast, also called rotten neck, is one of the most destructive diseases of Missouri rice. Losses due to this disease have been on the increase since 2000.

Rice blast can be controlled by a combination of preventive measures and foliar fungicides applied when rice is in the late boot stage and again when it is 80 to 90 percent headed. Whenever possible, consult an expert in plant disease diagnosis.

Symptoms

- Blast symptoms can occur on leaves, leaf collars, nodes and panicles.
- Leaf spots are typically elliptical (football shaped), with gray-white centers and brown to red-brown margins.
- Fully developed leaf lesions are approximately 0.4 to 0.7 inch long and 0.1 to 0.2 inch wide.
- Both the shape and the color vary depending on the environment, age of the lesion and rice variety.
- Lesions on leaf sheaths, which rarely develop, resemble those on leaves.

Prevention:

- Plant the least-susceptible varieties and use a broad-spectrum seed treatment.
- Plant the least-susceptible varieties and use a broad-spectrum seed treatment.
- Grow rice in open fields free of tree lines particularly on east and south sides
- Seed over a range of time to spread the heading dates. However, avoid planting late because blast will be more severe.
- Seed to a stand of 15 to 20 plants per square foot.
- Avoid excessive nitrogen application rates and apply no more than 30 pounds per acre of nitrogen per application at midseason. In fields with a history of blast, always split applications.
- If symptoms are found, prepare to use fungicides at the late boot stage and again when 80 to 90 percent of plants are headed.



Tomato_Target_Spot:

The Wheat Stem Fly (*Meromyza americana*) is a significant pest of wheat and other cereal crops, particularly in regions with cool and dry climates. Adult flies are small, slender, and yellowish, resembling tiny house flies. They lay their eggs on the leaves of young wheat plants.

Symptoms:

1. Stunted Growth:

- Infested wheat plants often exhibit stunted growth. The larvae feeding inside the stem disrupt the flow of nutrients and water, leading to reduced plant vigor and overall size.

2. Yellowing and Chlorosis:

- Leaves of infested plants may turn yellow, particularly the lower leaves. This chlorosis is a result of the impaired ability of the plant to transport nutrients due to the larval feeding inside the stem.

Prevention of Wheat Stem Fly:

1. Crop Rotation:

- Implement crop rotation by planting non-host crops (such as legumes or oilseeds) after wheat. This disrupts the lifecycle of the Wheat Stem Fly by reducing the availability of suitable hosts for egg-laying and larval development.

2. Resistant Varieties:

- While specific resistance to Wheat Stem Fly is not widely available, planting early-maturing or fast-growing wheat varieties can help reduce the window of vulnerability to infestation. Strong, vigorous plants are less likely to suffer severe damage.

3. Field Monitoring:

- Regularly monitor fields for the presence of Wheat Stem Fly adults and signs of larval infestation. Early detection allows for timely intervention, although chemical control is generally not recommended due to the difficulty of targeting larvae inside stems.



Tomato_Bacterial_spot:

Wheat Scab, also known as Fusarium Head Blight (FHB), is a destructive fungal disease caused by *Fusarium* species, primarily *Fusarium graminearum*. It affects wheat, barley, and other small grains, particularly in regions with warm, humid conditions during the flowering and grain-filling stages.

Symptoms of Wheat Scab:

1. Bleached Spikelets:

- One of the most distinctive symptoms of Wheat Scab is the bleaching of spikelets on the wheat head. Infected spikelets turn a pale, straw-colored or bleached appearance, while healthy parts of the head remain green.

2. Shriveled and Discolored Grains:

- Infected grains, often called "tombstones," are shriveled, lightweight, and may appear pinkish, white, or grayish due to fungal growth.

3. Premature Ripening:

- Infected heads may ripen prematurely, leading to uneven maturity across the field. This can complicate harvest timing and reduce overall yield and quality.

Prevention of Wheat Scab:

1. Resistant Varieties:

- Plant wheat varieties that are resistant or tolerant to Fusarium Head Blight. Breeding programs have developed several resistant varieties that can significantly reduce the impact of the disease.

2. Fungicide Application:

- Apply fungicides during the flowering stage (Feekes growth stage 10.5.1) when the wheat is most susceptible to infection. Fungicides containing triazoles have been shown to be effective in reducing disease severity.

3. Monitoring and Early Detection:

- Regularly scout fields during the flowering stage to detect early signs of Wheat Scab. Early detection allows for timely fungicide applications and other interventions to limit disease spread.

