


RESEARCH ARTICLES | MARCH 31 2015

Smoking Is Associated with Acute and Chronic Prostatic Inflammation: Results from the REDUCE Study FREE

Daniel M. Moreira ; J. Curtis Nickel; Leah Gerber; Roberto L. Muller; Gerald L. Andriole; Ramiro Castro-Santamaria; Stephen J. Freedland



[+ Author & Article Information](#)

Cancer Prev Res (Phila) (2015) 8 (4): 312–317.

<https://doi.org/10.1158/1940-6207.CAPR-14-0260>

[Article history](#) 



View Metrics

Citing Articles Via

Web Of Science (26)

Google Scholar

CrossRef (24)

Abstract

Both anti- and proinflammatory effects of cigarette smoking have been described. As prostate inflammation is common, we hypothesized smoking could contribute to prostate inflammation. Thus, we evaluated the association of smoking status with acute and chronic inflammation within the prostate of men undergoing prostate biopsy. We retrospectively analyzed 8,190 men ages 50 to 75 years with PSA levels between 2.5 and 10 ng/mL enrolled in the Reduction by Dutasteride of Prostate Cancer Events study. Smoking status was self-defined as never, former, or current. Prostate inflammation was assessed by systematic central review blinded to smoking status. The association of smoking with inflammation in the baseline, 2-year, and 4-year biopsies was evaluated with univariable and multivariable logistic regressions. At study enrollment, 1,233 (15%), 3,203 (39%), and 3,754 (46%) men were current, former, and never smokers, respectively. Current smokers were significantly younger and had smaller prostates than former and never smokers (all $P < 0.05$). Former smokers were significantly heavier than current and never smokers ($P < 0.001$). Acute and chronic prostate inflammations were identified in 1,261 (15%) and 6,352 (78%) baseline biopsies, respectively. In univariable analysis, current smokers were more likely to have acute inflammation than former (OR, 1.35; P , 0.001) and never smokers (OR, 1.36; P , 0.001). The results

Email Alerts

- Article Activity Alert
- Online First Alert
- eTOC Alert

Latest	Most Read	Most Cited
Triple Primary Cancers: An Analysis of Genetic and Environmental Factors		
Next-generation Multi-target Stool DNA Panel Accurately Detects Colorectal Cancer and Advanced Precancerous Lesions		
Acceptability of Personalized Lung Cancer Screening Program Among Primary Care Providers		
High-Fat Diet Promotes Acute		

were unchanged at 2- and 4-year biopsies. In contrast, current smoking was linked with chronic inflammation in the baseline biopsy, but not at 2- and 4-year biopsies. In conclusion, among men undergoing prostate biopsy, current smoking was independently associated with acute and possibly chronic prostate inflammations. *Cancer Prev Res*; 8(4); 312–7. ©2015 AACR.

Promyelocytic Leukemia through PPAR δ -Enhanced Self-renewal of Preleukemic Progenitors

Introduction

Cigarette smoking has been associated with higher prostate cancer mortality in all comers and more disease recurrence, metastasis, and cancer-specific mortality after surgery and radiotherapy for prostate cancer (1–5). Although, the pathways by which smoking affects prostate carcinogenesis and prostate cancer progression are largely unknown, several potential mechanisms have been proposed, including genetic and epigenetic changes, enhanced angiogenesis, life style factors, and inflammation. Indeed, the effects of cigarette smoking on the immune system are far-reaching and may affect the prostate. For example, there is evidence suggesting cigarette smoking promotes an overall systemic inflammatory state by increasing levels of proinflammatory markers (6). Moreover, inflammatory changes in distant organs, such as eyes, arteries, and pancreas, have been described among active smokers (7–9). Conversely, active smoking has been linked to a decreased risk of ulcerative colitis, an inflammatory bowel disease (10), and rosacea, an inflammatory skin disorder (11). Thus, the direct inflammatory effects of cigarette smoking on several organs, including the prostate, remain to be determined.

Acute and chronic inflammatory infiltrates are frequently found in the prostate. Previous studies indicate that between 35% and 100% of prostate biopsies done for suspected prostate cancer had some histologic evidence of inflammation (12–15).

However, factors associated with inflammatory changes in the prostate of men undergoing biopsy are largely unknown. To date, no studies have evaluated the association of cigarette smoking with histologic prostatic inflammation. Therefore, we sought to evaluate the association of smoking status and histologic inflammation in the prostates of men undergoing prostate biopsy in the REDuction by DUtasteride of Prostate Cancer Events (REDUCE) trial. The REDUCE trial was a 4-year, placebo-controlled study evaluating daily dutasteride to reduce the risk of biopsy detectable prostate cancer among patients with negative baseline prostate biopsy (16). Repeat prostate biopsies were performed at 2 and 4 years. Thus, REDUCE uniquely allows the examination of the relationship between baseline cigarette smoking and inflammatory changes on baseline-negative biopsies and subsequent repeat prostate biopsies. We hypothesize that smoking is associated with greater risk of inflammation in the prostate.

Materials and Methods

REDUCE design was published previously (16). In brief, eligible men were ages 50 to 75 years, had serum PSA ≥ 2.5 or 3.0 ng/mL according to age (50–60 and 60–75 years, respectively) but ≤ 10 ng/mL, one single negative prostate biopsy (6 to 12 cores) within 6 months of enrollment. Men were excluded if they had history of prostate cancer, high-grade intraepithelial neoplasia, atypical small acinar proliferation, prostate volume (PV) > 80 mL, had undergone previous prostate surgery or had International Prostate Symptom Score ≥ 25 or ≥ 20 on alpha-blockers, or were previously on finasteride or dutasteride. Medical history was collected at baseline. All men were randomized in double-blind fashion to receive orally either dutasteride 0.5 mg or placebo daily and followed every 6 months for 4 years. To maintain the blinded nature of the study, PSA levels in the

dutasteride-treated men were doubled (given dutasteride reduces PSA levels by approximately half) and randomly adjusted by 0.1 ng/mL so that the final reported values were equally even and odd. PV was measured by ultrasonography at the time of randomization and 2 and 4 years later. Ten-core transrectal, ultrasound-guided biopsies were performed as part of the protocol at 2 and 4 years. Baseline biopsies had been performed before the start of the study (and independently of the study) and were reread centrally (at Bostwick Laboratories). Biopsies that were performed as part of the study were also read centrally. Central pathology laboratory had no access to the randomization codes or data on smoking status. Prostate cancer, acute and chronic prostate inflammation were coded as present or absent. Inflammation was also coded based on severity as mild, moderate and marked according to the average cell density and extent of tissue involvement in each biopsy core. Chronic inflammation consisted mainly of lymphocytes and variable number of plasma cells and macrophages. Acute inflammation consisted of neutrophils. Mild inflammation was defined as small scattered or patchy aggregates and presence of nests of inflammatory cells, usually no more than 10 to 15 cells per nest. Moderate inflammation was characterized by larger aggregates and nests, usually more than about 15 cells, invariably multifocal. Moderate inflammation was noticeable at low magnification. Sheets of inflammatory cells or extensive multifocal confluent masses of cells were noted in severe inflammation, which was obvious and noticeable at any magnification. Additional criterion for severe acute inflammation was tissue destruction of any size (17). Self-reported smoking status was determined at baseline and categorized in one of three groups: current, former, or never smoker. The protocol was approved by the Institutional

Review Board at each research site, and all participants provided written informed consent. Of the 8,231 patients enrolled in the REDUCE trial, we excluded 41 (<1%) due to missing data including smoking status or histologic inflammation. This resulted in a study sample of 8,190 (99%) men with centrally reviewed baseline biopsies and complete data. Positive baseline biopsies were not included in the study (i.e., negative baseline biopsy for prostate cancer was part of the inclusion criteria in REDUCE). Moreover, patients were selected for baseline biopsies based on PSA levels, whereas repeat biopsies were done regardless of PSA. To overcome this limitation, we analyzed the 2- and 4-year repeat biopsies given they include both positive and negative biopsies for prostate cancer. A total of 6,311 (77%) and 4,619 (56%) had 2- and 4-year per protocol repeat biopsies and were included in the repeat biopsy analyses. The details of men who underwent at least one on-study biopsy have been described previously ([18](#)).

Univariable comparisons of baseline characteristics between current, former, and never smokers were performed using χ^2 for categorical data and ANOVA for continuous variables. Given the low numbers of patients with moderate and marked acute and chronic inflammation, inflammation was analyzed as present or absent. The association of smoking status with acute and chronic inflammation in baseline, 2-, and 4-year repeat prostate biopsies was evaluated using logistic regression in univariable analysis. Multivariable logistic regression models were used to determine the association between smoking and prostate inflammation adjusting for baseline characteristics, such as age (continuous, in years), race (Caucasian, African American, Asian, American Hispanic, or other), body mass index (BMI, continuous and log-transformed, in kg/m²),

digital rectal exam (DRE, coded as normal or abnormal), PV (continuous and log-transformed, in cm³), PSA (continuous, in ng/mL) to test the association of smoking status and baseline prostate inflammation. The selection of covariates in multivariable models was based on their observed and/or previously established associations with baseline smoking status and/or inflammation (19). The analysis of smoking and inflammation at 2- and 4-year biopsies was adjusted for treatment arm as well. All statistical analyses were two-tailed and performed using Stata 11.2 (StataCorp). A *P* < 0.05 was considered statistically significant.

Results

Baseline patient characteristics

Mean baseline age, BMI, PSA, and PV of our cohort were 62.8 years, 27.4 kg/m², 6.1 ng/dL, and 45.7 cm³, respectively. A total of 7,455 (91.0%) subjects were Caucasian. Abnormal DRE was identified in 284 (3.9%) patients. A total of 1,233 (15.1%), 3,203 (39.1%), and 3,754 (45.8%) men were current, former, and never smoker, respectively. Current smokers were significantly younger and had smaller prostates than former and never smokers (all *P* < 0.05) but had similar race, DRE, prebiopsy PSA, and treatment arm (all *P* > 0.05). Former smokers had significantly higher BMI than current and never smokers (*P* < 0.001; **Table 1**).

Table 1.

Baseline patient characteristics by smoking status

	Smoking status	
Characteristic	Current	Former
<i>N</i> (%)	1,233 (15.1)	3,203 (39.1)
Age (in years), mean (SD)	61.7 (6.0)	63.2 (6.0)
Race, <i>N</i> (%)		

White	1,118 (90.7)	2,923 (91.3)
Black	30 (2.4)	62 (1.9)
Asian	30 (2.4)	54 (1.7)
American Hispanic	44 (3.6)	129 (4.0)
Other	11 (0.9)	35 (1.1)
BMI (in kg/m ²), mean (SD)	27.0 (5.2)	27.7 (3.9)
DRE, <i>N</i> (%)		
Normal	1,055 (95.9)	2,807 (96.2)
Abnormal	45 (4.1)	112 (3.8)
PSA (in ng/mL), mean (SD)	6.0 (2.3)	6.1 (2.3)
PV (in cm ³), mean (SD)	44.2 (17.1)	45.9 (17.4)
Treatment arm, <i>N</i> (%)		
Placebo	630 (51.1)	1,627 (50.8)
Dutasteride	603 (48.9)	1,574 (49.2)

^a χ^2 test for categorical variables and ANOVA for continuous variables.

Baseline biopsy

At baseline biopsy where all biopsies were negative for cancer, acute and chronic inflammation was detected in 1,261 (15.4%) and 6,352 (77.6%) cases, respectively. A total of 1,196 (14.6%) baseline biopsies showed both acute and chronic inflammation and 1,773 (21.7%) had no inflammation (Supplementary Table S1). Acute prostate inflammation was graded as mild, moderate, and marked in 1,235 (97.9%), 24 (1.9%), and 2 (0.2%), respectively. Chronic prostate inflammation was graded as mild, moderate, and marked in 5,653 (89.0%), 681 (10.7%), and 18 (0.3%), respectively. In univariable analysis, current smokers were more likely to have acute inflammation than former (OR, 1.35; *P*, 0.001) and never smokers (OR, 1.36; *P*, 0.001) at baseline biopsy. Similarly, in multivariable analysis, current smokers had

significantly more acute inflammation than former (OR, 1.29; *P*, 0.008) and never smokers (OR, 1.30; *P*, 0.005; **Table 2**) at baseline. Though there were trends for current smokers to be more likely to have chronic inflammation at baseline biopsy on univariable analysis, this did not reach statistical significance (**Table 2**). However, on multivariable analysis, current smokers were more likely to have chronic inflammation at baseline biopsy relative to former (OR, 1.23; *P*, 0.017) and never smokers (OR, 1.24; *P*, 0.014; **Table 2**). Most of the change in the OR of chronic inflammation occurred with the addition of PV to the multivariable models.

Table 2.

Association of smoking status with baseline acute and chronic prostate inflammations

Smoking status	Acute inflammation (<i>N</i> = 1,261; 15.4%)		
	Univariable		Multivariable
	OR (95% CI)	<i>P</i>	OR (95% CI)
Former vs. never	0.99 (0.87–1.13)	0.901	1.01 (0.87–1.17)
Current vs. never	1.35 (1.14–1.59)	0.001	1.30 (1.09–1.55) ^a

NOTE: Never smokers: 3,754 (45.8%); former smokers: 3,203 (39.1%); current smokers: 1,233 (15.1%).

Abbreviation: CI, confidence interval.
^aAdjusted for age, race, BMI, DRE, PV, and PSA.

2- and 4-year biopsies

At 2-year repeat prostate biopsy, the prevalence of acute and chronic inflammation was 1,405 (22.3%) and 5,084 (80.6%) cases, respectively. At 4-year repeat prostate biopsy, prevalence of acute and chronic inflammation was 1,067 (23.0%) and 3,690 (79.7%) cases, respectively (Supplementary Table S1). Supplementary Table S2 shows the prevalence inflammation by treatment arm. At both 2- and 4-year repeat prostate biopsies, baseline current smokers had significantly more acute inflammation than never smokers in both uni- and

multivariable analyses (all $P < 0.05$, **Tables 3** and **4**). Likewise, baseline current smokers had more acute inflammation than former smokers in both uni- and multivariable analyses at 2- (univariable: OR, 1.16; P , 0.105; multivariable: OR, 1.23; P , 0.022) and 4-year (univariable: OR, 1.35; P , 0.003; multivariable: OR, 1.35; P , 0.004) repeat prostate biopsies, although the associations did not reach statistical significance at 2-year repeat biopsy (**Tables 3** and **4**). Baseline smoking status was not associated with chronic prostate inflammation on either 2- or 4-year repeat biopsy on either univariable or multivariable analysis (**Tables 3** and **4**). Supplementary Table S3 shows the univariable ORs of inflammation by smoking status stratified by treatment arm. Supplementary Tables S4 and S5 show the association of smoking status with acute and chronic prostate inflammations at 2- and 4-year repeat biopsies by cancer status. Analysis of smoking status and inflammation severity (acute and chronic) was not possible due to limited number of patients with moderate and marked acute and chronic inflammations.

Table 3.
Association of smoking status with acute and chronic prostate inflammations at 2-year repeat biopsy

	Acute inflammation (N = 1,405; 22.3%)		
	Univariable		Multivariable
Smoking status	OR (95% CI)	P	OR (95% CI)
Former vs. never	1.06 (0.93–1.21)	0.387	1.06 (0.93–1.21)
Current vs. never	1.23 (1.03–1.46)	0.022	1.23 (1.03–1.46)

NOTE: Never smokers: 2,908 (46.1%); former smokers: 2,490 (39.4%); current smokers: 913 (14.5%).

Abbreviation: CI, confidence interval.
^aAdjusted for age, race, BMI, DRE, PV, PSA, and treatment arm.

Table 4.
Association of smoking status with acute and chronic prostate

inflammations at 4-year repeat biopsy

	Acute inflammation (N = 1,067; 23.0%)		
	Univariable		Multivariable
Smoking status	OR (95% CI)	P	OR (95% CI)
Former vs. never	1.00 (0.86–1.16)	0.998	1.00 (0.86–1.16)
Current vs. never	1.35 (1.10–1.66)	0.004	1.32 (1.07–1.63)

NOTE: Never smokers: 2,178 (47.0%); former smokers: 1,823 (39.4%); current smokers: 631 (13.6%).

Abbreviation: CI, confidence interval.
^aAdjusted for age, race, BMI, DRE, PV, PSA, and treatment arm.

Discussion

Cigarette smoking has a major impact on health issues worldwide. It is the leading cause of preventable morbidity and mortality in the United States (20). Several studies suggest many of the adverse consequences of smoking might be due to its effects on oxidative stress and immune-inflammatory system (6). Indeed, cigarette smoking has been linked to inflammation of specific organs, such as eyes, arteries, and pancreas (7–9). However, direct inflammatory effects of cigarette smoking on urologic organs including the prostate have not been described. Therefore, we retrospectively analyzed 8,190 men ages 50 to 75 years with PSA levels between 2.5 and 10 ng/mL enrolled in the REDUCE study. In both uni- and multivariable analyses, current smokers were more likely to have acute prostate inflammation than former and never smokers. Results were virtually unchanged in multivariable analysis and at 2- and 4-year repeat prostate biopsies. In addition, we found smokers were more likely to have chronic inflammation in their baseline biopsies.

To date, no study has examined the link between cigarette smoking and objectively assessed


prostatic inflammation. In contrast, a few studies evaluated the association of cigarette smoking with prostatitis-related symptoms, but with mixed results. The largest study published by Liang and colleagues (21) evaluating 12,743 Chinese men found significant association between active smoking status and self-reported prostatitis-like symptoms. In addition, Bartoletti and colleagues (22) using data from 764 prostatitis cases and 152 controls found significant association between smoking and symptoms of chronic prostatitis/chronic pelvic pain syndrome. In opposition, Mi and colleagues (23) investigating 1,242 Chinese men failed to demonstrate any association between smoking and prostatitis-like symptoms. Moreover, Wallner and colleagues (24) studying 703 African American men did not find any association between self-reported smoking status and prostatitis. Similarly, Moon and colleagues (25) examining 184 self-answered questionnaires filled out by 20 to 49 years-old subjects failed to identify any correlation between smoking and urinary symptoms. Thus, it remains controversial whether smoking has any association with prostatitis-like symptoms. However, prostatitis symptoms may not be due to prostatic inflammation *per se* but could relate to bladder issues or even irritants in the urine from smoking. Indeed, symptoms compatible with prostatitis are not always accompanied by histologic inflammation and vice versa (26). As such, based on current literature, it is not possible to infer whether cigarette smoking is associated with histologic prostate inflammation or not. Thus, our study is the first to evaluate the association between smoking and prostatic inflammation, and, importantly, we found current smokers are more likely to have prostate inflammation than former and never smokers.


The effects of cigarette smoking on host immunity


including the innate and adaptive immune systems are far-reaching and occur at local and systemic levels. Many of the smoking-induced toxic effects, including the induction of carcinogenesis, result from direct genetic or epigenetic changes leading to abnormal gene function. In general, tissue damage associated with smoking is considered the main cause for increased numbers of neutrophils seen in acute inflammation. The oxidative stress causes elevated concentration of cytokines which in turn activates polymorphonuclear neutrophils and prolongs their survival. Smoking components (particularly reactive oxygen species) also activate intracellular signaling cascades generating inflammatory gene activation (such as production of interleukin-8 and tumor necrosis factor; ref. [27](#)). However, not all effects of smoking on host immunity are stimulatory. There is evidence showing smoking may impair the function of inflammatory cells. For example, smoking reduces the ability of neutrophils and macrophages to respond to stimuli, phagocytose, kill pathogens, and secrete cytokines. It also reduces proliferation of T cells and alters CD4 and CD8 cells ([6](#)). Unfortunately, there are no studies evaluating mechanisms linking smoking to inflammation specifically in the prostate. Our findings, however, provide objective evidence that smoking is linked to changes within the prostate, suggesting that chemicals within cigarette smoke may make it to the prostate and stimulate inflammatory changes. Thus, further study is required to understand the biology of how cigarette smoking promotes prostate inflammation.


The clinical implications of histologic prostate inflammation are not completely defined. However, there is increasing evidence linking histologic prostate inflammation with benign prostate hyperplasia (BPH) and degree of lower urinary tract symptoms ([28, 29](#)). For example, a previous


study using data from the REDUCE trial found significant correlation between chronic inflammation and urinary symptoms (17). Subgroup analysis of the Medical Therapies of Prostate Symptoms study found men with prostate inflammation were significantly more likely to experience worsening symptoms or acute urinary retention than those without inflammation (30). In our study, although smoking was significantly associated with prostate inflammation, smokers had smaller prostates compared with nonsmokers. Likewise, previous studies found an inverse correlation between smoking and BPH (31, 32). Thus, it seems the effects of smoking in PV and BPH may be mediated by other mechanisms than inflammation. Also, there is enough evidence to support that prostate inflammation is associated with elevated PSA levels (33–35). In our study, although smoking was significantly associated with prostate inflammation, no differences in PSA levels were seen between smokers and nonsmokers. In


 Split-Screen


 Views ▾

 PDF

 Share ▾

 Tools ▾

 Search Site

 Versions ▾

Volume 8, Issue 4
1 April 2015



[◀ Previous Article](#) [Next Article ▶](#)

Article Contents

- Abstract
- Introduction

Thus, it remains unclear whether smoking may affect PSA levels and if it does so by promoting prostate inflammation. Finally, the role of histologic prostate inflammation and smoking in prostate cancer is unknown. Multiple studies have suggested possible association between inflammation and clinical prostatitis with prostate cancer risk (37–40). For instance, several authors have demonstrated positive correlation between urogenital infection, prostatitis, and sexually transmitted disease with prostate cancer (4, 38, 40). Conversely, more recent data suggest histologic inflammation in the biopsy specimen may actually be associated with decreased risk of prostate cancer detection in subsequent biopsies (19, 41–43). Similarly, a meta-analysis of 24 cohort studies found an association between active smoking and prostate cancer risk (44). We have

Materials and Methods
Results
Discussion
Disclosure of Potential Conflicts of Interest
Authors' Contributions

also linked smoking to high-grade prostate cancer in REDUCE (45). However, others found decreased risk of prostate cancer among smokers (46). Thus, the clinical implication of the association between smoking and prostate inflammation could not be ascertained by the current study and require further investigation.

The REDUCE study has many strengths, including its large international and multicentric population, per-protocol biopsies regardless of PSA levels, prospective data acquisition, and central pathology review. This is crucial in that prostatic inflammation can lead to PSA elevations resulting in prostate biopsy. As such, uniform biopsy protocol not driven by PSA fluctuations is crucial to prevent biases from inflammatory-driven biopsies. The main limitation of the present study is that only patients with PSA values between 2.5 and 10 ng/dL were

This site uses cookies. By continuing to use our website, you are agreeing to [our privacy policy](#). **Accept**

acinar proliferation, PV >80 mL, had undergone previous prostate surgery, or had an International Prostate Symptom Score ≥ 25 or ≥ 20 on alpha-blockers were excluded. Thus, all baseline biopsies were negative for cancer. Although these exclusions increase the homogeneity of the sample, they limit the generalizability of our results. Furthermore, the limited number of non-Caucasian men prevented us from exploring whether race influenced the association between smoking and prostatic inflammation. We did not consider off-protocol biopsies, including transrectal/transperineal needle biopsies, transurethral resections, and prostatectomies of any kind. In addition, patients were followed for 4 years, thus we were unable to study long-term consequences of smoking on prostate inflammation nor long-term clinical implications of prostate inflammation. We did not evaluate cumulative exposure, smoking duration, and time

since quitting smoking given the data were not available. Lastly, smoking status was determined at baseline only; therefore, we were not able to analyze changes in smoking status during the study interval.

In conclusion, among men undergoing prostate biopsy, current smoking status was independently associated with inflammatory changes in the prostate. Although the clinical implications of this finding were not determined by the present study, these findings provide an objective evidence that smoking is linked to changes within the prostate, suggesting that chemicals within cigarette smoke may make it to the prostate. This might be a mechanistic explanation for how smoking may influence prostatic conditions such as prostatitis or prostate cancer.

Disclosure of Potential Conflicts of Interest

J.C. Nickel reports receiving a commercial research grant and has provided expert testimony for Glaxo Smith Kline. G.L. Andriole reports receiving other commercial research support from Johnson & Johnson, Medivation, and Trasxxson. He is also a consultant/advisory board member for Augmenix, Bayer, Genomic Health, GlaxoSmithKline, and Myriad Genetics. R. Castro-Santamaria has ownership interest (including patents) in GSK. S.J. Freedland reports receiving a commercial research grant from GSK. No potential conflicts of interest were disclosed by the other authors.

Authors' Contributions

Conception and design: D.M. Moreira, J.C. Nickel, G.L. Andriole, S.J. Freedland

Development of methodology: D.M. Moreira, J.C. Nickel, L. Gerber, G.L. Andriole, S.J.

Freedland

Acquisition of data (provided animals, acquired and managed patients, provided facilities, etc.): D.M. Moreira, J.C. Nickel, L. Gerber, G.L. Andriole, S.J. Freedland

Analysis and interpretation of data (e.g., statistical analysis, biostatistics, computational analysis): D.M. Moreira, J.C. Nickel, L. Gerber, R.L. Muller, G.L. Andriole, S.J. Freedland

Writing, review, and/or revision of the manuscript: D.M. Moreira, J.C. Nickel, R.L. Muller, G.L. Andriole, R. Castro-Santamaria, S.J. Freedland

Administrative, technical, or material support (i.e., reporting or organizing data, constructing databases): D.M. Moreira, L. Gerber, S.J. Freedland

Study supervision: S.J. Freedland

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

References

1. Moreira DM, Aronson WJ, Terris MK, Kane CJ, Amling CL, Cooperberg MR, et al Cigarette smoking is associated with an increased risk of biochemical disease recurrence, metastasis, castration-resistant prostate cancer, and mortality after radical prostatectomy: Results from the SEARCH database. *Cancer* 2014;120:197–204.
Google Scholar Crossref PubMed
2. Islami F, Moreira DM, Boffetta P, Freedland SJ. A systematic review and meta-analysis of tobacco use and prostate cancer

mortality and incidence in prospective cohort studies. *Eur Urol* 2014;66:1054–64.

[Google Scholar](#) [Crossref](#)
[PubMed](#)

3. Kenfield SA, Stampfer MJ, Chan JM, Giovannucci E. Smoking and prostate cancer survival and recurrence. *JAMA* 2011;305:2548–55.

[Google Scholar](#) [Crossref](#)
[PubMed](#)

4. Pantarotto J, Malone S, Dahrouge S, Gallant V, Eapen L. Smoking is associated with worse outcomes in patients with prostate cancer treated by radical radiotherapy. *BJU Int* 2007;99:564–9.

[Google Scholar](#) [Crossref](#)
[PubMed](#)

5. Pickles T, Liu M, Berthelet E, Kim-Sing C, Kwan W, Tyldesley S, et al The effect of smoking on outcome following external radiation for localized prostate cancer. *J Urol* 2004;171:1543–6.

[Google Scholar](#) [Crossref](#)
[PubMed](#)

6. Goncalves RB, Coletta RD, Silverio KG, Benevides L, Casati MZ, da Silva JS, et al Impact of smoking on inflammation: overview of molecular mechanisms. *Inflamm Res* 2011;60:409–24.

[Google Scholar](#) [Crossref](#)
[PubMed](#)

7. Malfertheiner P, Schutte K. Smoking—a trigger for chronic inflammation and cancer development in the pancreas. *Am J Gastroenterol* 2006;101:160–2.

[Google Scholar](#) [Crossref](#)
[PubMed](#)

8. Galor A, Feuer W, Kempen JH, Kacmaz RO, Liesegang TL, Suhler EB, et al Adverse effects of smoking on patients with ocular inflammation. *Br J Ophthalmol* 2010;94:848–53.

[Google Scholar](#) [Crossref](#)
[PubMed](#)

9. Zou N, Hong J, Dai QY. Passive cigarette smoking induces inflammatory injury in human arterial walls. *Chin Med J* 2009;122:444–8.
[Google Scholar](#) [PubMed](#)
10. Mahid SS, Minor KS, Soto RE, Hornung CA, Galandiuk S. Smoking and inflammatory bowel disease: a meta-analysis. *Mayo Clinic proceedings Mayo Clinic* 2006;81:1462–71.
[Google Scholar](#) [Crossref](#)
11. Spoenclin J, Voegel JJ, Jick SS, Meier CR. A study on the epidemiology of rosacea in the U.K. *Br J Dermatol* 2012;167:598–605.
[Google Scholar](#) [Crossref](#)
[PubMed](#)
12. Pepe P, Aragona F. Does an inflammatory pattern at primary biopsy suggest a lower risk for prostate cancer at repeated saturation prostate biopsy? *Urol Int* 2011;87:171–4.
[Google Scholar](#) [Crossref](#)
[PubMed](#)
13. Gui-Zhong L, Libo M, Guanglin H, Jianwei W. The correlation of extent and grade of inflammation with serum PSA levels in patients with IV prostatitis. *Int Urol Nephrol* 2011;43:295–301.
[Google Scholar](#) [Crossref](#)
[PubMed](#)
14. Sfanos KS, De Marzo AM. Prostate cancer and inflammation: the evidence. *Histopathology* 2012;60:199–215.
[Google Scholar](#) [Crossref](#)
[PubMed](#)
15. Schatteman PH, Hoekx L, Wyndaele JJ, Jeuris W, Van Marck E. Inflammation in prostate biopsies of men without prostatic malignancy or clinical prostatitis: correlation with total serum PSA and PSA density. *Eur Urol* 2000;37:404–12.
[Google Scholar](#) [Crossref](#)
[PubMed](#)
16. Andriole GL, Bostwick DG, Brawley OW, Gomella LG, Marberger M, Montorsi F, et al. Effect of dutasteride on the risk of prostate cancer. *N Engl J Med* 2010;362:1192–202.

[Google Scholar](#) [Crossref](#)
[PubMed](#)

17. Nickel JC, Roehrborn CG, O'Leary MP, Bostwick DG, Somerville MC, Rittmaster RS. The relationship between prostate inflammation and lower urinary tract symptoms: examination of baseline data from the REDUCE trial. *Eur Urol* 2008;54:1379–84.

[Google Scholar](#) [Crossref](#)
[PubMed](#)

18. Wu C, Moreira DM, Gerber L, Rittmaster RS, Andriole GL, Freedland SJ. Diabetes and prostate cancer risk in the REDUCE trial. *Prostate Cancer Prostatic Dis* 2011;14:326–31.

[Google Scholar](#) [Crossref](#)
[PubMed](#)

19. Moreira DM, Nickel JC, Gerber L, Muller RL, Andriole GL, Castro-Santamaria R, et al Baseline prostate inflammation is associated with a reduced risk of prostate cancer in men undergoing repeat prostate biopsy: Results from the REDUCE study. *Cancer* 2014;120:190–6.

[Google Scholar](#) [Crossref](#)
[PubMed](#)

20. Danaei G, Ding EL, Mozaffarian D, Taylor B, Rehm J, Murray CJ, et al The preventable causes of death in the United States: comparative risk assessment of dietary, lifestyle, and metabolic risk factors. *PLoS Med* 2009;6:e1000058.

[Google Scholar](#) [Crossref](#)
[PubMed](#)

21. Liang CZ, Li HJ, Wang ZP, Xing JP, Hu WL, Zhang TF, et al The prevalence of prostatitis-like symptoms in China. *J Urol* 2009;182:558–63.

[Google Scholar](#) [Crossref](#)
[PubMed](#)

22. Bartoletti R, Cai T, Mondaini N, Dinelli N, Pinzi N, Pavone C, et al Prevalence, incidence estimation, risk factors and characterization of chronic prostatitis/chronic pelvic pain syndrome in urological hospital outpatients in Italy: results of a multicenter case-control observational

study. *J Urol* 2007;178:2411–5; discussion 5.

[Google Scholar](#) [Crossref](#)

[PubMed](#)

23. Mi H, Gao Y, Yan Y, Wu Y, Tan A, Yang X, et al Research of correlation between the amount of leukocyte in EPS and NIH-CPSI: result from 1242 men in Fangchenggang Area in Guangxi Province. *Urology* 2012;79:403–8.

[Google Scholar](#) [Crossref](#)

[PubMed](#)

24. Wallner LP, Clemens JQ, Sarma AV. Prevalence of and risk factors for prostatitis in African American men: the Flint Men's Health Study. *Prostate* 2009;69:24–32.

[Google Scholar](#) [Crossref](#)

[PubMed](#)

25. Moon TD, Hagen L, Heisey DM. Urinary symptomatology in younger men. *Urology* 1997;50:700–3.

[Google Scholar](#) [Crossref](#)

[PubMed](#)

26. Nickel JC. The overlapping lower urinary tract symptoms of benign prostatic hyperplasia and prostatitis. *Curr Opin Urol* 2006;16:5–10.

[Google Scholar](#) [Crossref](#)

[PubMed](#)

27. Lee J, Taneja V, Vassallo R. Cigarette smoking and inflammation: cellular and molecular mechanisms. *J Dent Res* 2012;91:142–9.

[Google Scholar](#) [Crossref](#)

[PubMed](#)

28. Kim SH, Jung KI, Koh JS, Min KO, Cho SY, Kim HW. Lower urinary tract symptoms in benign prostatic hyperplasia patients: orchestrated by chronic prostatic inflammation and prostatic calculi. *Urol Int* 2013;90:144–9.

[Google Scholar](#) [Crossref](#)

[PubMed](#)

29. Kwak C, Ku JH, Kim T, Park DW, Choi KY, Lee E, et al Effect of subclinical prostatic inflammation on serum PSA levels in men with clinically undetectable prostate cancer. *Urology* 2003;62:854–9.

[Google Scholar](#) [Crossref](#)
[PubMed](#)

30. McConnell JD, Roehrborn CG, Bautista OM, Andriole GL Jr, Dixon CM, Kusek JW, et al The long-term effect of doxazosin, finasteride, and combination therapy on the clinical progression of benign prostatic hyperplasia. *N Engl J Med* 2003;349:2387–98.

[Google Scholar](#) [Crossref](#)
[PubMed](#)

31. Meigs JB, Mohr B, Barry MJ, Collins MM, McKinlay JB. Risk factors for clinical benign prostatic hyperplasia in a community-based population of healthy aging men. *J Clin Epidemiol* 2001;54:935–44.

[Google Scholar](#) [Crossref](#)
[PubMed](#)

32. Kupeli B, Soygur T, Aydos K, Ozdiler E, Kupeli S. The role of cigarette smoking in prostatic enlargement. *Br J Urol* 1997;80:201–4.

[Google Scholar](#) [Crossref](#)
[PubMed](#)

33. Okada K, Kojima M, Naya Y, Kamoi K, Yokoyama K, Takamatsu T, et al Correlation of histological inflammation in needle biopsy specimens with serum prostate-specific antigen levels in men with negative biopsy for prostate cancer. *Urology* 2000;55:892–8.

[Google Scholar](#) [Crossref](#)
[PubMed](#)

34. Nadler RB, Collins MM, Probert KJ, Mikolajczyk SD, Knauss JS, Landis JR, et al Prostate-specific antigen test in diagnostic evaluation of chronic prostatitis/chronic pelvic pain syndrome. *Urology* 2006;67:337–42.

[Google Scholar](#) [Crossref](#)
[PubMed](#)

35. Carver BS, Bozeman CB, Williams BJ, Venable DD. The prevalence of men with National Institutes of Health category IV prostatitis and association with serum prostate specific antigen. *J Urol* 2003;169:589–91.

[Google Scholar](#) [Crossref](#)
[PubMed](#)

36. Ankerst DP, Boeck A, Freedland SJ, Thompson IM, Cronin AM, Roobol MJ, et al. Evaluating the PCPT risk calculator in ten international biopsy cohorts: results from the Prostate Biopsy Collaborative Group. *World J Urol* 2012;30:181–7.
[Google Scholar](#) [Crossref](#)
[PubMed](#)
37. Rosenblatt KA, Wicklund KG, Stanford JL. Sexual factors and the risk of prostate cancer. *Am J Epidemiol* 2001;153:1152–8.
[Google Scholar](#) [Crossref](#)
[PubMed](#)
38. Sarma AV, McLaughlin JC, Wallner LP, Dunn RL, Cooney KA, Schottenfeld D, et al. Sexual behavior, sexually transmitted diseases and prostatitis: the risk of prostate cancer in black men. *J Urol* 2006;176:1108–13.
[Google Scholar](#) [Crossref](#)
[PubMed](#)
39. Dennis LK, Lynch CF, Torner JC. Epidemiologic association between prostatitis and prostate cancer. *Urology* 2002;60:78–83.
[Google Scholar](#) [Crossref](#)
[PubMed](#)
40. Cheng I, Witte JS, Jacobsen SJ, Haque R, Quinn VP, Quesenberry CP, et al. Prostatitis, sexually transmitted diseases, and prostate cancer: the California Men's Health Study. *PLoS ONE* 2010;5:e8736.
[Google Scholar](#) [Crossref](#)
[PubMed](#)
41. Gann PH, Fought A, Deaton R, Catalona WJ, Vonesh E. Risk factors for prostate cancer detection after a negative biopsy: a novel multivariable longitudinal approach. *J Clin Oncol* 2010;28:1714–20.
[Google Scholar](#) [Crossref](#)
[PubMed](#)
42. Fujita K, Hosomi M, Tanigawa G, Okumi M, Fushimi H, Yamaguchi S. Prostatic inflammation detected in initial biopsy specimens and urinary pyuria are predictors of negative repeat prostate biopsy. *J Urol* 2011;185:1722–7.

[Google Scholar](#) [Crossref](#)
[PubMed](#)

43. Kryvenko ON, Jankowski M, Chitale DA, Tang D, Rundle A, Trudeau S, et al Inflammation and preneoplastic lesions in benign prostate as risk factors for prostate cancer. *Mod Pathol* 2012;25:1023–32.

[Google Scholar](#) [Crossref](#)
[PubMed](#)

44. Huncharek M, Haddock KS, Reid R, Kupelnick B. Smoking as a risk factor for prostate cancer: a meta-analysis of 24 prospective cohort studies. *Am J Public Health* 2010;100:693–701.

[Google Scholar](#) [Crossref](#)
[PubMed](#)

45. Ho T, Howard LE, Vidal AC, Gerber L, Moreira D, McKeever M, et al Smoking and Risk of Low- and High-Grade Prostate Cancer: Results from the REDUCE Study. *Clin Cancer Res* 2014;20:5331–8.

[Google Scholar](#) [Crossref](#)
[PubMed](#)

46. Watters JL, Park Y, Hollenbeck A, Schatzkin A, Albanes D. Cigarette smoking and prostate cancer in a prospective US cohort study. *Cancer Epidemiol Biomarkers Prev* 2009;18:2427–35.

[Google Scholar](#) [Crossref](#)
[PubMed](#)

©2015 American Association for Cancer Research.

Supplementary data

Supplemental Table 1: Inflammation by smoking status				
Inflammation	Smoking status			Overall
	Never	Former	Current	
Baseline biopsy				
Acute inflammation				
Present	556 (15%)	471 (15%)	234 (19%)	1261 (15%)
Absent	3198 (85%)	2732 (85%)	999 (81%)	6929 (85%)
Chronic inflammation				
Present	854 (23%)	732 (23%)	252 (20%)	1838 (22%)
Absent	2900 (77%)	2471 (77%)	981 (80%)	6352 (78%)
2-year biopsy				
Acute inflammation				
Present	621 (21%)	556 (22%)	228 (25%)	1405 (22%)
Absent	2287 (79%)	1934 (78%)	685 (75%)	4906 (78%)
Chronic inflammation				
Present	2339 (80%)	2009 (81%)	736 (81%)	5084 (81%)
Absent	569 (20%)	481 (19%)	177 (19%)	1227 (19%)
4-year biopsy				
Acute inflammation				
Present	485 (22%)	406 (22%)	176 (28%)	1067 (23%)
Absent	1693 (78%)	1417 (78%)	455 (72%)	3565 (77%)

figshare



Share

Download

Supplemental tables 1-5
Supplemental tables 1-5. Supplemental Table 1:
Inflammation by smoking status. Supplemental Table
2: Inflammation by treatment arm. Supplemental Table
3: Univariable odds ratios of inflammation by
smoking status stratified by treatment arm.
Supplemental Table 4: Association of smoking status
with acute and chronic prostate inflammations at 2-
year repeat biopsy by cancer status. Supplemental
Table 5: Association of smoking status with acute and
chronic prostate inflammations at 4-year repeat
biopsy by cancer status.
- docx file

AACR Journals

- Blood Cancer Discovery
- Cancer Discovery
- Cancer Epidemiology, Biomarkers & Prevention
- Cancer Immunology Research
- Cancer Prevention Research
- Cancer Research
- Cancer Research Communications
- Clinical Cancer Research
- Molecular Cancer Research
- Molecular Cancer Therapeutics

- Info for Advertisers
- Information for Institutions/Librarians
- RSS Feeds
- Privacy Policy



Copyright © 2023 by the American Association for Cancer Research.