

Reductive groups of semisimple rank 1

SGA 3, Exposé XX

English draft

M. Demazure

About this draft

This is an unofficial English translation of Exposé XX, “Reductive groups of semisimple rank 1” from *Schémas en groupes* (SGA 3). The source is the recomposition edited by P. Gille and P. Polo, PDF Exp20-13oct24.pdf (35 pages), from <https://webusers.imj-prg.fr/~patrick.polo/SGA3/>. The seminar was directed by M. Demazure and A. Grothendieck (IHÉS, 1962–1964).

Statement numbers follow that PDF. A clear typographical slip is corrected in the English and marked with a % `typo:` comment; such slips are listed in the README. A slip that would change the mathematics is left as printed and marked % `typo?:`. Scholarly proofreading remains outstanding.

The translator’s contribution is licensed under the MIT License. The French original remains copyright of its authors and publishers.

Contents

About this draft	iii
Exposé XX. Reductive groups of semisimple rank 1	1
1. Elementary systems. The groups U_α and $U_{-\alpha}$	1
2. Structure of elementary systems	7
3. The Weyl group	14
4. The isomorphism theorem	20
5. Examples of elementary systems, applications	21
6. Generators and relations for an elementary system	26

Exposé XX. Reductive groups of semisimple rank 1

by M. Demazure

1. Elementary systems. The groups U_α and $U_{-\alpha}$

(0)

Reminder 1.1. Let $S = \text{Spec}(k)$, where k is an algebraically closed field, and let G be a reductive S -group of semisimple rank 1, T a maximal torus (necessarily split) of G . One then has

$$\mathfrak{g} = \mathfrak{t} \oplus \mathfrak{g}^\alpha \oplus \mathfrak{g}^{-\alpha},$$

where α and $-\alpha$ are the roots of G with respect to T . Moreover, there exist two monomorphisms of groups

$$p_\alpha : \mathbf{G}_{a,S} \longrightarrow G \quad \text{and} \quad p_{-\alpha} : \mathbf{G}_{a,S} \longrightarrow G$$

such that

$$tp_\alpha(x)t^{-1} = p(\alpha(t)x) \quad \text{and} \quad tp_{-\alpha}(x)t^{-1} = p_{-\alpha}(\alpha(t)^{-1}x),$$

for every $S' \longrightarrow S$ and all $t \in T(S')$, $x \in \mathbf{G}_a(S')$, and such that the morphism

$$\mathbf{G}_{a,S} \times_S T \times_S \mathbf{G}_{a,S} \longrightarrow G,$$

defined by $(y, t, x) \longmapsto p_{-\alpha}(y)tp_\alpha(x)$, is radicial and dominant (*Bible*, § 13.4, cor. 2 to th. 3).

Since the tangent map at the identity element is bijective, this morphism is also separable, hence birational; by Zariski's "Main Theorem" (EGA III₁, 4.4.9), it is therefore an open immersion.

Lemma 1.2. Let S be a scheme, G a group S -scheme, T a torus of G , Q a subtorus of T , α a character of T inducing on Q a character nontrivial on each fiber. Let $p_\alpha : \mathbf{G}_{a,S} \longrightarrow G$ (resp. $p_{-\alpha} : \mathbf{G}_{a,S} \longrightarrow G$) be a morphism of groups normalized by T with multiplier α (resp. $-\alpha$). Suppose that the morphism

$$u : \mathbf{G}_{a,S} \times_S T \times_S \mathbf{G}_{a,S} \longrightarrow G$$

defined set-theoretically by $u(y, t, x) = p_{-\alpha}(y)tp_\alpha(x)$ is an open immersion. Finally, let q be an integer ≥ 0 and

$$p : \mathbf{G}_{a,S} \longrightarrow G$$

a morphism of groups such that for every $S' \longrightarrow S$ and all $t \in Q(S')$, $x \in \mathbf{G}_a(S')$, one has

$$\text{int}(t)^q(p(x)) = p(\alpha(t)x).$$

There then exists a unique $\nu \in \mathbf{G}_a(S)$ such that $p(x) = p_\alpha(\nu x^q)$.

Indeed, let Ω be the image of u and $U = p^{-1}(\Omega)$. This is an open subset of $\mathbf{G}_{a,S}$ containing the zero section. For every section t of Q , the automorphism of $\mathbf{G}_{a,S}$ defined by multiplication by $\alpha(t)$ preserves U globally. One has $U = \mathbf{G}_{a,S}$; indeed, it suffices to check this when S is the spectrum of an algebraically closed field k ; then $\alpha : Q(k) \longrightarrow k^*$ is surjective, which proves at once that $U(k) \supset k^*$, hence $U = \mathbf{G}_{a,k}$. There therefore exist three morphisms

$$a : \mathbf{G}_{a,S} \longrightarrow \mathbf{G}_{a,S}, \quad b : \mathbf{G}_{a,S} \longrightarrow T, \quad c : \mathbf{G}_{a,S} \longrightarrow \mathbf{G}_{a,S},$$

(0)**N.D.E.** Version of 13/10/2024

such that

$$p(x) = p_{-\alpha}(a(x))b(x)p_{\alpha}(c(x)).$$

The condition on p becomes

$$\begin{aligned} a(\alpha(t)x) &= \alpha(t)^{-q}a(x), \\ b(\alpha(t)x) &= b(x), \\ c(\alpha(t)x) &= \alpha(t)^q c(x). \end{aligned}$$

For the same reason as above, for every $S' \rightarrow S$ and every $z \in \mathbf{G}_m(S')$, one therefore has

$$a(zx) = z^{-q}a(x), \quad b(zx) = b(x), \quad c(zx) = z^q c(x),$$

hence

$$z^q a(z) = a(1), \quad b(z) = b(1), \quad c(z) = z^q c(1).$$

Since $\mathbf{G}_{m,S}$ is schematically dense in $\mathbf{G}_{a,S}$, one has at once, for every $x \in \mathbf{G}_a(S')$, $S' \rightarrow S$:

$$\begin{aligned} x^q a(x) &= a(1) = a(0) = 0, & \text{whence } a &= 0, \\ c(x) &= x^q c(1) = \nu x^q, & \text{for some } \nu \in \mathbf{G}_a(S), \\ b(x) &= b(1) = b(0) = e, & \text{whence } b &= e, \end{aligned}$$

which completes the proof.

Definition 1.3. Let S be a scheme. An *elementary S -system* is a triple (G, T, α) where

- (i) G is a reductive S -group of semisimple rank 1 (Exp. XIX 2.7),
- (ii) T is a maximal torus of G ,
- (iii) α is a root of G with respect to T (Exp. XIX 3.2).

One therefore has a direct sum decomposition (Exp. XIX 3.5)⁽¹⁾

$$\mathfrak{g} = \mathfrak{t} \oplus \mathfrak{g}^{\alpha} \oplus \mathfrak{g}^{-\alpha},$$

$\mathfrak{g}^{\alpha}$ and $\mathfrak{g}^{-\alpha}$ being locally free of rank one.

1.4. — If (G, T, α) is an elementary S -system, then $(G_{S'}, T_{S'}, \alpha_{S'})$ is an elementary S' -system for every $S' \rightarrow S$. If (G, T, α) is an elementary S -system, then $(G, T, -\alpha)$ is also one.

If S is a scheme, G a reductive S -group, T a maximal torus of G , α a root of G with respect to T , then (Exp. XIX 3.9) (Z_{α}, T, α) is an elementary S -system.

Let (G, T, α) be an elementary S -system. The invertible module $\mathfrak{g}^{\alpha}$ is canonically equipped with a T -module structure. One therefore also has a T -module structure on the vector bundle $W(\mathfrak{g}^{\alpha})$. On the other hand, the inner automorphisms of T define on G a structure of group with operator group T .

Theorem 1.5. Let (G, T, α) be an elementary S -system.

- (i) There exists a unique morphism of groups with operator group T

$$\exp : W(\mathfrak{g}^{\alpha}) \rightarrow G$$

which induces on Lie algebras the canonical morphism $\mathfrak{g}^{\alpha} \rightarrow \mathfrak{g}$.⁽²⁾

In other words, $\exp$ is the unique morphism satisfying the following conditions: for every $S' \rightarrow S$ and all $t \in T(S')$, $X, X' \in W(\mathfrak{g}^{\alpha})(S')$, one has

$$\begin{aligned} \exp(X + X') &= \exp(X) \exp(X'), \\ \text{int}(t)(\exp(X)) &= \exp(\alpha(t)X), \\ \underline{\text{Lie}}(\exp)(X) &= X. \end{aligned}$$

⁽¹⁾**N.D.E.** Of $\mathcal{O}_S$ -modules.

⁽²⁾**N.D.E.** We shall see later (Cor. 5.9) that $\exp$ is an isomorphism from $W(\mathfrak{g}^{\alpha})$ onto a closed subgroup of G .

(ii) If one defines similarly (in the elementary S -system $(G, T, -\alpha)$)

$$\exp : W(\mathfrak{g}^{-\alpha}) \longrightarrow G,$$

then the morphism

$$W(\mathfrak{g}^{-\alpha}) \times_S T \times_S W(\mathfrak{g}^{\alpha}) \longrightarrow G$$

defined set-theoretically by $(Y, t, X) \mapsto \exp(Y) \cdot t \cdot \exp(X)$ is an open immersion.

Suppose we have proved the existence of the required morphisms $\exp$, and let us prove the other assertions of the theorem. Let us first prove (ii). Since both sides are of finite presentation and flat over S , it suffices to do so when S is the spectrum of an algebraically closed field (SGA 1, I 5.7 and VIII 5.5). Let then $S = \text{Spec } k$. Let $X \in \Gamma(S, \mathfrak{g}^{\alpha})^{\times}$, $Y \in \Gamma(S, \mathfrak{g}^{-\alpha})^{\times}$. It suffices to prove that the morphism

$$\mathbf{G}_{a,k} \times_k T \times_k \mathbf{G}_{a,k} \longrightarrow G, \quad (y, t, x) \mapsto \exp(yY)t \exp(xX)$$

is an open immersion. Now, by 1.1 and 1.2, there exist $a, b \in k$ such that

$$\exp(yY) = p_{-\alpha}(ay) \quad \text{and} \quad \exp(xX) = p_{\alpha}(bx).$$

Since $\exp : W(\mathfrak{g}^{-\alpha}) \longrightarrow G$ induces a monomorphism on Lie algebras, one has $a \neq 0$; similarly $b \neq 0$, and one is reduced to 1.1.

The uniqueness of the morphism $\exp$ can be proved locally on S ; one then reduces to the case where $\mathfrak{g}^{\alpha}$ and $\mathfrak{g}^{-\alpha}$ are free, and one need only apply 1.2 (with $Q = T$ and $q = 1$).

It therefore remains to prove the existence of the required morphism $\exp$. Note first that, by the theory of faithfully flat descent and the preceding uniqueness assertion, it suffices to prove this existence locally on S for the (fpqc) topology. By the usual arguments using finite presentation, one reduces to the case where S is noetherian, then to the case where it is local noetherian. By the preceding remark, one may therefore content oneself with proving the existence of the desired morphism $\exp$ when $S = \text{Spec}(A)$, A local noetherian complete with algebraically closed residue field k . Let then $p_0 : \mathbf{G}_{a,k} \longrightarrow G_k$ be a monomorphism of k -groups normalized by T_k with multiplier $\alpha_0 = \alpha \otimes_A k$ (one exists by 1.1). One knows (1.1 and 1.2) that the corresponding morphism $T_k \cdot_{\alpha_0} \mathbf{G}_{a,k} \longrightarrow G_k$ is an immersion, hence in particular a monomorphism. Let us provisionally admit the following two lemmas:

Lemma 1.6. *Let S be a scheme, G an S -group of finite presentation, T an S -torus, α a character nontrivial on each fiber of T , s_0 a point of S . Let*

$$f : T \cdot_{\alpha} \mathbf{G}_{a,S} \longrightarrow G$$

be a morphism of S -groups such that f_{s_0} is a monomorphism and the restriction of f to T is a monomorphism. There exists an open neighborhood U of s_0 such that $f|_U$ is a monomorphism.

Lemma 1.7. *Let A be a complete local noetherian ring with algebraically closed residue field k , (G, T, α) an elementary A -system, $p_0 : \mathbf{G}_{a,k} \longrightarrow G_k$ a morphism of k -groups normalized by T_k with multiplier $\alpha \otimes_A k$. There exists a morphism of groups $p : \mathbf{G}_{a,A} \longrightarrow G$ normalized by T with multiplier α .*

Let p be the morphism whose existence is asserted by 1.7. Let $f : T \cdot_{\alpha} \mathbf{G}_{a,S} \longrightarrow G$ be the corresponding morphism. It satisfies the hypotheses of 1.6, hence is a monomorphism; in particular p is a monomorphism. One then concludes by Exp. XIX 4.9.

Proof of 1.6. Let $\varepsilon : S \longrightarrow T \cdot_{\alpha} \mathbf{G}_{a,S}$ denote the unit section. Since f is unramified at $\varepsilon(s_0)$, it is so at $\varepsilon(s)$ for all s in an open neighborhood U of s_0 ; $f|_U$ is therefore unramified (Exp. X 3.5),⁽³⁾ hence its kernel $\text{Ker}(f)|_U$ is unramified over U . To prove that $f|_U$ is a monomorphism, it therefore suffices⁽⁴⁾ to prove that $\text{Ker}(f)|_U$ is radicial over U , which is a set-theoretic question. One is therefore reduced to proving:

⁽³⁾**N.D.E.** See also VI_B, 2.5.

⁽⁴⁾**N.D.E.** By EGA IV₄, 17.9.1.

Lemma 1.8. *Let k be an algebraically closed field; let N be an invariant subgroup of $T \cdot_\alpha \mathbf{G}_{a,k}$ (α a nontrivial character of the torus T), étale over k and such that $N \cap T = \{e\}$. Then $N = \{e\}$.*

One has $\text{int}(t')(t, x) = (t, \alpha(t')x)$. If (t, x) is a point of N , with $x \neq 0$, then (t, zx) is also a point of N for $z \in k^*$, and (t, x) is not isolated, hence N is not quasi-finite. One therefore has set-theoretically $N \subset T$, and the proof is finished.

Proof of 1.7. Let $\mathfrak{m}$ be the radical of A and $S_n = \text{Spec}(A/\mathfrak{m}^{n+1})$, $n \geq 0$. Let us first show, by induction on n , that p_0 can be extended for each n to a morphism of S_n -groups

$$p_n : \mathbf{G}_{a,S_n} \longrightarrow G_{S_n}$$

normalized by T_{S_n} with multiplier α_{S_n} , the p_n satisfying in addition $p_{n+1} \times_{S_{n+1}} S_n = p_n$.

Let $H = T \cdot_\alpha \mathbf{G}_{a,S}$. The morphism $H_{S_n} \longrightarrow G_{S_n}$ defined by p_n is denoted f_n . Let us admit the following lemma:

Lemma 1.9. *If (G, T, α) is an elementary k -system, k algebraically closed, and $p : \mathbf{G}_{a,k} \longrightarrow G$ is a monomorphism normalized by T with multiplier α , one has*

$$H^2(T \cdot_\alpha \mathbf{G}_{a,k}, \mathfrak{g}) = 0.$$

(One makes $T \cdot_\alpha \mathbf{G}_{a,k}$ act on $\mathfrak{g}$ through the morphism $T \cdot_\alpha \mathbf{G}_{a,k} \longrightarrow G$ defined by p , and the adjoint representation of G .)

Then, by Exp. III 2.8, f_n will extend to a morphism of S_{n+1} -groups

$$f'_{n+1} : H_{S_{n+1}} \longrightarrow G_{S_{n+1}}.$$

Now f'_{n+1} and the canonical immersion of $T_{S_{n+1}}$ into $G_{S_{n+1}}$ have the same restriction to T_{S_n} . By Exp. III 2.5, there exists an element $g \in G(S_{n+1})$ such that $g \times_{S_{n+1}} S_n = e$ and such that $f_{n+1} = \text{int}(g) \circ f'_{n+1}$ restricts to T_{n+1} by the canonical immersion of T_{n+1} . Let p_{n+1} be the restriction of f_{n+1} to $\mathbf{G}_{a,S_{n+1}}$. It is a morphism normalized by $T_{S_{n+1}}$ with multiplier $\alpha_{S_{n+1}}$, extending p_n .

One has therefore constructed a coherent system (f_n) , and one must now algebraize it. Now one has:

Lemma 1.10. *Let A be a complete local noetherian ring, $\mathfrak{m}$ its maximal ideal, $S = \text{Spec}(A)$, $S_n = \text{Spec}(A/\mathfrak{m}^{n+1})$, T an S -torus, α a nonzero character of T , X an affine S -scheme on which T acts. Let T act on $\mathbf{G}_{a,S}$ through α . Let q be an integer ≥ 0 , and let $(f_n)_{n \geq 0}$ be a coherent system of morphisms*

$$f_n : \mathbf{G}_{a,S_n}^q \longrightarrow X_{S_n}$$

of objects with operators T_{S_n} . There exists a unique morphism of objects with operators T

$$f : \mathbf{G}_{a,S}^q \longrightarrow X$$

inducing the f_n (compare with Exp. IX 7.1).

Corollary 1.11. *If X is a group with operator group T and the f_n are morphisms of groups, f is also one.*

It suffices to apply the uniqueness assertion of the lemma to the two morphisms $\mathbf{G}_{a,S}^{2q} \longrightarrow X$ deduced from f in the usual way.

Proof of 1.10. Suppose T split, which is moreover the case in the application of 1.10 to the proof of 1.5. One knows (Exp. I 4.7.3.1) that $X \longmapsto \mathcal{A}(X)$ gives an equivalence between the category of affine S -schemes equipped with an action of T and the opposite of the category of S -algebras graded of type $M = \text{Hom}_{S\text{-gr.}}(T, \mathbf{G}_{m,S})$.

One therefore has gradings

$$B = \mathcal{A}(X) = \coprod_{m \in M} B_m \quad \text{and} \quad C = \mathcal{A}(\mathbf{G}_{a,S}^q) = \coprod_{m \in M} C_m.$$

One sees at once that each C_m is *free of finite type* over A . (Indeed, $C_m = 0$ if m is not a multiple of α , and if $m = d\alpha$, C_m is isomorphic to the A -module of homogeneous polynomials of degree d in q variables.) Put

$$\begin{aligned}\widehat{B}_m &= \varprojlim_n B_m \otimes_A (A/\mathfrak{m}^{n+1}), \\ \widehat{C}_m &= \varprojlim_n C_m \otimes_A (A/\mathfrak{m}^{n+1}), \\ \widehat{B} &= \prod_{m \in M} \widehat{B}_m, \quad \widehat{C} = \prod_{m \in M} \widehat{C}_m.\end{aligned}$$

One then has canonical morphisms of algebras graded of type M

$$g_B : B \longrightarrow \widehat{B} \quad \text{and} \quad g_C : C \longrightarrow \widehat{C}.$$

It follows from the remark above that g_C is an isomorphism. Giving a coherent system (f_n) as in the statement is equivalent to giving a morphism of graded A -algebras

$$\widehat{F} : \widehat{B} \longrightarrow \widehat{C}.$$

Finding a morphism f as in the statement is equivalent to finding a morphism of graded A -algebras $F : B \longrightarrow C$ making the diagram commute:

$$\begin{array}{ccc} B & \xrightarrow{F} & C \\ g_B \downarrow & & \downarrow g_C \\ \widehat{B} & \xrightarrow{\widehat{F}} & \widehat{C}. \end{array}$$

Since g_C is an isomorphism, the existence and uniqueness of F are immediate. This proves 1.10.

To finish the proof of 1.5, it therefore remains only to prove 1.9.

1.12. — *Proof of 1.9.* One has $\mathfrak{g} = \mathfrak{t} \oplus \mathfrak{g}^\alpha \oplus \mathfrak{g}^{-\alpha}$. As explained in 1.9, consider $\mathfrak{g}$ as a $(T \cdot_\alpha \mathbf{G}_{a,k})$ -module. Clearly $\mathfrak{t} \oplus \mathfrak{g}^\alpha$ is a submodule of $\mathfrak{g}$, the quotient being isomorphic to $\mathfrak{g}^{-\alpha}$ as a k -vector space and even as a T -module. Clearly $\mathbf{G}_{a,k}$ acts trivially on this one-dimensional quotient (since every group morphism from $\mathbf{G}_{a,k}$ to $\mathbf{G}_{m,k}$ is trivial). Similarly, $\mathfrak{g}^\alpha$ is a submodule of $\mathfrak{t} \oplus \mathfrak{g}^\alpha$, the quotient being isomorphic to $\mathfrak{t}$ as a T -module, $\mathbf{G}_{a,k}$ acting trivially on it. To summarize:

Lemma 1.13. *Under the conditions of 1.9, $\mathfrak{g}$ has a composition series as a $(T \cdot_\alpha \mathbf{G}_{a,k})$ -module whose successive quotients are*

$$\mathfrak{g}^{-\alpha}, \quad \mathfrak{t}, \quad \mathfrak{g}^\alpha,$$

considered as $(T \cdot_\alpha \mathbf{G}_{a,k})$ -modules by the projection $T \cdot_\alpha \mathbf{G}_{a,k} \longrightarrow T$.

One is therefore reduced to calculating the cohomology of $T \cdot_\alpha \mathbf{G}_{a,k}$ acting through the projection $T \cdot_\alpha \mathbf{G}_{a,k} \longrightarrow T$ and the character β of T (here $\beta = 0, \alpha$ or $-\alpha$) on $W(k)$.⁽⁵⁾ Denote by $k[x_1, \dots, x_n]$ the polynomial algebra over k in n variables, and by $k_q[x_1, \dots, x_n]$ the subspace of homogeneous polynomials of degree q .

Lemma 1.14. *With the preceding notations, one has $H^n(T \cdot_\alpha \mathbf{G}_{a,k}, k) = H^n(C_{\alpha, \beta}^*)$, where the complex $C_{\alpha, \beta}^*$ is defined by*

$$C_{\alpha, \beta}^n = \begin{cases} k_q[x_1, \dots, x_n] & \text{if } \beta = q\alpha, \text{ with } q \in \mathbf{N}^*; \\ 0 & \text{otherwise,} \end{cases}$$

⁽⁵⁾**N.D.E.** The following sentence has been added.

and

$$\begin{aligned} \delta f(x_1, x_2, \dots, x_{n+1}) &= f(x_2, \dots, x_{n+1}) \\ &+ \sum_{i=1}^n (-1)^i f(x_1, \dots, x_i x_{i+1}, \dots, x_{n+1}) \\ &+ (-1)^{n+1} f(x_1, \dots, x_n). \end{aligned}$$

Indeed, the functor $M \mapsto H^0(T, M)$ is exact on the category of T -modules (and the $H^q(T, -)$ vanish), by Exp. I 5.3.2. It follows, as in the usual case of group cohomology, that $H^n(T \cdot_\alpha \mathbf{G}_{a,k}, k)$ can be calculated as the n th cohomology group of the complex of cochains from $\mathbf{G}_{a,k}$ to k , invariant under T , i.e. satisfying

$$f(\alpha(t)x_1, \dots, \alpha(t)x_n) = \beta(t)f(x_1, \dots, x_n).$$

This indeed gives the announced complex.

To prove 1.9, it therefore suffices to prove that $H^2(C_{\alpha,\beta}^*) = 0$ for $\beta = 0, \alpha, -\alpha$, which is immediate.

Remark 1.15. One can explicitly calculate the groups $H^n(C_{\alpha,\beta}^*)$ for $\beta = q\alpha$ (see M. Lazard, *Lois de groupes et analyseurs*, Annales E.N.S., 1955). In particular, one finds $H^n(C_{\alpha,q\alpha}^*) = 0$ for $n > q$.

Notation 1.16. The image of the canonical immersion

$$W(\mathfrak{g}^{-\alpha}) \times_S T \times_S W(\mathfrak{g}^\alpha) \longrightarrow G$$

will be denoted Ω . It is an open subset of G containing the unit section. The image of

$$W(\mathfrak{g}^{-\alpha}), \quad \text{resp. } W(\mathfrak{g}^\alpha), \quad \text{resp. } W(\mathfrak{g}^{-\alpha}) \times_S T, \quad \text{resp. } T \times_S W(\mathfrak{g}^\alpha)$$

will be denoted⁽⁶⁾

$$U_{-\alpha}, \quad \text{resp. } U_\alpha, \quad \text{resp. } U_{-\alpha} \cdot T, \quad \text{resp. } T \cdot U_\alpha.$$

Then U_α (resp. $U_{-\alpha}$) is a subgroup of G canonically equipped with a vector bundle structure, and one has

$$\text{int}(t)(x) = x^{\alpha(t)} \quad (\text{resp. } x^{-\alpha(t)}),$$

for all $S' \longrightarrow S$, $t \in T(S')$, $x \in U_\alpha(S')$ (resp. $x \in U_{-\alpha}(S')$).

One has canonical isomorphisms

$$T \cdot U_\alpha \simeq T \cdot_\alpha U_\alpha \quad \text{and} \quad T \cdot U_{-\alpha} \simeq T \cdot_{-\alpha} U_{-\alpha}.$$

The open subset Ω is stable under $\text{int}(T)$: one has

$$\text{int}(t')(y \cdot t \cdot x) = y^{-\alpha(t')} \cdot t \cdot x^{\alpha(t')}.$$

Corollary 1.17. *One has $\mathcal{L}ie(U_\alpha/S) = \mathfrak{g}^\alpha$ and $\mathcal{L}ie(U_{-\alpha}/S) = \mathfrak{g}^{-\alpha}$. The isomorphisms*

$$W(\mathfrak{g}^\alpha) \xrightarrow[\sim]{\exp} U_\alpha \quad \text{and} \quad W(\mathfrak{g}^{-\alpha}) \xrightarrow[\sim]{\exp} U_{-\alpha}$$

are those of Exp. XIX 4.2.

Corollary 1.18. *The open subset Ω is relatively schematically dense in G (cf. XVIII, § 1).*

Clear by Exp. XVIII, 1.3.⁽⁷⁾

Corollary 1.19. *The center of G is $\text{Centr}(G) = \text{Ker}(\alpha)$. It is therefore a closed subgroup of G , of multiplicative type and of finite type.*

The second assertion follows from the first by Exp. IX 2.7. Let us therefore prove the latter. The inner automorphism defined by a section of $\text{Ker}(\alpha)$ acts trivially on Ω (last

⁽⁶⁾**N.D.E.** $P_{-\alpha}$ and P_α have been replaced by $U_{-\alpha}$ and U_α .

⁽⁷⁾**N.D.E.** See also EGA IV₃, 11.10.10.

formula of 1.16), hence on G by 1.18. Conversely, if $g \in G(S)$ centralizes G , it centralizes T and U_α , hence is a section of T (Exp. XIX 2.8) annihilating α ; since this also holds after every base change, one indeed has $\text{Centr}(G) = \text{Ker}(\alpha)$.

Corollary 1.20. *For there to exist a monomorphism $p_\alpha : \mathbf{G}_{a,S} \rightarrow G$ normalized by T with multiplier α , it is necessary and sufficient that the $\mathcal{O}_S$ -module $\mathfrak{g}^\alpha$ be free. More precisely, one has a bijection given by*

$$X_\alpha \mapsto (x \mapsto \exp(xX_\alpha)) \quad \text{and} \quad p_\alpha \mapsto \text{Lie}(p_\alpha)$$

between $\Gamma(S, \mathfrak{g}^\alpha)^\times$ and the set of monomorphisms p_α as above (which is also the set of isomorphisms of vector group schemes $\mathbf{G}_{a,S} \xrightarrow{\sim} U_\alpha$).⁽⁸⁾

Corollary 1.21. *The subgroups U_α and $U_{-\alpha}$ of G do not commute on any fiber.*

Indeed, if $(U_\alpha)_s$ and $(U_{-\alpha})_s$ commute, Ω_s is a subgroup of G_s , hence $\Omega_s = G_s$ ⁽⁹⁾ and G_s is solvable, which contradicts the hypothesis that G_s is reductive of semisimple rank 1.

2. Structure of elementary systems

Theorem 2.1. *Let S be a scheme, (G, T, α) an elementary S -system. There exist a morphism of $\mathcal{O}_S$ -modules*

$$\mathfrak{g}^\alpha \otimes_{\mathcal{O}_S} \mathfrak{g}^{-\alpha} \rightarrow \mathcal{O}_S, \quad (X, Y) \mapsto \langle X, Y \rangle,$$

and a morphism of S -groups

$$\alpha^* : \mathbf{G}_{m,S} \rightarrow T$$

such that for every $S' \rightarrow S$ and all $X \in \Gamma(S', \mathfrak{g}^\alpha \otimes \mathcal{O}_{S'})$, $Y \in \Gamma(S', \mathfrak{g}^{-\alpha} \otimes \mathcal{O}_{S'})$ one has the equivalence

$$\exp(X) \cdot \exp(Y) \in \Omega(S') \iff 1 + \langle X, Y \rangle \in \mathbf{G}_m(S'),$$

and under these conditions one has the formula:

$$(F) \quad \boxed{\exp(X) \cdot \exp(Y) = \exp\left(\frac{Y}{1 + \langle X, Y \rangle}\right) \alpha^*(1 + \langle X, Y \rangle) \exp\left(\frac{X}{1 + \langle X, Y \rangle}\right).}$$

Moreover, the morphisms $(X, Y) \mapsto \langle X, Y \rangle$ and α^* are uniquely determined, the first is an isomorphism, hence puts the modules $\mathfrak{g}^\alpha$ and $\mathfrak{g}^{-\alpha}$ in duality, and one has $\alpha \circ \alpha^* = 2$ (squaring in $\mathbf{G}_{m,S}$).

In view of the uniqueness assertions of the theorem, it suffices to give the proof locally on S . One may therefore suppose that $\mathfrak{g}^\alpha$ and $\mathfrak{g}^{-\alpha}$ are free over S . Take then $X \in \Gamma(S, \mathfrak{g}^\alpha)^\times$, $Y \in \Gamma(S, \mathfrak{g}^{-\alpha})^\times$, and put $p_\alpha(x) = \exp(xX)$, $p_{-\alpha}(y) = \exp(yY)$ for $x, y \in \mathbf{G}_a(S')$, $S' \rightarrow S$. By 1.5 and 1.21, it suffices to prove:

Proposition 2.2. *Let S be a scheme, G an S -group, T a torus of G , α a character of T nontrivial on each fiber, $p_\alpha : \mathbf{G}_{a,S} \rightarrow G$ (resp. $p_{-\alpha} : \mathbf{G}_{a,S} \rightarrow G$) a monomorphism of groups normalized by T with multiplier α (resp. $-\alpha$). Suppose that:*

- (i) *The morphism $\mathbf{G}_{a,S} \times_S T \times_S \mathbf{G}_{a,S} \rightarrow G$ defined by $(y, t, x) \mapsto p_{-\alpha}(y)tp_\alpha(x)$ is an open immersion. Denote its image by Ω .*
- (ii) *For every $s \in S$, $(p_\alpha)_s(\mathbf{G}_{a,\kappa(s)})$ and $(p_{-\alpha})_s(\mathbf{G}_{a,\kappa(s)})$ do not commute.*

Then there exist uniquely determined $a \in \mathbf{G}_a(S)$ and $\alpha^ \in \text{Hom}_{S\text{-gr.}}(\mathbf{G}_{m,S}, T)$ with the following properties: for every $S' \rightarrow S$ and all $x, y \in \mathbf{G}_a(S')$, one has*

$$p_\alpha(x)p_{-\alpha}(y) \in \Omega(S') \iff 1 + axy \in \mathbf{G}_m(S'),$$

⁽⁸⁾**N.D.E.** Indeed, on the one hand, $\text{Lie}(\mathbf{G}_{a,S}) = \mathcal{O}_S$, and $\text{Lie}(p_\alpha)$ is an element of $\text{Hom}_{\mathcal{O}_S}(\mathcal{O}_S, \mathfrak{g}^\alpha) = \Gamma(S, \mathfrak{g}^\alpha)$.

⁽⁹⁾**N.D.E.** By 1.18.

and, under this condition, one has the formula

$$p_\alpha(x)p_{-\alpha}(y) = p_{-\alpha}\left(\frac{y}{1+axy}\right)\alpha^*(1+axy)p_\alpha\left(\frac{x}{1+axy}\right).$$

Moreover, a is invertible (i.e. $a \in \mathbf{G}_m(S)$) and $\alpha \circ \alpha^* = 2$.

Proof:

A) Consider the morphism

$$\mathbf{G}_{a,S}^2 \longrightarrow G$$

defined by $(x, y) \mapsto p_\alpha(x)p_{-\alpha}(y)$. Let U be the inverse image of Ω by this morphism. It is an open subset of $\mathbf{G}_{a,S}^2$ containing $0 \times_S \mathbf{G}_{a,S}$ and $\mathbf{G}_{a,S} \times_S 0$. There therefore exist uniquely determined morphisms of S -schemes

$$A : U \longrightarrow \mathbf{G}_{a,S}, \quad C : U \longrightarrow \mathbf{G}_{a,S}, \quad B : U \longrightarrow T$$

satisfying the set-theoretic relation:

$$p_\alpha(u)p_{-\alpha}(v) = p_{-\alpha}(A(u, v))B(u, v)p_\alpha(C(u, v)).$$

One immediately has the relations

$$\begin{aligned} A(0, v) &= v, & A(u, 0) &= 0, & C(u, 0) &= u, & C(0, v) &= 0, \\ B(u, 0) &= B(0, v) = e. \end{aligned}$$

Let S' be a separated S -scheme and $t \in T(S')$ a point of T . Since $\Omega_{S'}$ is stable under $\text{int}(t)$, by the last formula of 1.16, $U_{S'}$ is stable under the automorphism $(x, y) \mapsto (\alpha(t)x, \alpha(t)^{-1}y)$ of $\mathbf{G}_{a,S'}^2$, and one has the relations:

$$\begin{aligned} A(\alpha(t)u, \alpha(t)^{-1}v) &= \alpha(t)^{-1}A(u, v), \\ C(\alpha(t)u, \alpha(t)^{-1}v) &= \alpha(t)C(u, v), \\ B(\alpha(t)u, \alpha(t)^{-1}v) &= B(u, v). \end{aligned}$$

Since α is faithfully flat, one deduces that for every $S' \rightarrow S$ and every $z \in \mathbf{G}_m(S')$, $U_{S'}$ is stable under the transformation $(x, y) \mapsto (zx, z^{-1}y)$ and one has

$$\begin{aligned} A(zu, z^{-1}v) &= z^{-1}A(u, v), & C(zu, z^{-1}v) &= zC(u, v), \\ B(zu, z^{-1}v) &= B(u, v). \end{aligned}$$

Suppose first that v is invertible; putting $z = v$, one deduces that if (u, v) is a section of U , then $(uv, 1)$ is also one, and one has

$$A(uv, 1) = v^{-1}A(u, v), \quad B(uv, 1) = B(u, v).$$

Let then V be the open subset of $\mathbf{G}_{a,S}^2$ defined by⁽¹⁰⁾

$$(u, v) \in V(S') \iff (u, v), (uv, 1) \text{ and } (1, uv) \text{ belong to } U(S').$$

Since U is an open subset of $\mathbf{G}_{a,S}^2$ containing $0 \times_S \mathbf{G}_{a,S}$ and $\mathbf{G}_{a,S} \times_S 0$, V is a neighborhood of the zero section of $\mathbf{G}_{a,S}^2$, and we have just seen that the morphisms

$$\begin{aligned} (u, v) &\mapsto A(u, v) \quad \text{and} \quad (u, v) \mapsto vA(uv, 1), \\ \text{resp. } (u, v) &\mapsto B(u, v) \quad \text{and} \quad (u, v) \mapsto B(uv, 1) \end{aligned}$$

coincide in $V \cap (\mathbf{G}_{a,S} \times_S \mathbf{G}_{m,S})$. Since $\mathbf{G}_{a,S} \times_S \mathbf{G}_{m,S}$ is schematically dense in $\mathbf{G}_{a,S}^2$, these morphisms therefore coincide in V .

One knows that $A(0, 1) = 1$; it follows that there exists an open subset W_1 of $\mathbf{G}_{a,S}$ containing the zero section, such that for every section x of W_1 , $A(x, 1)$ is invertible; putting $A(x, 1)^{-1} = F(x)$, one obtains that if $(u, v) \in V(S')$ ⁽¹¹⁾ and $uv \in W_1(S')$, $S' \rightarrow S$, then

⁽¹⁰⁾**N.D.E.** The condition “ $(1, uv) \in U(S')$ ” has been added.

⁽¹¹⁾**N.D.E.** Here and in what follows, $U(S')$ has been replaced by $V(S')$.

$A(u, v) = vA(uv, 1) = vF(uv)^{-1}$. Reasoning similarly for C , one obtains that there exist an open subset W_2 of $\mathbf{G}_{a,S}$ containing the zero section and an element $E^{(12)}$ of $\mathcal{O}(W_2)^\times$ such that $C(u, v) = uC(1, uv) = uE(uv)^{-1}$ if $(u, v) \in V(S')$ and $uv \in W_2(S')$. Consequently, putting $W = W_1 \cap W_2$, one obtains:

There exist an open subset W of $\mathbf{G}_{a,S}$ containing the zero section, and S -morphisms

$$\begin{aligned} F : W &\longrightarrow \mathbf{G}_{m,S}, & F(0) &= 1, \\ H : W &\longrightarrow T, & H(0) &= e, \\ E : W &\longrightarrow \mathbf{G}_{m,S}, & E(0) &= 1, \end{aligned}$$

such that if $(u, v) \in V(S')$ and $uv \in W(S')$, $S' \longrightarrow S$, one has

$$(+) \quad p_\alpha(u)p_{-\alpha}(v) = p_{-\alpha}(vF(uv)^{-1})H(uv)p_\alpha(uE(uv)^{-1}).$$

B) Let us now use associativity in G to write

$$p_\alpha(u)p_{-\alpha}(v)p_{-\alpha}(w) = p_\alpha(u)p_{-\alpha}(v+w).$$

There exists an open subset L of $\mathbf{G}_{a,S}^3$, containing the unit section, such that $(u, v, w) \in L(S')$ is equivalent to

$$\begin{aligned} (u, v) &\in V(S'), & (uE(uv)^{-1}, w) &\in V(S'), & (u, v+w) &\in V(S'), \\ uv &\in W(S'), & uwE(uv)^{-1} &\in W(S'), & u(v+w) &\in W(S'). \end{aligned}$$

Using formula (+), one writes at once, for $(u, v, w) \in L(S')$, the relations:

$$\begin{aligned} (1) \quad & E(uv + uw) = E(uwE(uv)^{-1})E(uv), \\ (2) \quad & H(uv + uw) = H(uwE(uv)^{-1})H(uv), \\ (3) \quad & (v+w)F(uv + uw)^{-1} = \alpha(H(uv)^{-1})wF(uwE(uv)^{-1})^{-1} \\ & \quad + vF(uv)^{-1}. \end{aligned}$$

It is immediate from the definition of L that $(1, 0, 0) \in L(S)$. Consider therefore

$$L \cap (1 \times_S \mathbf{G}_{a,S} \times_S \mathbf{G}_{a,S}) = 1 \times_S M;$$

M is an open subset of $\mathbf{G}_{a,S}^2$ containing the section $(0, 0)$, and, for $(v, w) \in M(S')$, one has $v, wE(v)^{-1}, v+w \in W(S')$ and

$$\begin{aligned} (1') \quad & E(v+w) = E(wE(v)^{-1})E(v), \\ (2') \quad & H(v+w) = H(wE(v)^{-1})H(v), \\ (3') \quad & (v+w)F(v+w)^{-1} = \alpha(H(v))^{-1}wF(wE(v)^{-1})^{-1} + vF(v)^{-1}. \end{aligned}$$

Finally, consider the morphism from M to $\mathbf{G}_{a,S}^2$ defined set-theoretically by $(v, w) \mapsto (v, wE(v)^{-1})$.⁽¹³⁾ It preserves the section $(0, 0)$ and induces an isomorphism from M onto an open subset N of $\mathbf{G}_{a,S}^2$ containing the zero section (the inverse isomorphism being given by $(x, y) \mapsto (x, yE(x))$).⁽¹⁴⁾ One has therefore proved the following assertion:

There exists an open subset N of $\mathbf{G}_{a,S}^2$ containing the zero section such that if $(x, y) \in N(S')$, then x, y and $x + yE(x)$ ⁽¹⁵⁾ belong to $W(S')$ and:

$$\begin{aligned} (1'') \quad & E(x + yE(x)) = E(x)E(y), \\ (2'') \quad & H(x + yE(x)) = H(x)H(y), \\ (3'') \quad & (x + yE(x))F(x + yE(x))^{-1} = xF(x)^{-1} + \alpha(H(x))^{-1}yE(x)F(y)^{-1}. \end{aligned}$$

⁽¹²⁾**N.D.E.** The element denoted G in the original has been denoted E , since G already denotes the S -group under consideration.

⁽¹³⁾**N.D.E.** The following has been corrected.

⁽¹⁴⁾**N.D.E.** I.e. one has made the “change of variables” $x = v, y = wE(v)^{-1}$, that is, $v = x, w = yE(x)$.

⁽¹⁵⁾**N.D.E.** $yE(x)$ has been corrected to $x + yE(x)$.

C) Reasoning similarly with associativity on the left, one proves the following assertion:⁽¹⁶⁾

There exists an open subset N' of $\mathbf{G}_{a,S}^2$ containing the zero section such that if $(x, y) \in N'(S')$, then x, y and $x + yF(x)$ ⁽¹⁷⁾ belong to $W(S')$, and

$$(4'') \quad F(x + yF(x)) = F(x)F(y),$$

$$(5'') \quad H(x + yF(x)) = H(x)H(y),$$

$$(6'') \quad (x + yF(x))E(x + yF(x))^{-1} = xE(x)^{-1} + \alpha(H(x))^{-1}yF(x)E(y)^{-1}.$$

We are therefore led to solve the “functional equation” (1'').

Lemma 2.3. *Let S be a scheme, W an open subset of $\mathbf{G}_{a,S}$ containing the unit section, $F : W \rightarrow \mathbf{G}_{m,S}$ an S -morphism. Suppose that $F(0) = 1$ and that there exists an open subset N of $\mathbf{G}_{a,S}^2$ containing the zero section such that for $(x, y) \in N(S')$, x, y and $x + yF(x)$ ⁽¹⁷⁾ belong to $W(S')$ and one has:*

$$(\dagger) \quad F(x + yF(x)) = F(x)F(y).$$

(i) *If S is the spectrum of a field k , there exists $a \in k$ such that $F(x) = 1 + ax$.*

(ii) *If $a = F'(0) \in \Gamma(S, \mathcal{O}_S)$ is invertible, then $F(x) = 1 + ax$.*

By the hypotheses, we can differentiate the given equation at $x = 0$ (resp. at $y = 0$), and find that

$$(*) \quad F'(y)(1 + yF'(0)) = F'(0)F(y) \quad \text{for } (0, y) \in N(S'),$$

resp.

$$F'(x)F(x) = F(x)F'(0) \quad \text{for } (x, 0) \in N(S').$$

Since F takes values in $\mathbf{G}_m$, the second relation gives

$$(*') \quad F'(x) = F'(0) \quad \text{for } (x, 0) \in N(S');$$

whence, by the first,

$$F'(0)(1 + yF'(0)) = F'(0)F(y) \quad \text{for } (y, 0), (0, y) \in N(S').$$

If $a = F'(0)$ is invertible, this gives

$$F(y) = 1 + ay,$$

for y a section of an open subset of W containing the unit section, hence schematically dense in W , which proves (ii). This also proves (i) when $F'(0) \neq 0$.

If $F'(0) = 0$, then, by (*'), $F'(x) = 0$ when x is “near 0”, hence $F' = 0$ by schematic density. If k has characteristic 0, F is a rational fraction with zero derivative, hence constant and equal to $F(0) = 1$.

If k has characteristic p and F is not constant,⁽¹⁸⁾ there exist an integer $n > 0$ and a rational fraction $F_1 \in k(X)$ such that $F'_1(X) \neq 0$ and

$$F(X) = F_1(X^{p^n}) = F_1(X)^{p^n}.$$

Substituting in the functional equation, one finds

$$(\dagger_1) \quad F_1(x + yF_1(x)^{p^n}) = F_1(x)F_1(y).$$

Differentiating at $x = 0$, one finds

$$(*_1) \quad F'_1(y) = F'_1(0)F_1(y),$$

and differentiating $(\dagger_1)$ at $y = 0$, one obtains

$$(*'_1) \quad F'_1(x)F_1(x)^{p^n} = F_1(x)F'_1(0).$$

⁽¹⁶⁾**N.D.E.** I.e. one writes the equalities resulting from $p_\alpha(t)p_\alpha(u)p_{-\alpha}(v) = p_\alpha(t+u)p_{-\alpha}(v)$, and puts $v = 1$ and $x = u$, $t = yF(u)$ (i.e. $y = tF(u)^{-1}$).

⁽¹⁷⁾**N.D.E.** $yF(x)$ has been corrected to $x + yF(x)$.

⁽¹⁸⁾**N.D.E.** The following has been corrected.

Since, by hypothesis, $F'_1(X)$ is an invertible element of $k(X)$, one deduces from these two equalities that

$$F_1(X)^{p^n} = 1,$$

hence F_1 is a constant, contradicting the initial hypothesis. This shows that F is constant and equal to $1 = F(0)$.

D) Suppose that S is the spectrum of a field. If $F'(0) = 0$, then $F = 1$. Formula (5'') then gives $H(x+y) = H(x)H(y)$, which shows that H extends to a morphism of groups $\mathbf{G}_{a,S} \rightarrow T$ (Exp. XVIII 2.3), which is necessarily constant with value e . On the other hand, by lemma 2.3, one will also have $E(x) = 1 + bx$ for some $b \in k$. But then (6'') gives, for $(x, y) \in N(S')$,

$$(x+y)E(x+y)^{-1} = xE(x)^{-1} + yE(y)^{-1},$$

hence,⁽¹⁹⁾ by Exp. XVIII 2.3 again, $x \mapsto xE(x)^{-1}$ extends to a morphism of k -groups $\mathbf{G}_{a,k} \rightarrow \mathbf{G}_{a,k}$, hence $x/(1+bx) = cx$ for some $c \in k$, whence $b = 0$ (and $c = 1$).

This shows that F, H, E are constant with values $(1, e, 1)$ in a neighborhood of the unit section, hence everywhere, which by (+) shows that U_α and $U_{-\alpha}$ commute, contrary to hypothesis (ii).

If S is now arbitrary, one has therefore proved that $F'(0)$ vanishes on no fiber, hence is invertible. The same is clearly true of $E'(0)$, which, by lemma 2.3, shows that there exist $a, b \in \mathbf{G}_m(S)$ such that

$$(\diamond_1) \quad F(x) = 1 + ax, \quad E(x) = 1 + bx, \quad \text{for } x \in W(S').$$

E) The rest is now easy. Substituting the preceding results in (3''), one finds

$$y\alpha(H(x))(1+ay) = y(1+ax+ay(1+bx))(1+ax).$$

This formula is valid for every section (x, y) of N . But since $\mathbf{G}_{a,S} \times_S \mathbf{G}_{m,S}$ is schematically dense in $\mathbf{G}_{a,S}^2$, one deduces

$$(1+ay)\alpha(H(x)) = (1+ax+ay(1+bx))(1+ax).$$

Putting $y = 0$ gives $\alpha(H(x)) = (1+ax)^2$. Substituting this in the preceding equality, one finds⁽²⁰⁾

$$a^2xy = abxy.$$

Since $\mathbf{G}_{m,S}$ is schematically dense in $\mathbf{G}_{a,S}$, one deduces $a^2 = ab$, whence, since a is invertible,

$$(\diamond_2) \quad a = b \quad \text{and} \quad \alpha(H(x)) = (1+ax)^2.$$

Since a is invertible, $x \mapsto 1+ax$ is an automorphism of $\mathbf{G}_{a,S}$; one can therefore find an open subset W' of $\mathbf{G}_{a,S}$ containing the section 1 and a morphism

$$P : W' \rightarrow T$$

such that $P(1+ax) = H(x)$.⁽²¹⁾

Substituting in relation (2'), one finds at once, for $(x, y) \in N(S')$,

$$P(1+ax+ay) = P\left(\frac{1+ax+ay}{1+ax}\right)P(1+ax),$$

which proves that there exists an open neighborhood of 1 in $\mathbf{G}_{m,S}$ such that for x and y in this neighborhood one has $P(x)P(y) = P(xy)$. By Exp. XVIII 2.3, there exists a morphism of groups

$$(\diamond_3) \quad \alpha^* : \mathbf{G}_{m,S} \rightarrow T$$

⁽¹⁹⁾**N.D.E.** The following sentence has been added. One can also see by a direct calculation that the preceding equality implies $0 = xyb(2 + (x+y)b)$, whence $0 = b(2 + (x+y)b)$, and finally $b = 0$.

⁽²⁰⁾**N.D.E.** In the preceding three equalities, the original has been corrected by replacing the factor $(1+bx)$ on the right by $1+ax$. Substituting the third equality in the second and taking account of the invertibility of $1+ax$, one indeed obtains $a^2xy = abxy$.

⁽²¹⁾**N.D.E.** I.e. one has made the change of variables $x' = 1+ax$, that is, $x = (x' - 1)/a$.

extending P . Since $\alpha(H(x)) = (1 + ax)^2$ in a neighborhood of the zero section, one has $\alpha(\alpha^*(z)) = z^2$ in a neighborhood of the section 1, hence

$$(\diamond_4) \quad \alpha \circ \alpha^* = 2.$$

F) ⁽²²⁾Collecting results (+) and $(\diamond_1 - \diamond_4)$, one sees that there exist $a \in \mathbf{G}_m(S)$ and $\alpha^* \in \text{Hom}_{S\text{-gr.}}(\mathbf{G}_{m,S}, T)$ such that $\alpha \circ \alpha^* = 2$, and such that, if $(u, v) \in V(S')$ and $uv \in W(S')$, then $1 + auv$ is invertible and

$$p_\alpha(u)p_{-\alpha}(v) = p_{-\alpha}\left(\frac{v}{1 + auv}\right)\alpha^*(1 + auv)p_\alpha\left(\frac{u}{1 + auv}\right).$$

Consider the open subset V' of $\mathbf{G}_{a,S}^2$ defined by “ $1 + auv$ invertible”, i.e. $V' = (\mathbf{G}_{a,S}^2)_f$, where $f(u, v) = 1 + auv$. Both sides of the preceding formula define morphisms from V' to G which coincide in a neighborhood of the zero section, hence coincide in V' . The preceding formula is therefore valid for every section (u, v) of V' . In particular, it follows that $V' \subset U$, where U is the open subset introduced at the beginning of A).

Let us prove that $U = V'$. Returning to the notations of A), one has a morphism

$$A : U \longrightarrow \mathbf{G}_{a,S}$$

which, on V' , is defined by $A(u, v) = v(1 + auv)^{-1}$. To show that $U = V'$, which is a set-theoretic question, one reduces to the case where S is the spectrum of a field k , hence to the following obvious assertion: the domain of definition of the rational map $\mathbf{G}_{a,k}^2 \longrightarrow \mathbf{G}_{a,k}$ defined by the rational fraction $Y/(1 + aXY)$ is the open subset defined by the function $1 + aXY$.

G) One has therefore proved the existence of a and α^* , as well as the two supplementary properties announced. It remains to prove uniqueness. Let therefore a' and $\alpha^{*'} also satisfy the required conditions. If $u, v \in \mathbf{G}_a(S')^2$, one has at once:$

$$1 + auv \text{ invertible} \Rightarrow 1 + a'uv \text{ invertible and } \frac{v}{1 + auv} = \frac{v}{1 + a'uv};$$

one therefore has for every section u of $\mathbf{G}_a(S')$

$$1 + au \text{ invertible} \Rightarrow 1 + au = 1 + a'u,$$

which proves at once that $a = a'$.

With the same notations, one then has

$$1 + au \text{ invertible} \Rightarrow \alpha^*(1 + au) = \alpha^{*'}(1 + au),$$

hence also $\alpha^* = \alpha^{*'}$.

Corollary 2.4. *Let $\exp(Y)t\exp(X)$ and $\exp(Y')t'\exp(X')$ be two elements of $\Omega(S')$. Then their product is in $\Omega(S')$ if and only if $u = 1 + \langle X, Y' \rangle$ is invertible, and one then has*

$$(F') \quad \begin{aligned} \exp(Y)t\exp(X) \cdot \exp(Y')t'\exp(X') &= \exp(Y + u^{-1}\alpha(t)^{-1}Y') \\ &\quad \cdot tt'\alpha^*(u) \cdot \exp(u^{-1}\alpha(t')^{-1}X + X'). \end{aligned}$$

Remark 2.5. One can also write formula (F) of theorem 2.1 without using the morphisms $\exp$. Indeed, transporting by these morphisms the duality $\mathfrak{g}^\alpha \otimes \mathfrak{g}^{-\alpha} \longrightarrow \mathcal{O}_S$, one obtains a canonical pairing of vector bundles:

$$U_\alpha \times_S U_{-\alpha} \longrightarrow \mathbf{G}_{a,S},$$

which we shall again denote by $(x, y) \longmapsto \langle x, y \rangle$. One therefore has

$$\langle \exp X, \exp Y \rangle = \langle X, Y \rangle.$$

⁽²²⁾**N.D.E.** The following has been slightly modified, since an open subset V has already been introduced in A).

If $x \in U_\alpha(S')$, $y \in U_{-\alpha}(S')$ and $1 + \langle x, y \rangle \in \mathbf{G}_m(S')$, one has

$$(F) \quad x \cdot y = y^{(1+\langle x, y \rangle)^{-1}} \cdot \alpha^*(1 + \langle x, y \rangle) \cdot x^{(1+\langle x, y \rangle)^{-1}}.$$

Corollary 2.6. *The pairing*

$$W(\mathfrak{g}^\alpha) \times_S W(\mathfrak{g}^{-\alpha}) \longrightarrow \mathbf{G}_{a,S}$$

defines a pairing of principal bundles under $\mathbf{G}_{m,S}$

$$W(\mathfrak{g}^\alpha)^\times \times_S W(\mathfrak{g}^{-\alpha})^\times \longrightarrow \mathbf{G}_{m,S}.$$

This pairing will be denoted $(X, Y) \mapsto \langle X, Y \rangle$, or more simply $(X, Y) \mapsto XY$.

For every section $X \in \Gamma(S, \mathfrak{g}^\alpha)^\times$, there therefore exists a unique section X^{-1} of $\Gamma(S, \mathfrak{g}^{-\alpha})^\times$ such that $XX^{-1} = 1$. One has $(zX)^{-1} = z^{-1}X^{-1}$. The morphism

$$s : W(\mathfrak{g}^\alpha)^\times \longrightarrow W(\mathfrak{g}^{-\alpha})^\times$$

thus defined is therefore an isomorphism of schemes, compatible with the isomorphism $s : z \mapsto z^{-1}$ on the operator groups.

Definition 2.6.1. One will say that X and $s(X) = X^{-1}$ are *paired*.

Apply corollary 2.4 to $Y = 0 = X'$ and $Y' = aX^{-1}$, $a \in \mathcal{O}_S(S)$. Then $u = 1 + a$ and $u^{-1}Y' = u^{-1}(u - 1)X^{-1} = (1 - u^{-1})X^{-1}$, whence:

Corollary 2.7. *Let $X \in \Gamma(S, \mathfrak{g}^\alpha)^\times$ and $u \in \Gamma(S, \mathcal{O}_S)^\times$. One has*

$$\alpha^*(u) = \exp((u^{-1} - 1)X^{-1}) \exp(X) \exp((u - 1)X^{-1}) \exp(-u^{-1}X).$$

Definition 2.8. The morphism α^* is called the *coroot* associated with the root α .

Remark 2.9. If (G, T, α) is an elementary S -system, $(G, T, -\alpha)$ is also one. One therefore has by theorem 2.1 a duality between $\mathfrak{g}^{-\alpha}$ and $\mathfrak{g}^\alpha$, and a coroot $(-\alpha)^*$. Taking the inverse of formula (F), one proves at once

$$\langle X, Y \rangle = \langle Y, X \rangle, \quad (-\alpha)^* = -\alpha^*.$$

Let us now pass to the Lie algebra of G . The root α and the coroot α^* define the linear forms

$$\mathcal{O}_S \xrightarrow{\bar{\alpha}^*} \mathfrak{t} \xrightarrow{\bar{\alpha}} \mathcal{O}_S.$$

One puts $H_\alpha = \bar{\alpha}^*(1)$. One calls $\bar{\alpha}$ the *infinitesimal root* associated with α , and H_α the corresponding *infinitesimal coroot*.

Lemma 2.10. *Let $S' \longrightarrow S$, and $X, X' \in W(\mathfrak{g}^\alpha)(S')$, $H \in W(\mathfrak{t})(S')$, $Y, Y' \in W(\mathfrak{g}^{-\alpha})(S')$, $t \in T(S')$. One has*

- (1) $\text{Ad}(t)H = H, \quad \text{Ad}(t)X = \alpha(t)X, \quad \text{Ad}(t)Y = \alpha(t)^{-1}Y.$
- (2) $\begin{cases} \text{Ad}(\exp(X))H = H - \bar{\alpha}(H)X, & \text{Ad}(\exp(X))X' = X', \\ \text{Ad}(\exp(X))Y = Y + \langle X, Y \rangle H_\alpha - \langle X, Y \rangle X. \end{cases}$
- (2') $\begin{cases} \text{Ad}(\exp(Y))H = H + \bar{\alpha}(H)Y, & \text{Ad}(\exp(Y))Y' = Y', \\ \text{Ad}(\exp(Y))X = X + \langle X, Y \rangle H_{-\alpha} - \langle X, Y \rangle Y. \end{cases}$
- (3) $[H, X] = \bar{\alpha}(H)X, \quad [H, Y] = -\bar{\alpha}(H)Y, \quad [X, Y] = \langle X, Y \rangle H_\alpha.$
- (4) $H_{-\alpha} = -H_\alpha.$
- (5) $\bar{\alpha}(H_\alpha) = 2.$

The proof of these various formulas is either trivial or an immediate consequence of formula (F) of 2.1.

Corollary 2.11. *Suppose that H_α is nonzero on every fiber (which is in particular the case if 2 is invertible on S , by (5)). Then $X_\alpha \in \Gamma(S, \mathfrak{g}^\alpha)^\times$ and $X_{-\alpha} \in \Gamma(S, \mathfrak{g}^{-\alpha})^\times$ are paired if and only if $[X_\alpha, X_{-\alpha}] = H_\alpha$.*

2.12. — Let (G, T, α) be an elementary S -system. We know (1.19) that the center of G is $\text{Centr}(G) = \text{Ker}(\alpha)$, a group of multiplicative type and of finite type. If Q is a subgroup of multiplicative type of $\text{Centr}(G)$, the quotient G/Q is affine over S (Exp. IX 2.5), smooth over S (Exp. VI_B 9.2), with connected reductive fibers of semisimple rank 1 (Exp. XIX 1.8).

Put $G' = G/Q$; it is a reductive S -group of semisimple rank 1; $T' = T/Q$ is a maximal torus of it. The open subset $U_{-\alpha} \cdot T \cdot U_\alpha$ of G is stable under Q , and one sees at once that the quotient is isomorphic to $U_{-\alpha} \times_S (T/Q) \times_S U_\alpha$. If α' denotes the character of T' induced by α , it follows that the derived morphism of the canonical morphism $G \rightarrow G'$ induces isomorphisms

$$\mathfrak{g}^\alpha \xrightarrow{\sim} \mathfrak{g}'^{\alpha'} \quad \text{and} \quad \mathfrak{g}^{-\alpha} \xrightarrow{\sim} \mathfrak{g}'^{-\alpha'}.$$

In particular, α' is a root of G' with respect to T' . Thus, denoting by α/Q the character $T/Q \rightarrow \mathbf{G}_{m,S}$ induced by α , one has:

Lemma 2.13. *If Q is a subgroup of multiplicative type of $\text{Ker}(\alpha)$, then*

$$(G/Q, T/Q, \alpha/Q)$$

is an elementary system.

Lemma 2.14. *Under the preceding conditions, the following diagrams commute:*

$$\begin{array}{ccccc} W(\mathfrak{g}^\alpha) & \xrightarrow{\text{exp}} & G & \xleftarrow{\text{exp}} & W(\mathfrak{g}^{-\alpha}) \\ \text{can} \downarrow \simeq & & \downarrow \text{can} & & \simeq \downarrow \text{can} \\ W(\mathfrak{g}'^{\alpha'}) & \xrightarrow{\text{exp}} & G' & \xleftarrow{\text{exp}} & W(\mathfrak{g}'^{-\alpha'}) \\ \\ \mathfrak{g}^\alpha \otimes \mathfrak{g}^{-\alpha} & \xrightarrow{\sim} & \mathcal{O}_S & & \\ \text{can} \downarrow \simeq & & \downarrow \text{id} & & \\ \mathfrak{g}'^{\alpha'} \otimes \mathfrak{g}'^{-\alpha'} & \xrightarrow{\sim} & \mathcal{O}_S & & \\ \\ \begin{array}{ccc} & T & \\ \alpha^* \nearrow & \downarrow \text{can} & \searrow \alpha \\ \mathbf{G}_{m,S} & & \mathbf{G}_{m,S} \\ \alpha'^* \searrow & \downarrow & \nearrow \alpha' \\ & T' & \end{array} & & \end{array}.$$

3. The Weyl group

Notation 3.0. ⁽²³⁾If (G, T, α) is an elementary S -system, one puts

$$N = \underline{\text{Norm}}_G(T), \quad W = \underline{\text{Norm}}_G(T)/T$$

(cf. Exp. XIX 6.3); N is a closed subgroup of G , smooth over S . One denotes by $N^\times = N - T$ the open subscheme of N induced on the complement of T .⁽²⁴⁾ Let R be the (unique) maximal torus of $\text{Ker}(\alpha)$, and T' the image of $\alpha^* : \mathbf{G}_{m,S} \rightarrow T$, which is a one-dimensional subtorus of T .

⁽²³⁾**N.D.E.** The numbering 3.0 has been added for subsequent references.

⁽²⁴⁾**N.D.E.** Q has been replaced by the more suggestive notation $N^\times$.

The morphism

$$T' \times_S R \longrightarrow T$$

induced by the product in T is surjective (hence faithfully flat); indeed, one reduces to checking this on geometric fibers, and it follows at once from the formula $\alpha \circ \alpha^* = 2$.

Theorem 3.1. *With the preceding notations:*

- (i) W is isomorphic to the constant group $(\mathbf{Z}/2\mathbf{Z})_S$.
- (ii) $N^\times$ is a locally trivial principal homogeneous bundle under T , on the left by the law $(t, q) \mapsto tq$ (resp. on the right by the law $(q, t) \mapsto qt$).
- (iii) One has the formula

$$\text{int}(w)t = t \cdot \alpha^*(\alpha(t)^{-1})$$

for $w \in N^\times(S')$, $t \in T(S')$, $S' \longrightarrow S$. In the decomposition $T_{S'} = T'_{S'} \cdot R_{S'}$, $\text{int}(w)$ induces the identity on $R_{S'}$ and reflection on $T'_{S'}$. One has the relations

$$\alpha \circ \text{int}(w) = \alpha^{-1}, \quad \text{int}(w) \circ \alpha^* = (\alpha^*)^{-1}.$$

- (iv) For $X \in W(\mathfrak{g}^\alpha)^\times(S')$, $S' \longrightarrow S$, put

$$w_\alpha(X) = \exp(X) \exp(-X^{-1}) \exp(X).$$

Then $w_\alpha(X) \in N^\times(S')$, and the morphism $w_\alpha : W(\mathfrak{g}^\alpha)^\times \longrightarrow N^\times$ thus defined satisfies

$$w_\alpha(zX) = \alpha^*(z)w_\alpha(X) = w_\alpha(X)\alpha^*(z)^{-1},$$

for $z \in \mathbf{G}_m(S')$, $X \in W(\mathfrak{g}^\alpha)^\times(S')$, $S' \longrightarrow S$.

- (v) ⁽²⁵⁾For $X, Y \in W(\mathfrak{g}^\alpha)^\times(S')$, $S' \longrightarrow S$, with the notations of 2.6, one has

$$w_\alpha(X)w_\alpha(Y) = \alpha^\vee(-XY^{-1}).$$

In particular,

$$w_\alpha(X)^2 = \alpha^*(-1) \in {}_2T(S) \cap \underline{\text{Centr}}(G)(S),$$

$$w_\alpha(X)^{-1} = w_\alpha(-X) = \alpha^*(-1)w_\alpha(X).$$

- (vi) If one similarly defines, for $Y \in W(\mathfrak{g}^{-\alpha})^\times(S')$,

$$w_{-\alpha}(Y) = \exp(Y) \exp(-Y^{-1}) \exp(Y),$$

one has (in addition to formulas analogous to the preceding ones)

$$w_{-\alpha}(X^{-1}) = w_\alpha(X)^{-1} = w_\alpha(-X),$$

$$w_\alpha(X)w_{-\alpha}(Y) = \alpha^*(XY).$$

Proof. (i) was already seen in Exp. XIX 2.4; it follows at once that $N^\times$ is indeed a principal homogeneous bundle under T for the laws defined in (ii); its local triviality (for the Zariski topology) follows, in particular, from (iv).

Let us prove (iii); if $w \in N^\times(S)$, clearly $\alpha \circ \text{int}(w)$ is a root of G with respect to T , hence is locally equal to α or $-\alpha$; since on each fiber it is $-\alpha$ (*Bible*, 12-05, proof of the cor. to prop. 1), one has $\alpha \circ \text{int}(w) = -\alpha$. By transport of structure, one deduces

$$-\alpha^* = \text{int}(w)^{-1} \circ \alpha^* = \text{int}(w) \circ \alpha^*,$$

since $\text{int}(w)^2 = \text{int}(w^2)$ and w^2 is a section of T . Thus $\text{int}(w)$ induces reflection on T' ; since R is central, $\text{int}(w)$ induces the identity on R . The formula of (iii) defines a morphism $T \longrightarrow T$ with the same properties, hence coincides with $\text{int}(w)$.

Let us prove (iv). One has successively

$$\begin{aligned} w_\alpha(X)tw_\alpha(X)^{-1} &= \exp(X) \exp(-X^{-1}) \exp(X)t \exp(-X) \exp(X^{-1}) \exp(-X) \\ &= \exp(X) \exp(-X^{-1}) \exp(X - \alpha(t)X) \exp(\alpha(t)^{-1}X^{-1}) \exp(-\alpha(t)X)t. \end{aligned}$$

⁽²⁵⁾**N.D.E.** The first formula of the original has been corrected.

Applying formula (F), one has

$$\begin{aligned} \exp(-X^{-1}) \exp((1 - \alpha(t))X) &= \exp((\alpha(t)^{-1} - 1)X) \alpha^*(\alpha(t)^{-1}) \\ &\quad \cdot \exp(-\alpha(t)^{-1}X^{-1}). \end{aligned}$$

Substituting in the preceding relation, one finds

$$\begin{aligned} \text{int}(w_\alpha(X))t &= \exp(\alpha(t)^{-1}X) \alpha^*(\alpha(t)^{-1}) \exp(-\alpha(t)X)t \\ &= \exp(aX) \alpha^*(\alpha(t)^{-1})t, \end{aligned}$$

where

$$a = \alpha(t)^{-1} - (\alpha \circ \alpha^*)(\alpha(t)^{-1})\alpha(t),$$

but $\alpha \circ \alpha^* = 2$, which gives at once $a = 0$ and $w_\alpha(X) \in N^\times(S')$.

Let us now prove the second assertion of (iv). One has⁽²⁶⁾

$$\begin{aligned} \alpha^*(z)w_\alpha(X) &= \exp(z^2X) \exp(-z^{-2}X^{-1}) \exp(z^2X) \alpha^*(z) \\ &= \exp(zX) \exp((z^2 - z)X) \exp(-z^{-2}X^{-1}) \exp(z^2X) \alpha^*(z) \\ &= \exp(zX) \exp(-z^{-1}X^{-1}) \alpha^*(z)^{-1} \exp((z^3 - z^2)X) \exp(z^2X) \alpha^*(z) \\ &= \exp(zX) \exp(-z^{-1}X^{-1}) \exp(zX) = w_\alpha(zX). \end{aligned}$$

Let us prove (v). By the preceding result, the first formula of (v) follows at once from the second; let us prove the latter:

$$\begin{aligned} w_\alpha(X)^2 &= \exp(X) \exp(-X^{-1}) \exp(2X) \exp(-X^{-1}) \exp(X) \\ &= \exp(X) \exp(-X^{-1}) \exp(X^{-1}) \alpha^*(-1) \exp(-2X) \exp(X) \\ &= \exp(X) \alpha^*(-1) \exp(-X) = \alpha^*(-1), \end{aligned}$$

since $\alpha(\alpha^*(-1)) = (-1)^2 = 1$, which proves that $\alpha^*(-1) \in \text{Centr}(G)(S)$.

Finally, let us prove (vi). The first assertion is a particular case of the second; let us prove the latter. Both sides of this formula define morphisms from $W(\mathfrak{g}^\alpha)^\times \times_S W(\mathfrak{g}^{-\alpha})^\times$ to G . To prove that they coincide, it suffices to do so on an open subset nonempty on each fiber (Exp. XVIII 1.4); it therefore suffices to check the relation when $1 + XY$ is invertible. One then has successively:

$$\begin{aligned} w_\alpha(X)w_{-\alpha}(Y) &= \exp(X) \exp(-X^{-1}) \exp(X) \exp(Y) \exp(-Y^{-1}) \exp(Y) \\ &= \exp(X) \exp(-X^{-1}) \exp\left(\frac{Y}{1+XY}\right) \alpha^*(1+XY) \\ &\quad \exp\left(\frac{X}{1+XY}\right) \exp(-Y^{-1}) \exp(Y) \\ &= \exp(X) \exp\left(\frac{-X^{-1}}{1+XY}\right) \alpha^*(1+XY) \exp\left(\frac{-Y^{-1}}{1+XY}\right) \exp(Y) \\ &= \exp(-X^{-2}Y^{-1}) \alpha^*\left(\frac{XY}{1+XY}\right) \exp(X+Y^{-1}) \\ &\quad \alpha^*(1+XY) \exp\left(\frac{-Y^{-1}}{1+XY}\right) \exp(Y) \\ &= \exp(-X^{-2}Y^{-1}) \alpha^*(XY) \exp\left(\frac{Y^{-1}+X}{(1+XY)^2}\right) \exp\left(\frac{-Y^{-1}}{1+XY}\right) \exp(Y) \\ &= \alpha^*(XY) \exp(-Y) \exp(Y) = \alpha^*(XY). \end{aligned}$$

⁽²⁶⁾**N.D.E.** The first equality follows from 1.5 (i), which, combined with $\alpha \circ \alpha^* = 2$, gives the formulas

$$\begin{aligned} \alpha^*(z) \exp(X) \alpha^*(z)^{-1} &= \exp(z^2X), \\ \alpha^*(z) \exp(X^{-1}) \alpha^*(z)^{-1} &= \exp(z^{-2}X), \end{aligned}$$

the third equality follows from formula (F), and the fourth from (†), again. Finally, a similar calculation shows that $w_\alpha(X) \alpha^*(z^{-1}) = w_\alpha(zX)$.

Corollary 3.2. *Let $n \in \mathbf{Z}$, $n \neq 0$. For every $w \in G(S)$, the following conditions are equivalent:*

- (i) $w \in N^\times(S)$,
- (ii) *one has $\text{int}(w) \circ n\alpha^* = -n\alpha^*$ (recall that $(n\alpha^*)(z) = \alpha^*(z)^n$).*

One has (i) $\Rightarrow$ (ii) (assertion (iii) of theorem 3.1); conversely, one may suppose that $N^\times$ has a section, and one is reduced to proving:

Lemma 3.3. *One has $\underline{\text{Centr}}_G(n\alpha^*) = T$ for $n \neq 0$.*

Indeed, the image T' of $n\alpha^*$ is a subtorus of G . It follows (Exp. XIX 2.8) that $\underline{\text{Centr}}_G(n\alpha^*)$ is a reductive subgroup of G , containing T . Since on each fiber one has $\underline{\text{Centr}}_{G_{\bar{s}}}(n\alpha_{\bar{s}}^*) \neq G_{\bar{s}}$, one has $\underline{\text{Centr}}_{G_{\bar{s}}}(n\alpha_{\bar{s}}^*) = T_{\bar{s}}$ (Exp. XIX 1.6.3⁽²⁷⁾), hence $\underline{\text{Centr}}_G(n\alpha^*) = T$, since these are smooth subgroups of G .

Remark 3.4. The construction of w_α and the fact that $w_\alpha(X)$ normalizes T rely only on formula (F). In particular, if G is an S -group satisfying the conditions of 2.2, $\underline{\text{Norm}}_G(T)$ is different from T on each fiber. It follows that if G is an affine S -group with connected fibers satisfying the conditions of 2.2, it is reductive of semisimple rank 1. Indeed, it is smooth in a neighborhood of the unit section, hence smooth, and one can apply the criterion of Exp. XIX 1.11.

3.5. — Before stating the following theorem, let us make a few remarks. As usual, we identify $\mathfrak{g}^{-\alpha}$ with $(\mathfrak{g}^\alpha)^{\otimes -1}$. Similarly, we shall identify $\underline{\text{Hom}}_{\mathcal{O}_S}(\mathfrak{g}^{-\alpha}, \mathfrak{g}^\alpha)$ with $(\mathfrak{g}^\alpha)^{\otimes 2}$, and hence

$$\underline{\text{Isom}}_{\mathcal{O}_S\text{-mod.}}(\mathbf{W}(\mathfrak{g}^{-\alpha}), \mathbf{W}(\mathfrak{g}^\alpha)) \simeq \mathbf{W}((\mathfrak{g}^\alpha)^{\otimes 2})^\times.$$

If $w \in N^\times(S)$, $\text{Ad}(w)$ interchanges $\mathfrak{g}^\alpha$ and $\mathfrak{g}^{-\alpha}$ (3.1, (iii)), hence defines an isomorphism

$$a_\alpha(w) : \mathfrak{g}^{-\alpha} \xrightarrow{\sim} \mathfrak{g}^\alpha,$$

which we shall therefore identify with a section $a_\alpha(w) \in \Gamma(S, (\mathfrak{g}^\alpha)^{\otimes 2})^\times$. This construction is compatible with base change and therefore defines a morphism

$$a_\alpha : N^\times \longrightarrow \mathbf{W}((\mathfrak{g}^\alpha)^{\otimes 2})^\times,$$

such that $a_\alpha(w)Y = \text{Ad}(w)Y$ for all $w \in N^\times(S')$, $Y \in \Gamma(S', \mathfrak{g}^{-\alpha})^\times$, $S' \longrightarrow S$.

Theorem 3.6.

- (i) *One has*

$$\text{int}(w) \exp(Y) = \exp(a_\alpha(w)Y)$$

for every $S' \longrightarrow S$ and all $w \in N^\times(S')$, $Y \in \mathbf{W}(\mathfrak{g}^{-\alpha})(S')$.

- (ii) *One has*

$$a_\alpha(tw) = \alpha(t)a_\alpha(w), \quad a_\alpha(wt) = \alpha(t)^{-1}a_\alpha(w).$$

- (iii) *If one similarly defines $a_{-\alpha} : N^\times \longrightarrow \mathbf{W}((\mathfrak{g}^{-\alpha})^{\otimes 2})^\times$, one has*

$$a_{-\alpha}(w) = a_\alpha(w)^{-1}.$$

(28)

- (iv) *For every $X \in \mathbf{W}(\mathfrak{g}^\alpha)^\times(S')$, $S' \longrightarrow S$, one has*

$$a_\alpha(w_\alpha(X)) = -X^2.$$

⁽²⁷⁾**N.D.E.** The hypothesis $\underline{\text{Centr}}_{G_{\bar{s}}}(n\alpha_{\bar{s}}^*) \neq G_{\bar{s}}$ implies that $\dim \underline{\text{Centr}}_{G_{\bar{s}}}(n\alpha_{\bar{s}}^*) - \dim T_{\bar{s}} < 2$, whereas this difference is even, by loc. cit.

⁽²⁸⁾**N.D.E.** I.e. $a_{-\alpha}(w)$ and $a_{-\alpha}(w)$ are paired, cf. 2.6.1.

Assertion (i) is trivial by the characterization of the morphisms $\exp$ given in 1.5. Assertion (ii) is immediate, as is (iii). Let us prove (iv): let $X \in \Gamma(S', \mathfrak{g}^\alpha)^\times$, $Z \in \Gamma(S', \mathfrak{g}^\alpha)$; by definition, one has⁽²⁹⁾

$$\begin{aligned} a_\alpha(w_\alpha(X))^{-1}(Z) &= \text{Ad}(w_\alpha(X))(Z) \\ &= \text{Ad}(\exp(X)) \text{Ad}(\exp(-X^{-1})) \text{Ad}(\exp(X))(Z). \end{aligned}$$

Applying formulas (2') and (2) of lemma 2.10, as well as the equalities $H_{-\alpha} = -H_\alpha$, $\bar{\alpha}(H_\alpha) = 2$ (loc. cit. (4) and (5)) and $\langle X, X^{-1} \rangle = 1$ (2.6), one obtains that the right-hand term equals, successively:

$$\begin{aligned} \text{Ad}(\exp(X)) \text{Ad}(\exp(-X^{-1}))(Z) &= \text{Ad}(\exp(X))(Z + \langle X^{-1}, Z \rangle (H_\alpha - X^{-1})) \\ &= Z + \langle X^{-1}, Z \rangle (H_\alpha - 2X - X^{-1} - H_\alpha + X) \\ &= Z - \langle X^{-1}, Z \rangle X - \langle X^{-1}, Z \rangle X^{-1}. \end{aligned}$$

But $Z = \langle X^{-1}, Z \rangle X$ and $\langle X^{-1}, Z \rangle X^{-1} = X^{-2}Z$, hence this shows that $a_\alpha(w_\alpha(X))^{-1} = -X^{-2}$, whence $a_\alpha(w_\alpha(X)) = -X^2$.

Corollary 3.7. *In particular, one has*

$$\text{int}(w_\alpha(X)) \exp(X) = \exp(-X^{-1}),$$

whence (by the definition of $w_\alpha(X)$):

$$w_\alpha(X) \exp(X) w_\alpha(X)^{-1} = \exp(-X) w_\alpha(X) \exp(-X),$$

i.e. by an immediate calculation,

$$(w_\alpha(X) \exp(X))^3 = e.$$

Corollary 3.8. *Let $X \in \Gamma(S, \mathfrak{g}^\alpha)^\times$ and $n \in \mathbf{Z}$, $n \neq 0$. Then $w_\alpha(X)$ is the unique section $w \in G(S)$ satisfying*

- (i) $\text{int}(w) \circ n\alpha^* = -n\alpha^*$.
- (ii) $(w \exp(X))^3 = e$.

One knows that $w_\alpha(X)$ indeed satisfies these conditions. Conversely, let $w \in G(S)$ satisfy (i) and (ii). By 3.2 and 3.1 (ii), one knows that there exists $t \in T(S)$ such that $w = w_\alpha(X)t$. Put $u = \exp(X)$. One then has

$$wuw^{-1} = w_\alpha(X)t \exp(X)t^{-1}w_\alpha(X)^{-1} = \exp(-\alpha(t)X^{-1}),$$

and, on the other hand,

$$\begin{aligned} u^{-1}wu^{-1} &= \exp(-X)w_\alpha(X)t \exp(-X) \\ &= \exp(-X)w_\alpha(X) \exp(-X) \exp(X - \alpha(t)X)t \\ &= \exp(-X^{-1}) \exp(X - \alpha(t)X)t = \exp(-X^{-1})t \exp(H). \end{aligned}$$

Now $(wu)^3 = e \iff wuw^{-1} = u^{-1}wu^{-1}$; comparing the two decompositions of this element on $U_{-\alpha} \cdot T \cdot U_\alpha$, one obtains $t = e$.

Remark 3.9. One can summarize a number of the results of this section by the following diagram of principal homogeneous bundles (on the left):

$$\begin{array}{ccc} \mathbf{W}(\mathfrak{g}^\alpha)^\times & \xrightarrow{w_\alpha} & N^\times \xrightarrow{a_\alpha} \mathbf{W}((\mathfrak{g}^\alpha)^{\otimes 2})^\times \\ & & \\ \mathbf{G}_{\mathbf{m},S} & \xrightarrow{\alpha^*} & T \xrightarrow{\alpha} \mathbf{G}_{\mathbf{m},S}. \end{array}$$

⁽²⁹⁾**N.D.E.** The following has been expanded.

Note that a_α is faithfully flat (since α is) and that w_α is a monomorphism if and only if α^* is a monomorphism. We leave it to the reader to write the corresponding diagrams for the structures of principal bundles on the right, as well as diagrams of the same kind for the root $-\alpha$, and to study the relations between these various diagrams.

Lemma 3.10. *Let S be a scheme, q an integer > 0 such that $x \mapsto x^q$ defines an endomorphism of $\mathbf{G}_{a,S}$, (G, T, α) and (G', T', α') two elementary S -systems, $f : G \rightarrow G'$ a morphism of S -groups. Let*

$$h : (\mathfrak{g}^\alpha)^{\otimes q} \xrightarrow{\sim} \mathfrak{g}'^{\alpha'}$$

be an isomorphism of $\mathcal{O}_S$ -modules, and

$$h^\vee : (\mathfrak{g}^{-\alpha})^{\otimes q} \xrightarrow{\sim} \mathfrak{g}'^{-\alpha'}$$

the contragredient isomorphism. For every $S' \rightarrow S$ and every $X \in W(\mathfrak{g}^\alpha)(S')$, suppose that:

$$f(\exp(X)) = \exp(h(X^q)).$$

Then the following conditions are equivalent:

- (i) $f(\alpha^*(z)) = \alpha'^*(z)^q$.
- (ii) $f(w_\alpha(Z)) = w_{\alpha'}(h(Z^q))$.
- (iii) $f(\exp(Y)) = \exp(h^\vee(Y^q))$.

(Each condition is to be read: for every $S' \rightarrow S$ and every $z \in \mathbf{G}_m(S')$, $Z \in W(\mathfrak{g}^\alpha)^\times(S')$, $Y \in W(\mathfrak{g}^{-\alpha})(S')$, one has ...)

Indeed, (i) $\Rightarrow$ (ii) by 3.8, (ii) $\Rightarrow$ (iii) by 3.7, (iii) $\Rightarrow$ (i) by 2.7.

Proposition 3.11. *Let S be a scheme, $q \in \mathbf{Z}$, $q > 0$, such that $x \mapsto x^q$ defines an endomorphism of $\mathbf{G}_{a,S}$, (G, T, α) and (G', T', α') two elementary S -systems, $f : G \rightarrow G'$ a morphism of S -groups. The following conditions on f are equivalent:*

- (i) *The restriction of f to T factors as a morphism $f_T : T \rightarrow T'$ making the diagram commute:*

$$\begin{array}{ccccc} \mathbf{G}_{m,S} & \xrightarrow{\alpha^*} & T & \xrightarrow{\alpha} & \mathbf{G}_{m,S} \\ q \downarrow & & \downarrow f_T & & \downarrow q \\ \mathbf{G}_{m,S} & \xrightarrow{\alpha'^*} & T' & \xrightarrow{\alpha'} & \mathbf{G}_{m,S} \end{array}$$

- (ii) *There exists a (unique) isomorphism of $\mathcal{O}_S$ -modules*

$$h : (\mathfrak{g}^\alpha)^{\otimes q} \rightarrow \mathfrak{g}'^{\alpha'}$$

such that

$$f(\exp(X)) = \exp(h(X^q)), \quad f(\exp(Y)) = \exp(h^\vee(Y^q))$$

for all $X \in W(\mathfrak{g}^\alpha)(S')$, $Y \in W(\mathfrak{g}^{-\alpha})(S')$, $S' \rightarrow S$ (it follows that f also satisfies the equivalent conditions of 3.10).

One has (ii) $\Rightarrow$ (i). Indeed, by 3.10, condition (ii) implies $f \circ \alpha^* = q\alpha'^*$, hence, by 3.3, $f|_T$ factors through T' . It remains to prove that $\alpha'(f(t)) = \alpha(t)^q$, which follows at once from the fact that f induces a morphism of groups $T \cdot U_\alpha \rightarrow T' \cdot U_{\alpha'}$.

Let us prove (i) $\Rightarrow$ (ii). Let $X \in \Gamma(S, \mathfrak{g}^\alpha)$, $Y \in \Gamma(S, \mathfrak{g}^{-\alpha})$. Put $p_+(x) = f(\exp(xX))$ and $p_-(x) = f(\exp(yY))$; these are morphisms of groups

$$p_+, p_- : \mathbf{G}_{a,S} \rightarrow G.$$

Now one has

$$\begin{aligned} \text{int}(\alpha'^*(z))^q(p_+(x)) &= \text{int}(f_T(\alpha^*(z)))(f(\exp(xX))) \\ &= f(\text{int}(\alpha^*(z))(\exp(xX))) \\ &= f(\exp(z^2xX)) = p_+(z^2x). \end{aligned}$$

Applying lemma 1.2 (with $Q = \alpha'^*(\mathbf{G}_{m,S})$), one deduces that there exists a section $X' \in \Gamma(S, \mathfrak{g}^{\alpha'})$ such that

$$f(\exp(xX)) = p_+(x) = \exp(x^q X').$$

Similarly, there exists a section $Y' \in \Gamma(S, \mathfrak{g}^{-\alpha'})$ such that

$$f(\exp(yY)) = \exp(y^q Y').$$

Writing now that f is a morphism of groups, hence respects formula (F), one obtains at once

$$X^q Y^q = (XY)^q = X' Y'.$$

One easily concludes that $X^q \mapsto X'$ and $Y^q \mapsto Y'$ define isomorphisms h and $h^\vee$ as announced.

Proposition 3.12. *Let (G, T, α) be an elementary S -system, $w \in N^\times(S)$; put*

$$\Omega_0 = \Omega \cap \text{int}(w^{-1})(\Omega).$$

Let d be the function on Ω defined by

$$d(\exp(Y) \cdot t \cdot \exp(X)) = \alpha(t)^{-1} + XY.$$

Then $\Omega_0 = \Omega_d$, and for $\exp(Y) \cdot t \cdot \exp(X) \in \Omega_0(S')$ one has the following formula (put $z = d(\exp(Y) \cdot t \cdot \exp(X))$):

$$(\star) \quad \begin{aligned} \text{int}(w)(\exp(Y) \cdot t \cdot \exp(X)) &= \exp(z^{-1} a_\alpha(w)^{-1} X) \\ &\quad \cdot t \alpha^*(z) \cdot \exp(z^{-1} a_\alpha(w) Y). \end{aligned}$$

Moreover, $d \circ \text{int}(w) = d^{-1}$.

Indeed, one has at once⁽³⁰⁾

$$\begin{aligned} \text{int}(w)(\exp(Y) \cdot t \cdot \exp(X)) &= \exp(a_\alpha(w) Y) \cdot t \alpha^*(\alpha(t)^{-1}) \cdot \exp(a_\alpha(w)^{-1} X) \\ &= \exp(a_\alpha(w) Y) \cdot \exp(\alpha(t) a_\alpha(w)^{-1} X) \cdot t \alpha^*(\alpha(t)^{-1}). \end{aligned}$$

By 2.1, this is a section of Ω if and only if $1 + \alpha(t)XY$ is invertible, which indeed proves $\Omega_0 = \Omega_d$; then applying formula (F) of loc. cit., one deduces by an immediate calculation the announced formula $(\star)$. Finally, it follows from $(\star)$ that one has

$$\begin{aligned} (d \circ \text{int}(w))(\exp(Y) \cdot t \cdot \exp(X)) &= \alpha(t \alpha^*(z))^{-1} + z^{-2} XY \\ &= z^{-2} (\alpha(t)^{-1} + XY) = z^{-1}, \end{aligned}$$

whence the last assertion.

N. B. One will note that the function d is independent of the choice of w .

4. The isomorphism theorem

Theorem 4.1. *Let S be a scheme, $q \in \mathbf{Z}$, $q > 0$, such that $x \mapsto x^q$ is an endomorphism of $\mathbf{G}_{a,S}$, (G, T, α) and (G', T', α') two elementary S -systems. Let*

$$h : (\mathfrak{g}^\alpha)^{\otimes q} \longrightarrow \mathfrak{g}'^{\alpha'} \quad \text{and} \quad h^\vee : (\mathfrak{g}^{-\alpha})^{\otimes q} \longrightarrow \mathfrak{g}'^{-\alpha'}$$

⁽³⁰⁾**N.D.E.** The original has been corrected by interchanging $a_\alpha(w)$ and $a_\alpha(w)^{-1}$, and the proof of $d \circ \text{int}(w) = d^{-1}$ has been expanded.

be two mutually contragredient isomorphisms. Let $f_T : T \rightarrow T'$ be a morphism of S -groups making the diagram commute:

$$\begin{array}{ccc} \mathbf{G}_{m,S} & \xrightarrow{q} & \mathbf{G}_{m,S} \\ \alpha^* \downarrow & & \downarrow \alpha'^* \\ T & \xrightarrow{f_T} & T' \\ \alpha \downarrow & & \downarrow \alpha' \\ \mathbf{G}_{m,S} & \xrightarrow{q} & \mathbf{G}_{m,S}. \end{array}$$

There exists a unique morphism of S -groups $f : G \rightarrow G'$ extending f_T and satisfying

$$f(\exp(X)) = \exp(h(X^q))$$

for every $X \in W(\mathfrak{g}^\alpha)(S')$, $S' \rightarrow S$. Moreover, this morphism also satisfies

$$f(\exp(Y)) = \exp(h^\vee(Y^q)) \quad \text{and} \quad f(w_\alpha(Z)) = w_\alpha(h(Z^q)),$$

for every $S' \rightarrow S$ and all $Y \in \Gamma(S', \mathfrak{g}^{-\alpha})$, $Z \in \Gamma(S', \mathfrak{g}^\alpha)^\times$.

If $f : G \rightarrow G'$ extends f_T , then $f \circ \alpha^* = (\alpha'^*)^q$. If in addition f satisfies the second condition, it also satisfies the other two by 3.10. It follows that f is determined on Ω by the relation

$$f(\exp(Y)t\exp(X)) = \exp(h^\vee(Y^q))f_T(t)\exp(h(X^q)).$$

Since Ω is schematically dense in G , this already proves the uniqueness of f . To prove its existence, it suffices, by Exp. XVIII 2.3, to prove that the preceding formula defines a “generically multiplicative” morphism from Ω to G' . Now, by 2.4, this amounts to checking that $\alpha' \circ f = \alpha^q$, which follows from the fact that f extends f_T .

Scholium 4.2. One can also interpret 4.1 as follows: consider the category $\mathcal{E}$ of elementary S -systems and the category $\mathcal{D}$ of pairs

$$\left(\mathbf{G}_{m,S} \xrightarrow{\alpha^*} T \xrightarrow{\alpha} \mathbf{G}_{m,S}, \mathcal{L} \right),$$

where T is a torus, α and α^* morphisms of groups such that $\alpha \circ \alpha^* = 2$, and $\mathcal{L}$ an invertible $\mathcal{O}_S$ -module (the reader will specify the morphisms of the two categories considered). One defines a functor $\mathcal{E} \rightarrow \mathcal{D}$ by

$$(G, T, \alpha) \mapsto \left(\mathbf{G}_{m,S} \xrightarrow{\alpha^*} T \xrightarrow{\alpha} \mathbf{G}_{m,S}, \mathfrak{g}^\alpha \right).$$

The preceding theorem says that this functor is fully faithful. It is in fact an equivalence of categories, as will be seen in the next section. One already has:

Corollary 4.3. If $q = 1$ and f_T is an isomorphism, then f is an isomorphism.

Corollary 4.4. If $q = 1$ and f_T is faithfully flat with kernel Q (cf. Exp. IX 2.7), then f is faithfully flat (quasi-compact) with kernel Q , hence identifies G' with G/Q .

Indeed, if f_T is faithfully flat with kernel Q , then

$$Q = \text{Ker}(f_T) \subset \text{Ker}(f_T \circ \alpha') = \text{Ker}(\alpha).$$

Introducing the elementary S -system $(G/Q, T/Q, r/Q)$ of 2.13, one is reduced by 2.14 to proving that f/Q induces an isomorphism from G/Q onto G' , which follows at once from 4.3.

5. Examples of elementary systems, applications

5.1. — Let S be a scheme, $\mathcal{L}$ an invertible $\mathcal{O}_S$ -module. Consider the group $G_{\mathcal{L}}$ over S defined by

$$G_{\mathcal{L}}(S') = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \left| \begin{array}{l} a, d \in \mathbf{G}_a(S'), \quad b \in W(\mathcal{L})(S'), \\ c \in W(\mathcal{L}^{-1})(S'), \quad ad - bc \in \mathbf{G}_m(S') \end{array} \right. \right\},$$

equipped with the usual matrix multiplication law. It is locally isomorphic to $\mathrm{GL}_{2,S}$. It is therefore a group S -scheme, affine and smooth over S , with connected fibers.

Remark. Let $\mathcal{L}'$ and $\mathcal{L}''$ be two invertible sheaves on S such that $\mathcal{L} = \mathcal{L}' \otimes \mathcal{L}''^{-1}$.⁽³¹⁾ Then one has an isomorphism of S -groups:

$$G_{\mathcal{L}} \xrightarrow{\sim} \mathrm{GL}(\mathcal{L}' \oplus \mathcal{L}'')$$

defined as follows: if x (resp. y) is a section of $\mathcal{L}'$ (resp. $\mathcal{L}''$) over an open subset V of S , one has

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} ax + by \\ cx + dy \end{pmatrix}.$$

5.2. — One denotes by $S_{\mathcal{L}}$ the closed subgroup of $G_{\mathcal{L}}$ defined by $ad - bc = 1$. It is also a group S -scheme, affine and smooth over S , with connected fibers (isomorphic to $\mathrm{SL}(\mathcal{L}' \oplus \mathcal{L}'')$ by the preceding isomorphism).

Similarly, consider the morphism $\mathbf{G}_{m,S} \rightarrow G_{\mathcal{L}}$ defined by $z \mapsto \begin{pmatrix} z & 0 \\ 0 & z \end{pmatrix}$. It is a central monomorphism; passing to the quotient, one deduces a group $P_{\mathcal{L}}$, smooth and affine over S , with connected fibers (cf. Exp. VIII 5.7).

One can see that, passing to the quotient from the isomorphism of the preceding remark, $P_{\mathcal{L}}$ is identified with the group of automorphisms of the projective bundle $\mathbf{P}(\mathcal{L}' \oplus \mathcal{L}'')$ (cf. EGA, II 4.2.7). One denotes by i and p the canonical morphisms

$$S_{\mathcal{L}} \xrightarrow{i} G_{\mathcal{L}} \xrightarrow{p} P_{\mathcal{L}};$$

i is a closed immersion, p is faithfully flat and affine.

5.3. — Consider the morphisms of groups

$$\begin{aligned} t_G : \mathbf{G}_{m,S}^2 &\rightarrow G_{\mathcal{L}}, & t_G(z, z') &= \begin{pmatrix} z & 0 \\ 0 & z' \end{pmatrix}; \\ t_S : \mathbf{G}_{m,S} &\rightarrow S_{\mathcal{L}}, & t_S(z) &= \begin{pmatrix} z & 0 \\ 0 & z^{-1} \end{pmatrix}; \\ t_P : \mathbf{G}_{m,S} &\rightarrow P_{\mathcal{L}}, & t_P(z) &= p(t_G(z, 1)). \end{aligned}$$

These are monomorphisms of groups defining in each group a (split) torus of relative codimension 2. For every $s \in S$, let

$$X \in \Gamma(\bar{s}, \mathcal{L} \otimes \bar{s})^{\times};$$

then the section $\begin{pmatrix} 0 & X \\ X^{-1} & 0 \end{pmatrix}$ of $G_{\mathcal{L},\bar{s}}$ normalizes $t_G(\mathbf{G}_{m,\bar{s}}^2)$ and does not centralize it; one then concludes from Exp. XIX 1.6 that $G_{\mathcal{L}}$ is *reductive*, of *semisimple rank 1*, with *maximal torus* $t_G(\mathbf{G}_{m,S}^2)$.

One reasons similarly for $S_{\mathcal{L}}$ and $P_{\mathcal{L}}$, and sees that $S_{\mathcal{L}}$ (resp. $P_{\mathcal{L}}$) is *reductive*, of *semisimple rank 1*, with maximal torus $t_S(\mathbf{G}_{m,S})$ (resp. $t_P(\mathbf{G}_{m,S})$).

5.4. — Reasoning as usual, one immediately determines the Lie algebra of these various groups and the adjoint action of the chosen maximal torus. Let us do so for $G_{\mathcal{L}}$; this is immediate by Exp. II 4.8: $\mathrm{Lie}(G_{\mathcal{L}}/S)$ is the Lie algebra of the matrices below:

$$\mathrm{Lie}(G_{\mathcal{L}}/S) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \left| \begin{array}{l} a \text{ and } d \text{ sections of } \mathcal{O}_S, \quad b \text{ a section of } \mathcal{L}, \\ c \text{ a section of } \mathcal{L}^{-1} \end{array} \right. \right\}$$

⁽³¹⁾**N.D.E.** $\mathcal{L}'' \otimes \mathcal{L}'^{-1}$ has been corrected to $\mathcal{L}' \otimes \mathcal{L}''^{-1}$, and the following sentence has been expanded.

with the usual bracket; one has

$$\mathrm{Ad}(t_G(z, z')) \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} a & zz'^{-1}b \\ z'z^{-1}c & d \end{pmatrix}.$$

Put $\mathcal{L}ie(G_{\mathcal{L}}/S) = \mathfrak{g}$. Let $\alpha_G : t_G(\mathbf{G}_{\mathbf{m},S}^2) \rightarrow \mathbf{G}_{\mathbf{m},S}$ be the character defined by

$$\alpha_G(t_G(z, z')) = zz'^{-1}.$$

One sees at once from the preceding relation that α_G is a root of $G_{\mathcal{L}}$ with respect to $t_G(\mathbf{G}_{\mathbf{m},S}^2)$, and that the morphism

$$u : \mathcal{L} \rightarrow \mathfrak{g} \quad (\text{resp. } u_- : \mathcal{L}^{-1} \rightarrow \mathfrak{g})$$

defined by $u(X) = \begin{pmatrix} 0 & X \\ 0 & 0 \end{pmatrix}$ (resp. $u_-(X) = \begin{pmatrix} 0 & 0 \\ X & 0 \end{pmatrix}$) is an isomorphism from $\mathcal{L}$ onto $\mathfrak{g}^{\alpha_G}$ (resp. from $\mathcal{L}^{-1}$ onto $\mathfrak{g}^{-\alpha_G}$).

One has therefore proved that $(G, t_G(\mathbf{G}_{\mathbf{m},S}^2), \alpha_G)$ is an elementary S -system.

Putting similarly

$$\alpha_S(t_S(z)) = z^2, \quad \alpha_P(t_P(z)) = z,$$

one proves that $(S_{\mathcal{L}}, t_S(\mathbf{G}_{\mathbf{m},S}), \alpha_S)$ and $(P_{\mathcal{L}}, t_P(\mathbf{G}_{\mathbf{m},S}), \alpha_P)$ are elementary systems, and defines isomorphisms of $\mathcal{L}$ (resp. $\mathcal{L}^{-1}$) with the corresponding direct summands of the Lie algebras of $S_{\mathcal{L}}$ and $P_{\mathcal{L}}$.

5.5. — Put $\exp \begin{pmatrix} 0 & X \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} 1 & X \\ 0 & 1 \end{pmatrix}$. One has thus defined a morphism

$$W(\mathfrak{g}^{\alpha_G}) \rightarrow G_{\mathcal{L}}$$

which induces on Lie algebras the canonical morphism, hence is the unique morphism of this type (1.5). Similarly, put $\exp \begin{pmatrix} 0 & 0 \\ Y & 0 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ Y & 1 \end{pmatrix}$. Calculating formula (F) explicitly, one finds

$$\left\langle \begin{pmatrix} 0 & X \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ Y & 0 \end{pmatrix} \right\rangle = XY, \quad \alpha_G^*(z) = \begin{pmatrix} z & 0 \\ 0 & 1/z \end{pmatrix} = t_G(z, z^{-1}).$$

⁽³²⁾The open subset $N^\times = N_G^\times$ (defined before 3.1) is:

$$N_G^\times(S') = \left\{ \begin{pmatrix} 0 & P \\ Q & 0 \end{pmatrix} \mid P \in W(\mathfrak{g}^\alpha)^\times(S'), Q \in W(\mathfrak{g}^{-\alpha})^\times(S') \right\},$$

the morphism w_{α_G} (cf. 3.1 (iv)) is given, for every $X \in W(\mathfrak{g}^\alpha)^\times(S')$, by

$$w_{\alpha_G}(X) = \begin{pmatrix} 0 & X \\ -X^{-1} & 0 \end{pmatrix};$$

the morphism a_{α_G} (cf. 3.5) is given by:

$$\text{if } w = \begin{pmatrix} 0 & P \\ Q & 0 \end{pmatrix} \in N_G^\times(S'), \quad \text{then } a_{\alpha_G}(w) = PQ^{-1} \in W((\mathfrak{g}^\alpha)^{\otimes 2})^\times(S'),$$

i.e. for every $Y \in W(\mathfrak{g}^{-\alpha})^\times(S')$, one has $a_{\alpha_G}(w)(Y) = PQ^{-1}Y \in W(\mathfrak{g}^\alpha)^\times(S')$.

5.6. — We leave it to the reader to perform the same calculations in $S_{\mathcal{L}}$ and $P_{\mathcal{L}}$. One finds the same duality formula and the coroots

$$\alpha_S^*(z) = t_S(z), \quad \alpha_P^*(z) = t_P(z^2).$$

Let p_T be the morphism induced by $p : G_{\mathcal{L}} \rightarrow P_{\mathcal{L}}$ on $t_S(\mathbf{G}_{\mathbf{m},S})$, i.e.

$$p_T(t_S(z)) = t_P(z^2).$$

⁽³²⁾**N.D.E.** The following has been expanded.

One therefore has the commutative diagram:⁽³³⁾

$$\begin{array}{ccccc}
 & & \mathbf{G}_{m,S} & & \\
 & \swarrow \text{id} & \searrow \alpha_S^* & \swarrow \alpha_P^* & \searrow 2 \\
 \mathbf{G}_{m,S} & \xrightarrow{t_S} & t_S(\mathbf{G}_{m,S}) & \xrightarrow{p_T} & t_P(\mathbf{G}_{m,S}) \xrightarrow{t_P} \mathbf{G}_{m,S} \\
 & \searrow 2 & \swarrow \alpha_S & \swarrow \alpha_P & \searrow \text{id} \\
 & & \mathbf{G}_{m,S} & &
 \end{array}$$

One recognizes in the central part the commutative diagram of 4.1⁽³⁴⁾ relative to the canonical morphism $p \circ i : S_{\mathcal{L}} \rightarrow P_{\mathcal{L}}$, which induces a morphism of the preceding elementary S -systems.

5.7. — Let now (G, T, α) be any elementary S -system. Consider the commutative diagram:

$$\begin{array}{ccccc}
 & & \mathbf{G}_{m,S} & & \\
 & \swarrow \text{id} & \downarrow \alpha^* & \swarrow 2 & \\
 \mathbf{G}_{m,S} & \xrightarrow{\alpha^*} & T & \xrightarrow{\alpha} & \mathbf{G}_{m,S} \\
 & \searrow 2 & \downarrow \alpha & \swarrow \text{id} & \\
 & & \mathbf{G}_{m,S} & &
 \end{array}$$

Combining the preceding two diagrams, one obtains a commutative diagram:

$$\begin{array}{ccccc}
 & & \mathbf{G}_{m,S} & & \\
 & \swarrow \alpha_S^* & \downarrow \alpha^* & \swarrow \alpha_P^* & \\
 t_S(\mathbf{G}_{m,S}) & \xrightarrow{\alpha^* \circ t_S^{-1}} & T & \xrightarrow{t_P \circ \alpha} & t_P(\mathbf{G}_{m,S}) \\
 & \searrow \alpha_S & \downarrow \alpha & \swarrow \alpha_P & \\
 & & \mathbf{G}_{m,S} & &
 \end{array}$$

Using 4.1, one therefore has:

Proposition 5.8. *Let S be a scheme, (G, T, α) an elementary S -system. Put $\mathcal{L} = \mathfrak{g}^\alpha$ (and hence $\mathcal{L}^{-1} = \mathfrak{g}^{-\alpha}$).*

(i) *There exists a unique morphism of groups $f : S_{\mathcal{L}} \rightarrow G$ satisfying the following equivalent conditions:*

$$\begin{aligned}
 (a) \quad & f \begin{pmatrix} z & 0 \\ 0 & z^{-1} \end{pmatrix} = \alpha^*(z), & f \begin{pmatrix} 1 & X \\ 0 & 1 \end{pmatrix} = \exp(X); \\
 (b) \quad & f \begin{pmatrix} 1 & X \\ 0 & 1 \end{pmatrix} = \exp(X), & f \begin{pmatrix} 1 & 0 \\ Y & 1 \end{pmatrix} = \exp(Y); \\
 (c) \quad & f \begin{pmatrix} 1 & X \\ 0 & 1 \end{pmatrix} = \exp(X), & f \begin{pmatrix} 0 & X \\ -X^{-1} & 0 \end{pmatrix} = w_\alpha(X).
 \end{aligned}$$

(ii) *There exists a unique morphism of groups $g : G \rightarrow P_{\mathcal{L}}$ satisfying*

$$g(t) = \begin{pmatrix} \alpha(t) & 0 \\ 0 & 1 \end{pmatrix}, \quad g(\exp(X)) = p \begin{pmatrix} 1 & X \\ 0 & 1 \end{pmatrix}.$$

⁽³³⁾**N.D.E.** Where t_S and t_P are isomorphisms.

⁽³⁴⁾**N.D.E.** With $q = 1$.

Moreover, one has

$$g(\exp(Y)) = p \begin{pmatrix} 1 & 0 \\ Y & 1 \end{pmatrix}, \quad g(w_\alpha(X)) = p \begin{pmatrix} 0 & X \\ -X^{-1} & 0 \end{pmatrix}.$$

The morphism g is faithfully flat quasi-compact with kernel $\text{Ker}(\alpha) = \underline{\text{Centr}}(G)$, and $g \circ f$ is the canonical morphism $S_{\mathcal{L}} \rightarrow P_{\mathcal{L}}$.

Note that conditions (b) of (i) give an explicit description of the duality between $\mathfrak{g}^\alpha$ and $\mathfrak{g}^{-\alpha}$.

Corollary 5.9. *Let (G, T, α) be an elementary S -system. The subgroups $T \cdot U_\alpha$, $T \cdot U_{-\alpha}$, U_α and $U_{-\alpha}$ are closed.*

Since U_α is a closed subgroup scheme of $T \cdot U_\alpha$, it suffices to check this for the latter. By Noether's theorem (Exp. IV 5.3.1 and 6.4.1), it suffices to prove that $(T \cdot U_\alpha)/\text{Ker}(\alpha)$ is a closed subgroup of $G/\text{Ker}(\alpha)$. By 5.8, one is therefore reduced to proving that the subgroup of $P_{\mathcal{L}}$ (or of $G_{\mathcal{L}}$, which amounts to the same thing by another application of Noether's theorem) defined by $c = 0$ is closed, which is trivial.

Consequently, the morphisms $\exp$ of theorem 1.5 (i) are closed immersions.

N. B. The corollary also follows from the fact that $T \cdot U_\alpha$ and $T \cdot U_{-\alpha}$ are “Borel subgroups” of G (cf. Exp. XII 7.10).

5.10. — Let $\mathcal{L}$ be an invertible $\mathcal{O}_S$ -module and

$$\mathbf{G}_{m,S} \xrightarrow{\alpha^*} T \xrightarrow{\alpha} \mathbf{G}_{m,S}$$

a diagram of groups⁽³⁵⁾ such that $\alpha \circ \alpha^* = 2$. Let R be the maximal torus of $\text{Ker}(\alpha)$ and $K = \alpha^{*-1}(R)$. Then K is a subgroup of multiplicative type of $\mathbf{G}_{m,S}$; by $\alpha \circ \alpha^* = 2$, it is even a subgroup of $\mu_{2,S}$. In particular, the morphism

$$K \rightarrow S_{\mathcal{L}}, \quad z \mapsto \begin{pmatrix} z & 0 \\ 0 & z^{-1} \end{pmatrix}$$

is central. One therefore has a central monomorphism of groups:

$$K \rightarrow R \times S_{\mathcal{L}}, \quad z \mapsto \left(\alpha^*(z), \begin{pmatrix} z & 0 \\ 0 & z^{-1} \end{pmatrix} \right).$$

Consider the group $G = (R \times S_{\mathcal{L}})/K$ obtained by passing to the quotient. It is a group affine and smooth over S , with connected fibers. It is immediate that the sequence

$$1 \rightarrow K \rightarrow R \times t_S(\mathbf{G}_{m,S}) \rightarrow T \rightarrow 1$$

where $u(x, t_S(z)) = x\alpha^*(z)$ is exact. The image of $R \times t_S(\mathbf{G}_{m,S})$ in G is therefore a torus T' isomorphic to T . One now shows without difficulty that if α' is the character of T' deduced from α by the preceding isomorphism, (G, T', α') is an elementary S -system, that $\mathfrak{g}^{\alpha'}$ is isomorphic to $\mathcal{L}$, and that α'^* is obtained from α^* by the isomorphism $T \xrightarrow{\sim} T'$. One has therefore constructed an elementary S -system (G, T', α') such that the corresponding object $(\mathbf{G}_{m,S} \xrightarrow{\alpha'^*} T' \xrightarrow{\alpha'} \mathbf{G}_{m,S}, \mathfrak{g}^{\alpha'})$ of the category $\mathcal{D}$ defined in 4.2 is isomorphic to $(\mathbf{G}_{m,S} \xrightarrow{\alpha^*} T \xrightarrow{\alpha} \mathbf{G}_{m,S}, \mathcal{L})$. One has therefore proved:

Theorem 5.11. *With the notations of 4.2, the functor*

$$(G, T, \alpha) \mapsto (\mathbf{G}_{m,S} \xrightarrow{\alpha^*} T \xrightarrow{\alpha} \mathbf{G}_{m,S}, \mathfrak{g}^\alpha)$$

is an equivalence of categories between $\mathcal{E}$ and $\mathcal{D}$.

6. Generators and relations for an elementary system

⁽³⁵⁾N.D.E. T being a torus.

6.1. — Let S be a scheme, (G, T, α) an elementary S -system. Let $X \in W(\mathfrak{g}^\alpha)^\times(S)$ and $u = \exp(X)$; we saw in 3.8 that the element $w = w_\alpha(X)$ satisfies, in particular, the relation

$$(wu)^3 = e.$$

⁽³⁶⁾Let s_α denote the automorphism of T induced by $\text{int}(w)$; by theorem 3.1 (iii), for every $S' \rightarrow S$ and $t \in T(S')$, one has

$$s_\alpha(t) = \text{int}(w)(t) = t \cdot \alpha^*(\alpha(t)^{-1}).$$

Theorem 6.2. *Let H be a sheaf of S -groups for (fppf). Let*

$$f_T : T \rightarrow H, \quad f_\alpha : U_\alpha \rightarrow H$$

be morphisms of groups and $h \in H(S)$ a section of H . For there to exist a morphism of groups (necessarily unique)

$$f : G \rightarrow H$$

extending f_T and f_α and satisfying $f(w) = h$, it is necessary and sufficient that the following conditions hold:

(i) *For every $S' \rightarrow S$, every $t \in T(S')$ and every $x \in U_\alpha(S')$, one has*

$$(1) \quad f_T(t)f_\alpha(x)f_T(t)^{-1} = f_\alpha(txt^{-1}) = f_\alpha(x^{\alpha(t)}).$$

(in other words, f_T and f_α extend to a morphism of groups from the semidirect product $T \cdot U_\alpha$ to H).

(ii) *For every $S' \rightarrow S$ and every $t \in T(S')$, one has*

$$(2) \quad hf_T(t)h^{-1} = f_T(s_\alpha(t)) = f_T(t \cdot \alpha^*(\alpha(t)^{-1})).$$

(iii) *One has the two relations in $H(S)$:*

$$(3) \quad h^2 = f_T(\alpha^*(-1)),$$

$$(4) \quad (hf_\alpha(u))^3 = e.$$

Proof. Let us write U_α and $U_{-\alpha}$ additively, and their vector structure multiplicatively. If f satisfies the conditions of the statement, one necessarily has, for every $y \in U_{-\alpha}(S')$,

$$f(y) = f(w^{-1}yw^{-1}w) = hf_\alpha(w^{-1}yw)h^{-1}.$$

Let therefore $f_{-\alpha} : U_{-\alpha} \rightarrow H$ be the morphism defined by

$$(*_1) \quad f_{-\alpha}(y) = hf_\alpha(w^{-1}yw)h^{-1}.$$

It is a morphism of groups. On the other hand, f is determined on the big cell Ω by

$$f(ytx) = f_{-\alpha}(y)f_T(t)f_\alpha(x).$$

This shows the uniqueness of f ; since the conditions of the statement are manifestly necessary, let us show that they are sufficient.

By (4), one has

$$hf_\alpha(u)h^{-1}h^2 = f_\alpha(-u)h^{-1}f_\alpha(-u).$$

Now, by (3) and (1), $h^2 = h^{-2}$ commutes with $f_\alpha(-u)$, which gives

$$hf_\alpha(u)h^{-1} = f_\alpha(-u)hf_\alpha(-u).$$

But, by definition, $hf_\alpha(u)h^{-1} = f_{-\alpha}(wuw^{-1})$; by 3.7, since $u = \exp(X)$ and $w = w_\alpha(X)$, one has

$$(*_2) \quad wuw^{-1} = -\tilde{u},$$

where $\tilde{u}$ denotes the element paired with u . One therefore obtains:

$$(*_3) \quad f_{-\alpha}(-\tilde{u}) = f_\alpha(-u)hf_\alpha(-u).$$

⁽³⁶⁾**N.D.E.** The following sentence has been added.

Let now t be a section of T over a variable $S' \rightarrow S$. Let $\text{int}(f_T(t))$ act on the preceding formula. On the left-hand side, one obtains⁽³⁷⁾

$$\begin{aligned} f_T(t)f_{-\alpha}(-\tilde{u})f_T(t)^{-1} &= f_T(t)hf_{\alpha}(u)h^{-1}f_T(t)^{-1} \\ &= h(h^{-1}f_T(t)h)f_{\alpha}(u)(h^{-1}f_T(t)^{-1}h)h^{-1} \\ &= hf_T(s_{\alpha}(t))f_{\alpha}(u)f_T(s_{\alpha}(t))^{-1}h^{-1} \\ &= hf_{\alpha}(\alpha(s_{\alpha}(t))u)h^{-1} \end{aligned}$$

by (2) and (1); then, since $s_{\alpha}(t) = t \cdot \alpha^*(\alpha(t)^{-1})$ and $\alpha \circ \alpha^* = 2$, this equals

$$hf_{\alpha}(\alpha(t)^{-1}u)h^{-1}.$$

Finally, by $(*_1)$ and $(*_2)$, one has

$$hf_{\alpha}(\alpha(t)^{-1}u)h^{-1} = f_{-\alpha}(\alpha(t)^{-1}uwuw^{-1}) = f_{-\alpha}(-\alpha(t)^{-1}\tilde{u}).$$

The right-hand side of $(*_3)$ gives

$$f_{\alpha}(-\alpha(t)u) \cdot f_T(t)hf_T(t)^{-1}h^{-1} \cdot h \cdot f_{\alpha}(-\alpha(t)u),$$

and since $hf_T(t)^{-1}h^{-1} = f_T(s_{\alpha}(t^{-1})) = f_T(t \cdot \alpha^*(\alpha(t)))$, this equals

$$f_{\alpha}(-\alpha(t)u) \cdot f_T(\alpha^*(\alpha(t))) \cdot h \cdot f_{\alpha}(-\alpha(t)u).$$

Comparing the two expressions obtained, one obtains

$$f_{-\alpha}(-\alpha(t)^{-1}\tilde{u}) = f_{\alpha}(-\alpha(t)u) \cdot f_T(\alpha^*(\alpha(t))) \cdot h \cdot f_{\alpha}(-\alpha(t)u).$$

Since $\alpha : T \rightarrow \mathbf{G}_{m,S}$ is faithfully flat and H is a separated presheaf, one may replace $-\alpha(t)^{-1}$ by an arbitrary section of $\mathbf{G}_{m,S}$ and obtains:

Lemma 6.2.1. *For every $z \in \mathbf{G}_m(S')$, $S' \rightarrow S$, one has*

$$f_{-\alpha}(z\tilde{u}) = f_{\alpha}(z^{-1}u) \cdot f_T(\alpha^*(-z^{-1})) \cdot h \cdot f_{\alpha}(z^{-1}u).$$

Let now $x, y \in \mathbf{G}_a(S')$, $S' \rightarrow S$; suppose y and $(1 + xy)$ invertible. First applying the lemma with $z = y$, one obtains⁽³⁸⁾

$$f_{\alpha}(xu)f_{-\alpha}(y\tilde{u}) = f_{\alpha}((x + y^{-1})u) \cdot f_T(\alpha(-y^{-1})) \cdot h \cdot f_{\alpha}(y^{-1}u).$$

Now $x + y^{-1} = y^{-1}(1 + xy)$. Applying the lemma with $z = y/(1 + xy)$, one finds

$$\begin{aligned} f_{\alpha}((x + y^{-1})u) &= f_{-\alpha}\left(\frac{y}{1 + xy}\tilde{u}\right) f_{\alpha}(-(x + y^{-1})u) \\ &\quad \cdot h^{-1} \cdot f_T\left(\alpha^*\left(\frac{-y}{1 + xy}\right)\right). \end{aligned}$$

Substituting this in the preceding equality, one obtains

$$\begin{aligned} f_{\alpha}(xu)f_{-\alpha}(y\tilde{u}) &= f_{-\alpha}\left(\frac{y}{1 + xy}\tilde{u}\right) f_{\alpha}(-(x + y^{-1})u) \\ &\quad \cdot h^{-1} \cdot f_T(\alpha^*(1 + xy)^{-1}) \cdot h \cdot f_{\alpha}(y^{-1}u). \end{aligned}$$

Since $h^{-1}f_T(t)h = f_T(s_{\alpha}(t))$ by (2) (note that $s_{\alpha}^2 = \text{id}$), and since $s_{\alpha} \circ \alpha^* = -\alpha^*$ (cf. 6.2.1), this equals

$$f_{-\alpha}\left(\frac{y}{1 + xy}\tilde{u}\right) f_{\alpha}(-(x + y^{-1})u) \cdot f_T(\alpha^*(1 + xy)) \cdot f_{\alpha}(y^{-1}u).$$

Finally, since for all $x' \in U_{\alpha}(S')$ and $z \in \mathbf{G}_m(S')$, one has

$$f_{\alpha}(x')f_T(\alpha^*(z)) = f_T(\alpha^*(z))f_{\alpha}(z^{-2}x'),$$

⁽³⁷⁾**N.D.E.** The following has been corrected.

⁽³⁸⁾**N.D.E.** The following calculations have been expanded.

one obtains

$$\begin{aligned} f_\alpha(xu)f_{-\alpha}(y\tilde{u}) &= f_{-\alpha}\left(\frac{y}{1+xy}\tilde{u}\right) \cdot f_T(\alpha^*(1+xy)) \cdot f_\alpha\left(\left(\frac{-y^{-1}(1+xy)}{(1+xy)^2} + y^{-1}\right)u\right) \\ &= f_{-\alpha}\left(\frac{y}{1+xy}\tilde{u}\right) \cdot f_T(\alpha^*(1+xy)) \cdot f_\alpha\left(\frac{x}{1+xy}u\right). \end{aligned}$$

One has therefore proved:

Lemma 6.2.2. *Let $S' \rightarrow S$. If $a \in U_\alpha(S')$, $b \in U_{-\alpha}^\times(S')$, and $1+ab \in \mathbf{G}_m(S')$, one has*

$$f_\alpha(a)f_{-\alpha}(b) = f_{-\alpha}\left(\frac{b}{1+ab}\right) f_T(\alpha^*(1+ab)) f_\alpha\left(\frac{a}{1+ab}\right).$$

By schematic density, this formula remains valid when $b \in U_{-\alpha}(S')$, $1+ab$ still being invertible. Consider then the morphism

$$f : \Omega \rightarrow H$$

defined by $f(ytx) = f_{-\alpha}(y)f_T(t)f_\alpha(x)$.

It follows at once from 6.2.2, condition 6.2 (i), and formula (F') of 2.4, that if $g, g' \in \Omega(S')$ and $gg' \in \Omega(S')$, one has $f(gg') = f(g)f(g')$. By Exp. XVIII 2.3 (iii) and 2.4,⁽³⁹⁾ there therefore exists a morphism of groups $G \rightarrow H$ extending f . Denote it also by f ; it answers the question; indeed, it suffices to prove that $f(w_\alpha) = h$. Now $w_\alpha = u \cdot (-\tilde{u}) \cdot u$, whence, by $(*)_3$:⁽⁴⁰⁾

$$f(w_\alpha) = f_\alpha(u)f_{-\alpha}(-\tilde{u})f_\alpha(u) = h.$$

Remark 6.3. We shall supplement these results in Exp. XXIII 3.5.

⁽³⁹⁾**N.D.E.** Note that each geometric fiber of G is connected, for example by 1.1.

⁽⁴⁰⁾**N.D.E.** The original has been simplified by invoking $(*)_3$.