

Introduction to Data Mining

03 - Statistical Testing

Benjamin Paaßen

WS 2023/2024, Bielefeld University

- ▶ First sheet is out TODAY with deadline friday in 11 days (Nov 03, noon, i.e. 12:00)

- ▶ First sheet is out TODAY with deadline friday in 11 days (Nov 03, noon, i.e. 12:00)
- ▶ ChatGPT/external help policy: language models are permitted IF you tell us where you used it (indicate in your solution) and you could still present the solution yourself; ideally: link the chatlog

- ▶ First sheet is out TODAY with deadline friday in 11 days (Nov 03, noon, i.e. 12:00)
- ▶ ChatGPT/external help policy: language models are permitted IF you tell us where you used it (indicate in your solution) and you could still present the solution yourself; ideally: link the chatlog Why?

- ▶ First sheet is out TODAY with deadline friday in 11 days (Nov 03, noon, i.e. 12:00)
- ▶ ChatGPT/external help policy: language models are permitted IF you tell us where you used it (indicate in your solution) and you could still present the solution yourself; ideally: link the chatlog Why?
 - ▶ sometimes, language models get it wrong and you need to be able to check the result

- ▶ First sheet is out TODAY with deadline friday in 11 days (Nov 03, noon, i.e. 12:00)
- ▶ ChatGPT/external help policy: language models are permitted IF you tell us where you used it (indicate in your solution) and you could still present the solution yourself; ideally: link the chatlog Why?
 - ▶ sometimes, language models get it wrong and you need to be able to check the result
 - ▶ you need to take full responsibility for correctness of the result

- ▶ First sheet is out TODAY with deadline friday in 11 days (Nov 03, noon, i.e. 12:00)
- ▶ ChatGPT/external help policy: language models are permitted IF you tell us where you used it (indicate in your solution) and you could still present the solution yourself; ideally: link the chatlog Why?
 - ▶ sometimes, language models get it wrong and you need to be able to check the result
 - ▶ you need to take full responsibility for correctness of the result
 - ▶ we want to understand how students use the available tools

Example

B

Write python code to automatically identify outliers, which are defined as any students with a pre-test score higher than 3 standard deviations above the mean. Write python code that removes these outliers from the data set.



**UNIVERSITÄT
BIELEFELD**

Faculty of Technology

Example

B

Write python code to automatically identify outliers, which are defined as any students with a pre-test score higher than 3 standard deviations above the mean. Write python code that removes these outliers from the data set.



You can identify and remove outliers from a dataset using Python with the help of libraries like NumPy and pandas. Here's an example code to do this:

python Copy code

```
import numpy as np
import pandas as pd

# Generate a sample dataset (you can replace this with your own data)
data = {
    'StudentID': [1, 2, 3, 4, 5, 6, 7, 8, 9, 10],
    'PreTestScore': [85, 92, 88, 110, 60, 105, 95, 102, 80, 120]
}

# Create a DataFrame from the data
df = pd.DataFrame(data)

# Calculate the mean and standard deviation of the PreTestScore
mean_score = df['PreTestScore'].mean()
std_dev = df['PreTestScore'].std()

# Define a threshold for identifying outliers
threshold = mean_score + 3 * std_dev

# Identify outliers
outliers = df[df['PreTestScore'] > threshold]

# Remove outliers from the dataset
df = df[df['PreTestScore'] <= threshold]
```



UNIVERSITÄT
BIELEFELD

Faculty of Technology

- ▶ First tutorial sessions: Python introduction with our intelligent tutoring system (bring your own device!)

- ▶ First tutorial sessions: Python introduction with our intelligent tutoring system (bring your own device!)
- ▶ Who wants to take part today, right after the lecture?

- ▶ First tutorial sessions: Python introduction with our intelligent tutoring system (bring your own device!)
- ▶ Who wants to take part today, right after the lecture?
- ▶ Who wants to take part Thursday, 16-18

- ▶ First tutorial sessions: Python introduction with our intelligent tutoring system (bring your own device!)
- ▶ Who wants to take part today, right after the lecture?
- ▶ Who wants to take part Thursday, 16-18
- ▶ Who wants to take part next monday?

- ▶ First tutorial sessions: Python introduction with our intelligent tutoring system (bring your own device!)
- ▶ Who wants to take part today, right after the lecture?
- ▶ Who wants to take part Thursday, 16-18
- ▶ Who wants to take part next monday?
- ▶ Who wants to take part next thursday?

- ▶ We define random variables via a domain of possible values \mathcal{X} and a probability mass/density function p assigning probability mass to each possible value

- ▶ We define random variables via a domain of possible values \mathcal{X} and a probability mass/density function p assigning probability mass to each possible value
- ▶ discrete random variables for finite/countably infinite sets; continuous random variables for (compact subsets of) the real numbers

- ▶ We define random variables via a domain of possible values \mathcal{X} and a probability mass/density function p assigning probability mass to each possible value
- ▶ discrete random variables for finite/countably infinite sets; continuous random variables for (compact subsets of) the real numbers
- ▶ marginal probabilities for single variables, joint/conditional probabilities for combinations of variables

- ▶ We define random variables via a domain of possible values \mathcal{X} and a probability mass/density function p assigning probability mass to each possible value
- ▶ discrete random variables for finite/countably infinite sets; continuous random variables for (compact subsets of) the real numbers
- ▶ marginal probabilities for single variables, joint/conditional probabilities for combinations of variables
- ▶ most important density functions for continuous variables (in this lecture): Gaussian (!), t -distribution

- ▶ We define random variables via a domain of possible values \mathcal{X} and a probability mass/density function p assigning probability mass to each possible value
- ▶ discrete random variables for finite/countably infinite sets; continuous random variables for (compact subsets of) the real numbers
- ▶ marginal probabilities for single variables, joint/conditional probabilities for combinations of variables
- ▶ most important density functions for continuous variables (in this lecture): Gaussian (!), t -distribution
- ▶ means of random variables tend to be Gaussian (central limit theorem)

Preamble: Study Design

Setup

Imagine you have invented a cool pedagogical intervention



Child with VR glasses by Julia M Cameron ([Link](#)); Usage according to pexels license.

Setup

Imagine you have invented a cool pedagogical intervention




Child with VR glasses by Julia M Cameron ([Link](#)); Usage according to pexels license.

... but how do you know if it is effective?


Attempt 1: Single Class Study

class of students


Attempt 1: Single Class Study



Attempt 1: Single Class Study




Attempt 1: Single Class Study



What are the problems here?

Attempt 1: Single Class Study




What are the problems here?

- ▶ Maybe the students already knew everything before the intervention


Attempt 2: Pre- and post-test

class of students


Attempt 2: Pre- and post-test




Attempt 2: Pre- and post-test



Attempt 2: Pre- and post-test




Attempt 2: Pre- and post-test



What are the problems here?

Attempt 2: Pre- and post-test




What are the problems here?

- Maybe the learning is not due to the intervention but something else


Attempt 3: Controlled study

intervention group


Attempt 3: Controlled study




Attempt 3: Controlled study



Attempt 3: Controlled study




Attempt 3: Controlled study



What are the problems here?


Attempt 3: Controlled study




What are the problems here?

- ▶ Maybe students in the intervention condition are just better

Attempt 4: Controlled pre-/post-test study



Attempt 4: Controlled pre-/post-test study



- ▶ Regression to the mean if intervention group are the (seemingly) weaker students

- ▶ Regression to the mean if intervention group are the (seemingly) weaker students
- ▶ Are the study participants representative of the underlying population?

- ▶ Regression to the mean if intervention group are the (seemingly) weaker students
- ▶ Are the study participants representative of the underlying population?
- ▶ Are there additional confounders? \Rightarrow randomize groups and blind participants & teachers, if possible

- ▶ Regression to the mean if intervention group are the (seemingly) weaker students
- ▶ Are the study participants representative of the underlying population?
- ▶ Are there additional confounders? \Rightarrow randomize groups and blind participants & teachers, if possible
- ▶ Are the tests representative of what you want to test?

- ▶ Regression to the mean if intervention group are the (seemingly) weaker students
- ▶ Are the study participants representative of the underlying population?
- ▶ Are there additional confounders? \Rightarrow randomize groups and blind participants & teachers, if possible
- ▶ Are the tests representative of what you want to test?
- ▶ ... many more (Kulik and Fletcher 2016)

Example data

- ▶ Let's say we have the following data from a study

Example data

- ▶ Let's say we have the following data from a study

student	control condition				
	1	2	3	4	5
pre-test score	24	17	29	85	31
post-test score	67	60	-	80	75

- Let's say we have the following data from a study

control condition					
student	1	2	3	4	5
pre-test score	24	17	29	85	31
post-test score	67	60	-	80	75

intervention condition				
student	6	7	8	9
pre-test score	20	27	23	16
post-test score	71	75	73	68

- Let's say we have the following data from a study

control condition					
student	1	2	3	4	5
pre-test score	24	17	29	85	31
post-test score	67	60	-	80	75

intervention condition				
student	6	7	8	9
pre-test score	20	27	23	16
post-test score	71	75	73	68

attention! This fictional data set is too small for actual statistics! This is only for illustration purposes!

Assume we have recorded pre- and post-test scores for control and intervention group

Assume we have recorded pre- and post-test scores for control and intervention group

- ▶ Are there students with implausibly high pre-test scores? (outlier detection)

Assume we have recorded pre- and post-test scores for control and intervention group

- ▶ Are there students with implausibly high pre-test scores? (outlier detection)
- ▶ If students have missing post-test values, what can we most plausibly fill in? (imputation)

Assume we have recorded pre- and post-test scores for control and intervention group

- ▶ Are there students with implausibly high pre-test scores? (outlier detection)
- ▶ If students have missing post-test values, what can we most plausibly fill in? (imputation)
- ▶ Do students with high pre-test scores also tend to have high post-test scores? (correlation)

Assume we have recorded pre- and post-test scores for control and intervention group


- ▶ Are there students with implausibly high pre-test scores? (outlier detection)
- ▶ If students have missing post-test values, what can we most plausibly fill in? (imputation)
- ▶ Do students with high pre-test scores also tend to have high post-test scores? (correlation)
- ▶ Do students in the intervention group improve their scores more than in the control group? (statistical testing)

Assume we have recorded pre- and post-test scores for control and intervention group


- ▶ Are there students with implausibly high pre-test scores? (outlier detection)
- ▶ If students have missing post-test values, what can we most plausibly fill in? (imputation)
- ▶ Do students with high pre-test scores also tend to have high post-test scores? (correlation)
- ▶ Do students in the intervention group improve their scores more than in the control group? (statistical testing)
- ▶ Do students improve between pre- and post-test? (paired statistical testing)

Outlier detection


Example: Outlier detection



Example: Outlier detection



Example: Outlier detection



Outlier Detection

Assume example data $x_1, \dots, x_m \in \mathcal{X}$.

Outlier Detection

Assume example data $x_1, \dots, x_m \in \mathcal{X}$.

Assume a probability density p fitted to that data.

Outlier Detection

Assume example data $x_1, \dots, x_m \in \mathcal{X}$.

Assume a probability density p fitted to that data.

We call points x_i **outliers** for threshold $\epsilon > 0$ if $p(x_i) < \epsilon$.

- ▶ Fit Gaussian, i.e. compute empiric mean μ and standard deviation σ of data

- ▶ Fit Gaussian, i.e. compute empiric mean μ and standard deviation σ of data
- ▶ Compute z -score for each data point x_1, \dots, x_m :

$$z = \frac{|x_i - \mu|}{\sigma}$$

- ▶ Fit Gaussian, i.e. compute empiric mean μ and standard deviation σ of data
- ▶ Compute z -score for each data point x_1, \dots, x_m :

$$z = \frac{|x_i - \mu|}{\sigma}$$

- ▶ Remove data point if $z > k$ for some threshold k

- ▶ Fit Gaussian, i.e. compute empiric mean μ and standard deviation σ of data
- ▶ Compute z -score for each data point x_1, \dots, x_m :

$$z = \frac{|x_i - \mu|}{\sigma}$$

- ▶ Remove data point if $z > k$ for some threshold k
- ▶ $\epsilon = \frac{1}{\sqrt{2\pi} \cdot \sigma} \cdot \exp(-\frac{1}{2}k^2)$


- ▶ Approach 1: perform outlier detection separately for each dimension

- ▶ Approach 1: perform outlier detection separately for each dimension
- ▶ Approach 2: Compute covariance matrix Σ ; z -score becomes Mahalanobis distance:

$$z = \sqrt{(x_i - \mu)^T \cdot \Sigma^{-1} \cdot (x_i - \mu)}$$

- ▶ Approach 1: perform outlier detection separately for each dimension
- ▶ Approach 2: Compute covariance matrix Σ ; z -score becomes Mahalanobis distance:


$$z = \sqrt{(x_i - \mu)^T \cdot \Sigma^{-1} \cdot (x_i - \mu)}$$



- ▶ Approach 1: perform outlier detection separately for each dimension
- ▶ Approach 2: Compute covariance matrix Σ ; z -score becomes Mahalanobis distance:

$$z = \sqrt{(x_i - \mu)^T \cdot \Sigma^{-1} \cdot (x_i - \mu)}$$

Gaussian density; iso-line for 1σ




- ▶ Approach 1: perform outlier detection separately for each dimension
- ▶ Approach 2: Compute covariance matrix Σ ; z -score becomes Mahalanobis distance:

$$z = \sqrt{(x_i - \mu)^T \cdot \Sigma^{-1} \cdot (x_i - \mu)}$$

Gaussian density; iso-line for 1σ


2σ



- ▶ Approach 1: perform outlier detection separately for each dimension
- ▶ Approach 2: Compute covariance matrix Σ ; z -score becomes Mahalanobis distance:

$$z = \sqrt{(x_i - \mu)^T \cdot \Sigma^{-1} \cdot (x_i - \mu)}$$

Gaussian density; iso-line for 1σ



- ▶ p needs to be an accurate model of the data – how to ensure that?

► p needs to be an accurate model of the data – how to ensure that?

⇒ e.g. generative models, density estimation

- ▶ p needs to be an accurate model of the data – how to ensure that?
- ⇒ e.g. generative models, density estimation
- ▶ We fit p to the data **including** outliers; chicken-and-egg problem

- ▶ p needs to be an accurate model of the data – how to ensure that?

- ⇒ e.g. generative models, density estimation


- ▶ We fit p to the data **including** outliers; chicken-and-egg problem

- ⇒ iterative models (remove an outlier, fit again, etc.)


- ▶ p needs to be an accurate model of the data – how to ensure that?
 - ⇒ e.g. generative models, density estimation
- ▶ We fit p to the data **including** outliers; chicken-and-egg problem
 - ⇒ iterative models (remove an outlier, fit again, etc.)
 - ⇒ density estimation & outlier exclusion in one go, e.g. one-class SVM

Imputation


Example: Imputation



Example: Imputation



Example: Imputation



Outlier Detection

Assume example data $x_1, \dots, x_m \in (\mathbb{R} \cup \{\text{NaN}\})^n$.

Outlier Detection

Assume example data $x_1, \dots, x_m \in (\mathbb{R} \cup \{\text{NaN}\})^n$.
We call a value $x_{i,j} = \text{NaN}$ a **missing value**.

Outlier Detection

Assume example data $x_1, \dots, x_m \in (\mathbb{R} \cup \{\text{NaN}\})^n$.

We call a value $x_{i,j} = \text{NaN}$ a **missing value**.

Assume a probability density p fitted to the data.

Outlier Detection

Assume example data $x_1, \dots, x_m \in (\mathbb{R} \cup \{\text{NaN}\})^n$.

We call a value $x_{i,j} = \text{NaN}$ a **missing value**.

Assume a probability density p fitted to the data.

Imputation means to replace a missing $x_{i,j}$ with one that maximizes p .


- ▶ Fit Gaussian, i.e. compute empiric mean μ and standard deviation σ of data, excluding missing values

- ▶ Fit Gaussian, i.e. compute empiric mean μ and standard deviation σ of data, excluding missing values
- ▶ Replace NaNs with $\mu \Rightarrow$ maximizes $p(x_{i,j})$


- ▶ Fit Gaussian, i.e. compute empiric mean μ and standard deviation σ of data, excluding missing values
- ▶ Replace NaNs with $\mu \Rightarrow$ maximizes $p(x_{i,j})$
- ▶ Approach 1 for multi-dimensional data: treat each dimension separately

- ▶ Fit Gaussian, i.e. compute empiric mean μ and standard deviation σ of data, excluding missing values
- ▶ Replace NaNs with $\mu \Rightarrow$ maximizes $p(x_{i,j})$
- ▶ Approach 1 for multi-dimensional data: treat each dimension separately
- ▶ Approach 2: maximize conditional $p(x_{i,j} | x_{i,1}, \dots, x_{i,j-1}, x_{i,j+1}, \dots, x_{i,n})$

- ▶ Fit Gaussian, i.e. compute empiric mean μ and standard deviation σ of data, excluding missing values
- ▶ Replace NaNs with $\mu \Rightarrow$ maximizes $p(x_{i,j})$
- ▶ Approach 1 for multi-dimensional data: treat each dimension separately
- ▶ Approach 2: maximize conditional $p(x_{i,j} | x_{i,1}, \dots, x_{i,j-1}, x_{i,j+1}, \dots, x_{i,n})$



- ▶ Fit Gaussian, i.e. compute empiric mean μ and standard deviation σ of data, excluding missing values
- ▶ Replace NaNs with $\mu \Rightarrow$ maximizes $p(x_{i,j})$
- ▶ Approach 1 for multi-dimensional data: treat each dimension separately
- ▶ Approach 2: maximize conditional $p(x_{i,j} | x_{i,1}, \dots, x_{i,j-1}, x_{i,j+1}, \dots, x_{i,n})$



- ▶ Constant value imputation artificially decreases variance of data

- ▶ Constant value imputation artificially decreases variance of data
- ⇒ Sample from p instead of always taking argmax

- ▶ Constant value imputation artificially decreases variance of data
- ⇒ Sample from p instead of always taking argmax
- ▶ better models of conditionals?

- ▶ Constant value imputation artificially decreases variance of data
- ⇒ Sample from p instead of always taking argmax
- ▶ better models of conditionals?
- ⇒ Regression approaches

Correlations

Correlation: Example


- ▶ Intuition: How strongly are two random variables associated?

Correlation: Example

- ▶ Intuition: How strongly are two random variables associated?
- ▶ Attempt 1: Slope of linear regression line


Correlation: Example

- ▶ Intuition: How strongly are two random variables associated?
- ▶ Attempt 1: Slope of linear regression line




Correlation: Example

- ▶ Intuition: How strongly are two random variables associated?
- ▶ Attempt 1: Slope of linear regression line



Correlation: Example


- ▶ Intuition: How strongly are two random variables associated?
- ▶ Attempt 1: Slope of linear regression line



- ▶ Problem: What about the **scaling** of the data?

Correlation: Example

- ▶ Intuition: How strongly are two random variables associated?
- ▶ Attempt 1: Slope of linear regression line




- ▶ Problem: What about the **scaling** of the data?

⇒ normalize data beforehand: $\tilde{x} = (x - \mu_x)/\sigma_x$ and $\tilde{y} = (y - \mu_y)/\sigma_y$.

Sir Francis Galton *FRS FRAI* ([/ˈɡɔːltən/](#); 16 February 1822 – 17 January 1911) was a British [polymath](#) and the originator of the [eugenics movement](#) during the [Victorian era](#).^{[1][2]}

Galton produced over 340 papers and books. He also developed the statistical concept of [correlation](#) and widely promoted [regression toward the mean](#).

In recent years, he has received significant criticism for being a proponent of [social Darwinism](#), [eugenics](#), and [scientific racism](#); he was a pioneer of [eugenics](#), coining the term itself in 1883.



Linear/Pearson correlation

Karl Pearson [FRS FRSE](#)^[1] ([/'piərsən/](#); born **Carl Pearson**; 27 March 1857 – 27 April 1936^[2]) was an English [mathematician](#) and [biostatistician](#). He has been credited with establishing the discipline of [mathematical statistics](#).^{[3][4]} He founded the world's first university statistics department at [University College London](#) in 1911, and contributed significantly to the field of [biometrics](#) and [meteorology](#). Pearson was also a proponent of [social Darwinism](#) and [eugenics](#), and his thought is an example of what is today described as [scientific racism](#).



Linear correlation coefficient

Let $(x_1, y_1), \dots, (x_m, y_m) \in \mathbb{R}^2$.

Linear correlation coefficient

Let $(x_1, y_1), \dots, (x_m, y_m) \in \mathbb{R}^2$.

the linear correlation coefficient is defined as:

$$r = \frac{1}{m} \sum_{i=1}^m \frac{x_i - \mu_x}{\sigma_x} \cdot \frac{y_i - \mu_y}{\sigma_y},$$

where μ_x and μ_y are the means and σ_x and σ_y are the standard deviations of the data.

Linear correlation coefficient

Let $(x_1, y_1), \dots, (x_m, y_m) \in \mathbb{R}^2$.

the linear correlation coefficient is defined as:

$$r = \frac{1}{m} \sum_{i=1}^m \frac{x_i - \mu_x}{\sigma_x} \cdot \frac{y_i - \mu_y}{\sigma_y},$$

where μ_x and μ_y are the means and σ_x and σ_y are the standard deviations of the data.

- range: $[-1, +1]$; symmetric for x and y

Linear correlation coefficient

Let $(x_1, y_1), \dots, (x_m, y_m) \in \mathbb{R}^2$.

the linear correlation coefficient is defined as:

$$r = \frac{1}{m} \sum_{i=1}^m \frac{x_i - \mu_x}{\sigma_x} \cdot \frac{y_i - \mu_y}{\sigma_y},$$

where μ_x and μ_y are the means and σ_x and σ_y are the standard deviations of the data.

- ▶ range: $[-1, +1]$; symmetric for x and y
- ▶ interpretation: slope of the linear regression line **after** normalization of x and y

Linear correlation coefficient

Let $(x_1, y_1), \dots, (x_m, y_m) \in \mathbb{R}^2$.


the linear correlation coefficient is defined as:

$$r = \frac{1}{m} \sum_{i=1}^m \frac{x_i - \mu_x}{\sigma_x} \cdot \frac{y_i - \mu_y}{\sigma_y},$$

where μ_x and μ_y are the means and σ_x and σ_y are the standard deviations of the data.

- ▶ range: $[-1, +1]$; symmetric for x and y
- ▶ interpretation: slope of the linear regression line **after** normalization of x and y
- ▶ Rules of thumb: $r \leq 0.3$ is very small, $r \in (0.3, 0.5]$ is small, $r \in (0.5, 0.7]$ is moderate, $r \in (0.7, 0.9]$ is high, $r \in (0.9, 1.0]$ is very high (Mukaka 2012)

Problems: Nonlinear correlations



Charles Edward Spearman, [FRS^{\[1\]\[3\]}](#) (10 September 1863 – 17 September 1945) was an English psychologist known for work in [statistics](#), as a pioneer of [factor analysis](#), and for [Spearman's rank correlation coefficient](#).

The Eugenics Review


[Eugen Rev.](#) 1914 Oct; 6(3): 219–237.

PMCID: PMC2987066

PMID: [21259592](#)

The heredity of abilities

[C. Spearman](#)



Rank correlation coefficient

Let $(x_1, y_1), \dots, (x_m, y_m) \in \mathbb{R}^2$.

Rank correlation coefficient

Let $(x_1, y_1), \dots, (x_m, y_m) \in \mathbb{R}^2$.

Replace x_i and y_i by their **rank**.

Rank correlation coefficient

Let $(x_1, y_1), \dots, (x_m, y_m) \in \mathbb{R}^2$.

Replace x_i and y_i by their **rank**.

The rank correlation coefficient is then defined as r on the ranks.

Rank correlation coefficient

Let $(x_1, y_1), \dots, (x_m, y_m) \in \mathbb{R}^2$.

Replace x_i and y_i by their **rank**.

The rank correlation coefficient is then defined as r on the ranks.

- ▶ similar interpretation to linear correlation coefficient r – but in rank space

Rank correlation coefficient

Let $(x_1, y_1), \dots, (x_m, y_m) \in \mathbb{R}^2$.

Replace x_i and y_i by their **rank**.

The rank correlation coefficient is then defined as r on the ranks.

- ▶ similar interpretation to linear correlation coefficient r – but in rank space
- ▶ no direct relation to regression line on the original data; no “drawing” of fit \Rightarrow “non-parametric statistic”

Rank correlation coefficient


Let $(x_1, y_1), \dots, (x_m, y_m) \in \mathbb{R}^2$.

Replace x_i and y_i by their **rank**.


The rank correlation coefficient is then defined as r on the ranks.


- ▶ similar interpretation to linear correlation coefficient r – but in rank space
- ▶ no direct relation to regression line on the original data; no “drawing” of fit \Rightarrow “non-parametric statistic”
- ▶ alternatives: Cohen’s κ , Krippendorff’s α , ...

Non-monotonic relations




Non-monotonic relations





- correlation measures capture only **monotonic** relationships




► correlation measures capture only **monotonic** relationships


⇒ independence \Rightarrow no correlation, but not vice versa

Statistical Tests


Example: Bar plot




Example: Bar plot




Example: Bar plot




Example: Bar plot




Example: Bar plot




Example (continued)




Example (continued)




Example (continued)




Example (continued)




Example (continued)



Example (continued)



Example (continued)



- What is the probability of $\frac{\mu_y - \mu_x}{\sigma}$ being that large just by random chance?

- ▶ We want to be certain that an observed effect could not also have been generated by mere chance

- ▶ We want to be certain that an observed effect could not also have been generated by mere chance – more precisely: by a random baseline model (the **null hypothesis**)

- ▶ We want to be certain that an observed effect could not also have been generated by mere chance – more precisely: by a random baseline model (the **null hypothesis**)
- ▶ If we are certain that the null hypothesis could not have generated the observed effect, we **reject the null hypothesis**

- ▶ We want to be certain that an observed effect could not also have been generated by mere chance – more precisely: by a random baseline model (the **null hypothesis**)
- ▶ If we are certain that the null hypothesis could not have generated the observed effect, we **reject the null hypothesis**
- ▶ Type I error: Rejecting the null hypothesis even though it did generate the observed effect (p value)

- ▶ We want to be certain that an observed effect could not also have been generated by mere chance – more precisely: by a random baseline model (the **null hypothesis**)
- ▶ If we are certain that the null hypothesis could not have generated the observed effect, we **reject the null hypothesis**
- ▶ Type I error: Rejecting the null hypothesis even though it did generate the observed effect (p value)
- ▶ Type II error: Failing to reject the null hypothesis even though it did not generate the observed effect (statistical power)

Null hypothesis testing

To quantify an effect, we compute a statistic \hat{s} .

Null hypothesis testing

To quantify an effect, we compute a statistic \hat{s} .

The null hypothesis is that \hat{s} is generated just due to random chance.

Null hypothesis testing

To quantify an effect, we compute a statistic \hat{s} .

The null hypothesis is that \hat{s} is generated just due to random chance.

We model the probability $p = P(s \geq \hat{s})$ under the null hypothesis.

Null hypothesis testing

To quantify an effect, we compute a statistic \hat{s} .

The null hypothesis is that \hat{s} is generated just due to random chance.

We model the probability $p = P(s \geq \hat{s})$ under the null hypothesis.

We reject the null hypothesis if p is sufficiently small (e.g. $p < 0.05$). We say the effect is **significant** (at 0.05-level).

Null hypothesis testing

To quantify an effect, we compute a statistic \hat{s} .

The null hypothesis is that \hat{s} is generated just due to random chance.

We model the probability $p = P(s \geq \hat{s})$ under the null hypothesis.

We reject the null hypothesis if p is sufficiently small (e.g. $p < 0.05$). We say the effect is **significant** (at 0.05-level).

- Key advantage: We only need a model of the statistic under the null hypothesis

Null hypothesis testing

To quantify an effect, we compute a statistic \hat{s} .

The null hypothesis is that \hat{s} is generated just due to random chance.

We model the probability $p = P(s \geq \hat{s})$ under the null hypothesis.

We reject the null hypothesis if p is sufficiently small (e.g. $p < 0.05$). We say the effect is **significant** (at 0.05-level).

- ▶ Key advantage: We only need a model of the statistic under the null hypothesis
- ▶ Tests differ in statistic, assumptions, and model

t -Test (Welch version)

William Sealy Gosset (13 June 1876 – 16 October 1937) was an English statistician, chemist and brewer who served as **Head Brewer** of **Guinness** and Head Experimental Brewer of Guinness and was a pioneer of modern statistics. He pioneered small sample experimental design and analysis with an economic approach to the logic of uncertainty. Gosset published under the **pen name Student** and developed most famously **Student's t-distribution** – originally called Student's "z" – and "Student's test of **statistical significance**".^[1]



t -Test (Welch version)

(Student 1908; Welch 1947)

- ▶ Assume two samples with n_x and n_y data points, means μ_x and μ_y , and standard deviations σ_x and σ_y

- ▶ Assume two samples with n_x and n_y data points, means μ_x and μ_y , and standard deviations σ_x and σ_y
- ▶ Test statistic: mean difference scaled by standard error: $t = \frac{\mu_x - \mu_y}{\tilde{\sigma}}$

- ▶ Assume two samples with n_x and n_y data points, means μ_x and μ_y , and standard deviations σ_x and σ_y
- ▶ Test statistic: mean difference scaled by standard error: $t = \frac{\mu_x - \mu_y}{\tilde{\sigma}}$
- ▶ Problem: Which $\tilde{\sigma}$ do we take?

$$\tilde{\sigma} = \sqrt{\frac{\sigma_x^2}{n_x} + \frac{\sigma_y^2}{n_y}}$$

- ▶ Assume two samples with n_x and n_y data points, means μ_x and μ_y , and standard deviations σ_x and σ_y
- ▶ Test statistic: mean difference scaled by standard error: $t = \frac{\mu_x - \mu_y}{\tilde{\sigma}}$
- ▶ Problem: Which $\tilde{\sigma}$ do we take? \Rightarrow compromise between standard error of sample one and sample two:

$$\tilde{\sigma} = \sqrt{\frac{\sigma_x^2}{n_x} + \frac{\sigma_y^2}{n_y}}$$

- ▶ Assume two samples with n_x and n_y data points, means μ_x and μ_y , and standard deviations σ_x and σ_y
- ▶ Test statistic: mean difference scaled by standard error: $t = \frac{\mu_x - \mu_y}{\tilde{\sigma}}$
- ▶ Problem: Which $\tilde{\sigma}$ do we take? \Rightarrow compromise between standard error of sample one and sample two:


$$\tilde{\sigma} = \sqrt{\frac{\sigma_x^2}{n_x} + \frac{\sigma_y^2}{n_y}}$$

- ▶ Under the null hypothesis, μ_x and μ_y stem from the same Gaussian with $\tilde{\sigma}$ and t is t -distributed with parameter


$$\nu = \frac{\tilde{\sigma}^4}{\frac{\sigma_x^4}{n_x^2 \cdot (n_x - 1)} + \frac{\sigma_y^4}{n_y^2 \cdot (n_y - 1)}}$$

- For our toy data set: $\tilde{\sigma} \approx 1$, $t = 7.39$, $\nu = 4.93$, $p = 0.0007$

- For our toy data set: $\tilde{\sigma} \approx 1$, $t = 7.39$, $\nu = 4.93$, $p = 0.0007$



- For our toy data set: $\tilde{\sigma} \approx 1$, $t = 7.39$, $\nu = 4.93$, $p = 0.0007$



Assumptions of the t -test

- ▶ Samples are independent

Assumptions of the t -test

- ▶ Samples are independent
- ▶ Sample means are Gaussian distributed (Central limit theorem)

- ▶ Samples are independent
 - ▶ Sample means are Gaussian distributed (Central limit theorem)
- ⇒ one way of verification: check normality of original data distribution via Shapiro-Wilk-test (or other tests) ⇒ mean of Gaussian variables is Gaussian

- ▶ Problem: significance testing only tells us whether an effect could be generated by the null hypothesis – not how strong it is

- ▶ Problem: significance testing only tells us whether an effect could be generated by the null hypothesis – not how strong it is
- ▶ Effect size: Cohen's d , defined as $d = \frac{|\mu_1 - \mu_2|}{\sigma}$

- ▶ Problem: significance testing only tells us whether an effect could be generated by the null hypothesis – not how strong it is
- ▶ Effect size: Cohen's d , defined as $d = \frac{|\mu_1 - \mu_2|}{\sigma}$
- ▶ Again: σ needs to be a compromise between σ_1 and σ_2 , e.g.

$$\sigma = \sqrt{\frac{(n_1 - 1) \cdot \sigma_1^2 + (n_2 - 1) \cdot \sigma_2^2}{n_1 + n_2 - 2}}$$

- ▶ Problem: significance testing only tells us whether an effect could be generated by the null hypothesis – not how strong it is
- ▶ Effect size: Cohen's d , defined as $d = \frac{|\mu_1 - \mu_2|}{\sigma}$
- ▶ Again: σ needs to be a compromise between σ_1 and σ_2 , e.g.

$$\sigma = \sqrt{\frac{(n_1 - 1) \cdot \sigma_1^2 + (n_2 - 1) \cdot \sigma_2^2}{n_1 + n_2 - 2}}$$

- ▶ Rules of thumb: $d < 0.2$ is very small, $d \in [0.2, 0.5)$ is small, $d \in [0.5, 0.8)$ is moderate, $d \in [0.8, 1.2)$ is large, $d \in [1.2, 2.0)$ is very large, $d \geq 2.0$ is huge (Sawilowsky 2009)

- ▶ Problem: significance testing only tells us whether an effect could be generated by the null hypothesis – not how strong it is
- ▶ Effect size: Cohen's d , defined as $d = \frac{|\mu_1 - \mu_2|}{\sigma}$
- ▶ Again: σ needs to be a compromise between σ_1 and σ_2 , e.g.

$$\sigma = \sqrt{\frac{(n_1 - 1) \cdot \sigma_1^2 + (n_2 - 1) \cdot \sigma_2^2}{n_1 + n_2 - 2}}$$

- ▶ Rules of thumb: $d < 0.2$ is very small, $d \in [0.2, 0.5)$ is small, $d \in [0.5, 0.8)$ is moderate, $d \in [0.8, 1.2)$ is large, $d \in [1.2, 2.0)$ is very large, $d \geq 2.0$ is huge (Sawilowsky 2009)
- ▶ Side note: largest measured effect sizes in psychology for educational studies

Example: signed rank test

- ▶ Question: Are post-test scores significantly higher than pre-test scores?

Example: signed rank test

► Question: Are post-test scores significantly higher than pre-test scores?

⇒ **paired** data

Example: signed rank test

► Question: Are post-test scores significantly higher than pre-test scores?

⇒ **paired** data

student	1	2	3	4	5	6	7	8	9
pre-test score	24	17	29	85	31	20	27	23	16
post-test score	67	60	70	80	75	71	75	73	68

Example: signed rank test

► Question: Are post-test scores significantly higher than pre-test scores?

⇒ **paired** data

student	1	2	3	4	5	6	7	8	9
pre-test score	24	17	29	85	31	20	27	23	16
post-test score	67	60	70	80	75	71	75	73	68
improvement	43	43	41	-5	44	51	48	50	52

Example: signed rank test

- Question: Are post-test scores significantly higher than pre-test scores?

⇒ **paired** data

student	1	2	3	4	5	6	7	8	9
pre-test score	24	17	29	85	31	20	27	23	16
post-test score	67	60	70	80	75	71	75	73	68
improvement	43	43	41	-5	44	51	48	50	52

- Idea 1: If there are more positive than negative numbers, probably yes

Example: signed rank test

- ▶ Question: Are post-test scores significantly higher than pre-test scores?

⇒ **paired** data

student	1	2	3	4	5	6	7	8	9
pre-test score	24	17	29	85	31	20	27	23	16
post-test score	67	60	70	80	75	71	75	73	68
improvement	43	43	41	-5	44	51	48	50	52

- ▶ Idea 1: If there are more positive than negative numbers, probably yes
- ▶ Idea 2: Put higher weights on bigger numbers

Wilcoxon signed rank test (Wilcoxon 1945)

- ▶ Sort improvements acc. to absolute value

Wilcoxon signed rank test (Wilcoxon 1945)

- ▶ Sort improvements acc. to absolute value
- ▶ test statistic T : sum of ranks, with signs

- ▶ Sort improvements acc. to absolute value
- ▶ test statistic T : sum of ranks, with signs

improvement 43 43 41 -5 44 51 48 50 52

- ▶ Sort improvements acc. to absolute value
- ▶ test statistic T : sum of ranks, with signs

improvement	43	43	41	-5	44	51	48	50	52
signed rank	3.5	3.5	2	-1	5	8	6	7	9

- ▶ Sort improvements acc. to absolute value
- ▶ test statistic T : sum of ranks, with signs

improvement	43	43	41	-5	44	51	48	50	52
signed rank	3.5	3.5	2	-1	5	8	6	7	9
$\Rightarrow T = 43$									

- ▶ Sort improvements acc. to absolute value
- ▶ test statistic T : sum of ranks, with signs

improvement	43	43	41	-5	44	51	48	50	52
signed rank	3.5	3.5	2	-1	5	8	6	7	9
$\Rightarrow T = 43$									

- ▶ Under null hypothesis, p_T can be computed exactly (for small n) – or via Gaussian approximation (for large n)

- Under the null hypothesis, any combination of signs amongst the ranks is equally likely

rank	sign combination							
1	-	+	-	-	+	+	-	+
2	-	-	+	-	+	-	+	+
3	-	-	-	+	-	+	+	+
T	-6	-4	-2	0	0	+2	+4	+6

- Under the null hypothesis, any combination of signs amongst the ranks is equally likely

rank	sign combination							
1	-	+	-	-	+	+	-	+
2	-	-	+	-	+	-	+	+
3	-	-	-	+	-	+	+	+
T	-6	-4	-2	0	0	+2	+4	+6

- $p(-6) = p(-4) = p(-2) = p(+2) = p(+4) = p(+6) = \frac{1}{8}, p(0) = \frac{1}{4}$

Assumptions of Wilcoxon signed rank test

- ▶ Samples are independent

Assumptions of Wilcoxon signed rank test

- ▶ Samples are independent
- ▶ Differences are symmetrically distributed

- ▶ This visualization summarizes all key concepts from null hypothesis testing neatly:

<https://rpsychologist.com/d3/nhst/>

- Cohen, Jacob (1988). **Statistical Power Analysis for the Behavioral Sciences**. New York, NY, USA: Routledge. DOI: 10.4324/9780203771587.
- Galton, Francis (1889). “I. Co-relations and their measurement, chiefly from anthropometric data”. In: **Proceedings of the Royal Society of London** 45.273-279, pp. 135–145. DOI: 10.1098/rsp1.1888.0082.
- Kulik, James A and JD Fletcher (2016). “Effectiveness of intelligent tutoring systems: a meta-analytic review”. In: **Review of educational research** 86.1, pp. 42–78. DOI: 10.3102/0034654315581420. URL: https://www.researchgate.net/profile/J-D-Fletcher/publication/277636218_Effectiveness_of_Intelligent_Tutoring_Systems_A_Meta-Analytic_Review/links/5707a6fe08ae8883a1f7e55f/Effectiveness-of-Intelligent-Tutoring-Systems-A-Meta-Analytic-Review.pdf.

- Mukaka, Mavuto M (2012). "A guide to appropriate use of correlation coefficient in medical research". In: **Malawi medical journal** 24.3, pp. 69–71. URL: <https://www.ajol.info/index.php/mmj/article/view/81576>.
- Sawilowsky, Shlomo S. (2009). "New effect size rules of thumb". In: **Journal of modern applied statistical methods** 8.2, p. 26. DOI: 10.22237/jmasm/1257035100.
- Spearman, Charles (1904). "The Proof and Measurement of Association between Two Things". In: **The American Journal of Psychology** 15.1, pp. 72–101. DOI: 10.2307/1412159.
- Student (1908). "The Probable Error of a Mean". In: **Biometrika** 6.1, pp. 1–25. DOI: 10.2307/2331554.
- Welch, Bernard Lewis (1947). "The Genralization of Student's Problem when several different population variances are involved". In: **Biometrika** 34.1-2, pp. 28–35. DOI: 10.1093/biomet/34.1-2.28.

Wilcoxon, Frank (1945). "Individual Comparisons by Ranking Methods". In: **Biometrics Bulletin** 1.6, pp. 80–83. DOI: 10.2307/3001968. URL: <http://www.jstor.org/stable/3001968>.