
1

Assignment

Dhruv Parashar-EE22BTECH11019

Question:- The transition diagram of a discrete memoryless channel with three input symbols and three output symbols is shown in the figure. The transition probabilities are as marked. The parameter α lies in the interval [0.25, 1]. The value of α for which the capacity of this channel is maximized, is **Solution:**



Variable	Description	Value
X	Input variable	$X = \{0, 1, 2\}$
Y	Output variable	$Y = \{0, 1, 2\}$
С	Channel Capacity	C
I	Mutual Information	I
Н	Entropy	Н

$$C = \max_{p(X,Y)} I(X,Y) \tag{1}$$

$$I(X,Y) = \sum_{x,y} p(x,y) \log_2 \frac{p(x,y)}{p(x)p(y)}$$
(2)

$$= \sum_{x,y} p(x,y) \log_2 \frac{p(y|x)}{p(y)}$$
(3)

$$= -\sum_{x,y} p(x,y) \log_2 p(y) + \sum_{x,y} p(x,y) \log_2 p(y|x)$$
(4)

$$= -\sum_{y} p(y) \log_2 p(y) - \left(-\sum_{x,y} p(x,y) \log_2 p(y|x)\right)$$
(5)

$$=H(Y)-H(Y|X) \tag{6}$$

Now.

$$p_X(0) = p_X(1) = p_X(2) = \frac{1}{3}$$
 (7)

For X = 0

$$p_Y(0) = \sum_{y=0}^{2} p_{Y|X}(y|x) p_X(x)$$
 (8)

$$= \frac{1}{3} \sum_{y=0}^{2} p_{Y|X}(y|x)$$
 (9)

$$=\frac{1}{3}\tag{10}$$

Similarly for X = 1, 2

$$p_Y(0) = p_Y(1) = p_Y(2) = \frac{1}{3}$$
 (11)

$$H(Y) = -\sum_{y=0}^{2} p_Y(y) \log_2 p_Y(y)$$
 (12)

$$= -\frac{1}{3}\log_2\frac{1}{3} \times 3\tag{13}$$

$$= \log_2 3 \tag{14}$$

$$H(Y|X) = -\sum_{x=0}^{2} \sum_{y=0}^{2} p_X(x) p_{Y|X}(y|x) \log_2(p_{Y|X}(y|x))$$
(15)

$$= -\sum_{x=0}^{2} p_X(x) \sum_{y=0}^{2} p_Y(y|X) \log_2(p_Y(y|X))$$
 (16)

$$= -3 \times \frac{1}{3} \left(\alpha \log_2 \alpha + (1 - \alpha) \log_2 (1 - \alpha) \right) \tag{17}$$

Using (14) and (17) in (6)

$$I(X, Y) = \log_2 3 + \alpha \log_2 \alpha + (1 - \alpha) \log_2 (1 - \alpha)$$
(18)

$$\implies \frac{d}{d\alpha}I(X,Y) = (\log_2\alpha - \log_2(1-\alpha)) \tag{19}$$

For maxima or minima $\frac{d}{d\alpha}I(X,Y) = 0$

At $\alpha = \frac{1}{2}$, I(X, Y) is minimum As I(X, Y) is maximum at $\alpha = 0, 1$ But $\alpha \in [0.25, 1]$,

$$\alpha = 1 \tag{20}$$

