

## Assignment 6

Indian Institute of Science Education and Research

CHM202: Energetics and dynamics of chemical reactions

Instructor: Dr. Arijit K. De

**Ques. 1** Suppose that (a) the attractive interactions between gas particles can be neglected, (b) the attractive interaction is dominant in a van der Waals gas, and the pressure is low enough to make the approximation  $4aP \ll (RT)^2$ . Find expressions for the fugacity coefficient of a van der Waals gas in terms of the pressure for both the cases.

**Ques. 2** Given that  $p^*(\text{H}_2\text{O}) = 0.02308 \text{ atm}$  and  $p(\text{H}_2\text{O}) = 0.02239 \text{ atm}$  in a solution in which 0.122 kg of a non-volatile solute ( $M = 241 \text{ g mol}^{-1}$ ) is dissolved in 0.920 kg water at 293 K, Calculate the activity and activity coefficient of water in the solution.

**Ques. 3** The equilibrium constant of a reaction is found to fit the expression:

$$\ln K = A + \frac{B}{T} + \frac{C}{T^3}$$

between 400 K and 500 K with  $A = -2.04$ ,  $B = -1176 \text{ K}$ , and  $C = 2.1 \times 10^7 \text{ K}^3$ . Calculate the standard reaction enthalpy and standard reaction entropy at 450 K.

**Ques. 4** By measuring the equilibrium between liquid and vapour phases of a solution at 30°C at 1.00 atm, it was found that  $x_A = 0.220$  when  $y_A = 0.314$ . Calculate the activities and activity coefficients of both components in this solution on the Raoult's law basis. The vapour pressures of the pure components at this temperature are:  $p_A^* = 73.0 \text{ kPa}$  and  $p_B^* = 92.1 \text{ kPa}$ . ( $x_A$  is the mole fraction in the liquid and  $y_A$  the mole fraction in the vapour)

**Ques. 5** The standard enthalpy of a certain reaction is approximately constant at  $+125 \text{ kJ mol}^{-1}$  from 800 K up to 1500 K. The standard reaction Gibbs energy is  $+22 \text{ kJ mol}^{-1}$  at 1120 K. Estimate the temperature at which the equilibrium constant becomes greater than 1.