



The Central Dogma of Molecular Biology

The Central Dogma of Molecular Biology



Importance of Proteins

- There are three main kinds:
 - structural - make up most body parts
 - hormone - chemical that controls the body
 - enzyme - catalyst - speeds up chemical reactions



Insulin,
a protein


Importance of Proteins

- Without proteins there would be no life
- All cells make proteins
- Proteins in your body make up your:
 - Hair
 - Nails
 - Muscles
 - Skin
 - Cartilage



Discovery of DNA (deoxyribonucleic acid)

- 1953 Watson, Crick and Wilkins determined the structure of DNA to be a **double helix**
- They won a Nobel Prize for their work



DNA and RNA Compared

■ DNA

- Found only in nucleus*
- Double helix
- Bases = ATGC
- Sugar = Deoxyribose

* DNA is also found in a select number of other organelles


■ RNA

- Found in ribosomes, nucleolus
- Single helix
- Bases = AUGC (URACIL)
- Sugar = Ribose

Introns and Exons

- **Introns** – sequences in the DNA that are NOT used to make mRNA or to make a protein
- **Exons** – sequences in the DNA that are expressed or used to make mRNA and ultimately are used to make a protein

Introns and Exons




Ribonucleic Acid (RNA)

- Three types of RNA:
 - **rRNA** = **ribosomal RNA** - makes up the ribosome
 - **mRNA** = **messenger RNA** - is the message from DNA for the construction of the new protein molecule
 - **tRNA** = **transfer RNA** - carries amino acids to ribosomes


Ribonucleic Acid (RNA)

- Ribosomal RNA is used to make a ribosome
- The ribosome “**reads**” the mRNA plan for the new protein



Ribonucleic Acid (RNA)

- Messenger RNA





Ribonucleic Acid (RNA)

■ Transfer RNA

- Each tRNA holds one amino acid
- Every tRNA has a special region called the anti-codon (3 bases)
- A tRNA anti-codon "mates" with codon on the mRNA molecule
- There are 61 different tRNA molecules, yet only about 20 amino acids

Ribonucleic Acid (RNA)

■ Transfer RNA




Transcription

- **Transcription** is the special copying of one side of the DNA molecule (the sense strand) that results in a single strand of RNA
- The original DNA is not changed
- This process can be repeated
- The amount of DNA that is transcribed is usually one gene

Transcription

- Process of Transcription
 - DNA is unzipped by an enzyme
 - Only one side fills with RNA nucleotides by the action of another enzyme **RNA polymerase**
A-U, G-C (**NO THYMINE = T**)
 - As the RNA strand separates the DNA strands reattach as before the process started
 - The result is the original DNA **plus a new RNA strand**


Transcription




Transcription

- Uracil – a base only found in RNA


Uracil (U)




Transcription



Transcription




Overview: the roles of transcription and translation in the flow of genetic information (layer 1)



Prokaryotic cell. In a cell lacking a nucleus, mRNA produced by transcription is immediately translated without additional processing.

Eukaryotic cell. The nucleus provides a separate compartment for transcription. The original RNA transcript, called pre-mRNA, is processed in various ways before leaving the nucleus as mRNA.


Overview: the roles of transcription and translation in the flow of genetic information (layer 2)



Prokaryotic cell. In a cell lacking a nucleus, mRNA produced by transcription is immediately translated without additional processing.

Eukaryotic cell. The nucleus provides a separate compartment for transcription. The original RNA transcript, called pre-mRNA, is processed in various ways before leaving the nucleus as mRNA.


Overview: the roles of transcription and translation in the flow of genetic information (layer 3)



Prokaryotic cell. In a cell lacking a nucleus, mRNA produced by transcription is immediately translated without additional processing.

Eukaryotic cell. The nucleus provides a separate compartment for transcription. The original RNA transcript, called pre-mRNA, is processed in various ways before leaving the nucleus as mRNA.


Overview: the roles of transcription and translation in the flow of genetic information (layer 4)



Prokaryotic cell. In a cell lacking a nucleus, mRNA produced by transcription is immediately translated without additional processing.

Eukaryotic cell. The nucleus provides a separate compartment for transcription. The original RNA transcript, called pre-mRNA, is processed in various ways before leaving the nucleus as mRNA.

Overview: the roles of transcription and translation in the flow of genetic information (layer 5)




Prokaryotic cell. In a cell lacking a nucleus, mRNA produced by transcription is immediately translated without additional processing.


Eukaryotic cell. The nucleus provides a separate compartment for transcription. The original RNA transcript, called pre-mRNA, is processed in various ways before leaving the nucleus as mRNA.

Translation

- **Translation** is the reading of the RNA code to make proteins or polypeptides
- Translation is often called **protein synthesis**




The triplet code



Translation

- **Triplet codons** - groups of three bases on mRNA that code for specific amino acids




The dictionary of the genetic code

		Second mRNA base					
		U	C	A	G		
First mRNA base (5' end)	U	UUU	UCU	UAU	UGU	U	Third mRNA base (3' end)
		UUC	UCC	UAC	UGC	C	
		UUA	UCA	UAA Stop	UGA Stop	A	
		UUG	UCG	UAG Stop	UGG Trp	G	
	C	CUU	CCU	CAU	CGU	U	
		CUC	CCC	CAC	CGC	C	
		CUA	CCA	CAA	CGA	A	
		CUG	CCG	CAG	CGG	G	
	A	AUU	ACU	AAU	AGU	U	
		AUC	ACC	AAC	AGC	C	
		AUA	ACA	AAA	AGA	A	
		AUG Met or start	ACG	AAG	AGG	G	
	G	GUU	GCU	GAU	GGU	U	
		GUC	GCC	GAC	GGC	C	
		GUA	GCA	GAA	GGA	A	
		GUG	GCG	GAG	GGG	G	

Translation

- mRNA is the message (the plan for the protein)
- rRNA "reads" the mRNA (the ribosome)
- tRNA molecules carry amino acids to the ribosome for assembly into proteins
- The ribosome allows only the correct tRNA to add its amino acid – others are rejected

Translation



Translation

