


PHY102 : Quiz 1

1. A spherical charge distribution has a density ρ that is constant from $r = 0$ out to $r = R$ and is zero beyond. What is the electric field for all values of r , both less than and greater than R ? [2.5]

Field of a pt charge moving with constant velocity

Take a pt. charge Q at rest at the origin in the frame F .

Electric field is directed radially outwards & at pt P , we have,



$$E_x = \frac{1}{4\pi\epsilon_0} \frac{Q}{r^2} \cos\theta = \frac{1}{4\pi\epsilon_0} \frac{Qx}{(x^2 + z^2)^{3/2}}$$

$$E_z = \frac{1}{4\pi\epsilon_0} \frac{Q}{r^2} \sin\theta = \frac{1}{4\pi\epsilon_0} \frac{Qz}{(x^2 + z^2)^{3/2}}$$


Since, $r = (x^2 + z^2)^{1/2}$ & $\cos\theta = \frac{x}{\sqrt{x^2 + z^2}}$, $\sin\theta = \frac{z}{\sqrt{x^2 + z^2}}$

Consider now a frame F' which is moving in the $-ve$ x direction with speed v , w.r.t frame F .

\therefore To an observer at rest in F' ,

the charge Q is moving in

the $+ve$ x direction with speed v .



The Lorentz transformations that we had written in class were for a frame F' which was moving in the $+ve$ x direction w.r.t frame F .

2. Designate the corners of a square, l on a side, in clockwise order, A, B, C, D . Put charges $2q$ at A and $-3q$ at B . Determine the value of the line integral of \mathbf{E} , from point C to point D . (No actual integration needed!) What is the numerical answer if $q = 10^{-9}C$ and $l = 5cm$? [2.5]

Therefore, in this case, where frame F' is moving in -ve x direction w.r.t. frame F , these transformations become,

$$x = \gamma(x' - vt'), \quad y = y', \quad z = z', \quad t = \gamma(t' - \frac{v}{c^2}x')$$

Note that we are ~~assume~~ going to assume that origin of two frames coincide at time zero according to observers in both frames.

Now, transformations for electric fields are,

$$E_z' = \gamma E_z \quad \& \quad E_x' = E_x$$

\therefore For the instant $t' = 0$, when $x = \gamma x'$, we have,

$$E_x' = E_x = \frac{1}{4\pi\epsilon_0} \frac{Qx}{(x^2 + z^2)^{3/2}} = \frac{1}{4\pi\epsilon_0} \frac{Q(\gamma x')}{[(\gamma x')^2 + z'^2]^{3/2}}$$

$$E_z' = \gamma E_z = \frac{1}{4\pi\epsilon_0} \frac{\gamma Qz}{(x^2 + z^2)^{3/2}} = \frac{1}{4\pi\epsilon_0} \frac{\gamma Qz'}{[(\gamma x')^2 + z'^2]^{3/2}}$$


$$\therefore \frac{E_z'}{E_x'} = \frac{z'}{x'} = \tan\theta'_2$$

\therefore Vector, \vec{E}' makes same angle with x' axis as does the radius vector \vec{r}' !

PHY102 : Quiz 1

1. A spherical charge distribution has a density ρ that is constant from $r = 0$ out to $r = R$ and is zero beyond. What is the electric field for all values of r , both less than and greater than R ? [2.5]

$\therefore \vec{E}'$ points radially outward along a line drawn from the ~~is~~ instantaneous position of Q .



(For implications of this read Purcell, sec. 5.6 & 5.7).

What about the strength of this electric field \vec{E}' ?

$$\begin{aligned} E'^2 &= E_x'^2 + E_z'^2 = \frac{1}{(4\pi\epsilon_0)^2} \cdot \frac{Q^2 (x'^2 + z'^2)}{[(rx')^2 + z'^2]^3} \\ &= \frac{1}{(4\pi\epsilon_0)^2} \cdot \frac{r^2 Q^2 (x'^2 + z'^2)}{\left[\frac{x'^2}{1-\beta^2} + z'^2\right]^3} \quad \left(\because r^2 = \frac{1}{1-\beta^2}\right) \\ &= \frac{1}{(4\pi\epsilon_0)^2} \cdot \frac{r^2 Q^2 (x'^2 + z'^2)}{r^6 [x'^2 + z'^2 - z'^2 \beta^2]^3} \\ &= \frac{1}{(4\pi\epsilon_0)^2} \cdot \frac{1}{r^4} \cdot \frac{Q^2 (x'^2 + z'^2)}{(x'^2 + z'^2)^3 \left[1 - \beta^2 \frac{z'^2}{x'^2 + z'^2}\right]^3} \\ &= \frac{1}{4\pi\epsilon_0} \cdot \frac{Q^2 (1-\beta^4)^2}{(x'^2 + z'^2)^2 \left[1 - \beta^2 \frac{z'^2}{x'^2 + z'^2}\right]^3} \end{aligned}$$

Since, $\sin \theta' = \frac{z'}{(x'^2 + z'^2)^{1/2}}$ & $r' = (x'^2 + z'^2)^{1/2}$

2. Designate the corners of a square, l on a side, in clockwise order, A, B, C, D . Put charges $2q$ at A and $-3q$ at B . Determine the value of the line integral of \mathbf{E} , from point C to point D . (No actual integration needed!) What is the numerical answer if $q = 10^{-9} \text{ C}$ and $l = 5 \text{ cm}$? [2.5]

$$\therefore E' = \frac{1}{(4\pi\epsilon_0)^2} \cdot \frac{Q^2(1-\beta^2)^2}{r'^4 (1-\beta^2 \sin^2 \theta')^3}$$

$$\Rightarrow \boxed{E' = \frac{1}{4\pi\epsilon_0} \cdot \frac{Q}{r'^2} \cdot \frac{1-\beta^2}{(1-\beta^2 \sin^2 \theta')^{3/2}}}$$

For low speeds, $\beta \approx 0 \Rightarrow E' \approx \frac{1}{4\pi\epsilon_0} \frac{Q}{r'^2}$ (expected)

For high enough speeds, because of the $\sin \theta'$ factor, the field is stronger at right angles to motion, ($\theta' = \pi/2$) than in the direction of motion ($\theta' = 0$)!

\therefore field lines concentrated in a pancake \perp to direction of motion. — not spherically symmetric!

Symmetric about a plane \perp to direction of motion of charge

