




Igneous Petrology



Formation of Igneous Rocks



Emplacement of magma below the surface followed by cooling and solidification- **intrusive/plutonic rocks.**

Emplacement of magma at shallow levels followed by cooling and solidification- **hypabyssal rocks.**

Eruption of magma on the surface followed by cooling and solidification- **volcanic rocks.**

A heterogeneous rock body, comprising minerals with different melting points, will undergo **partial melting**, i.e. the minerals with low melting point will melt first followed by the minerals having higher melting point. The melt that is generated and the “residue solid” both have different composition than the parent rock.

Magma type	Ultramafic	Mafic
SiO ₂	42–48	46–54
MgO + FeO + MnO + Fe ₂ O ₃	35–46	15–28
Na ₂ O + K ₂ O	<1	2–3.5
Major minerals	Olivine (generally dominant) + pyroxenes	Plagioclase + pyroxene
Volcanic	Komatiite	Basalt
Hypabyssal	Komatiite	Diabase
Plutonic	Peridotite	Gabbro


Magma type	Intermediate	Felsic
SiO ₂	60–65	>70
MgO + FeO + MnO + Fe ₂ O ₃	10–21	<3
Na ₂ O + K ₂ O	3–6	5–10
Major minerals	Pyroxene + plagioclase + amphibole	Alkali feldspar + quartz
Volcanic	Andesite	Rhyolite
Hypabyssal		
Plutonic	Diorite	Granite



A. Fine-grained



B. Coarse-grained



Extrusive
igneous
rocks

Intrusive
igneous
rocks



C. Glassy (pumice)













D. Porphyritic

The plutonic rocks are usually **coarse-grained** which form at greater depths due to slow cooling.

The volcanic rocks are **fine-grained** formed at the surface by quenching of lava.


The hypabyssal rocks are characterized by porphyritic texture which results due to two-phase cooling- the coarse grains are formed at greater depths (known as **phenocrysts**) and the fine-grained matter (known as **groundmass**) is formed at shallow depths due to faster cooling.

Texture	Composition		
Phaneritic (course-grained)	Felsic (Granitic)	Intermediate (Andesitic)	Mafic (Basaltic)
	 <p data-bbox="556 539 653 564">Granite</p>	 <p data-bbox="1025 539 1103 564">Diorite</p>	 <p data-bbox="1450 539 1528 564">Gabbro</p>
	 <p data-bbox="556 903 653 928">Rhyolite</p>	 <p data-bbox="1016 903 1112 928">Andesite</p>	 <p data-bbox="1460 903 1537 928">Basalt</p>
Porphyritic	 <p data-bbox="513 1268 697 1292">Granite porphyry</p>	 <p data-bbox="967 1268 1170 1292">Andesite porphyry</p>	 <p data-bbox="1412 1268 1595 1292">Basalt porphyry</p>




The sequential crystallization of minerals due to cooling of magma. In the discontinuous series, the high temperature minerals are eventually consumed and forms the next mineral in the series (for e.g. for a certain span of time, olivine and pyroxene will coexist and as the temperature further drops, olivine will be consumed completely leaving behind pyroxene). However, the plagioclase feldspar has a different pattern of evolution wherein the Ca is replaced by Na without forming any distinctly different mineral.

Bowen's Reaction in action





An olivine crystal surrounded by pyroxene in an extrusive (volcanic) igneous rock.



A zoned crystal of plagioclase in an igneous rock. The center (core) is Ca-rich and the edge (rim) is Na-rich.

Structures of Igneous Rocks- Plutonic Bodies



Sill is a concordant body formed by the penetration of magma parallel to the bedding planes. Basic magmas have low viscosity hence can flow easily and mostly occur as sills. **Dykes** are discordant bodies that cut across bedding planes. Sills and dykes are formed when during upward journey of magma, weak planes such as fractures, joints or crevices are encountered. **Laccolith** is a concordant igneous body with a flat base and convex upward top. When viscous magma rises through country rock, laccoliths are formed since the magma cannot flow easily and through longer distances and accumulates forming the laccoliths. Laccoliths particularly form when the supply of magma from below is more than can be accommodated by lateral spreading. When sedimentary rocks are folded, the crests and troughs may be occupied by magmatic intrusions which look like crescentic or lens shaped bodies known as **phacoliths**. **Batholiths** are largest intrusive igneous bodies that characteristically occur in mountain regions and are considered as roots of mountains. They extend at great depths and since these are deep-seated bodies, they might be exposed only after prolonged erosion.

Devil's Tower- A Laccolith or Volcanic Neck?

