

Assignment 11

①

$$1. \quad v_1 = (1, -1), \quad v_2 = (2, -1), \quad v_3 = (-3, 2) \\ w_1 = (1, 0), \quad w_2 = (0, 1), \quad w_3 = (1, 1)$$

$$\text{If } Tv_1 = w_1 \text{ and } Tv_2 = w_2$$

$$\text{then } Tv_1 + Tv_2 = w_1 + w_2 = w_3$$

$$\Rightarrow T(v_1 + v_2) = w_3.$$

~~But~~ On the other hand

$$(-3, 2) = v_3 = -v_1 - v_2 \\ = (-1, 1) + (-2, 1).$$

$$\therefore Tv_3 = -w_1 - w_2 = -w_3$$

Hence there cannot exist a linear transformation $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ st $T(v_i) = w_i$,
i=1, 2, 3.

$$2. \quad \text{Let } T: \mathbb{R}^2 \rightarrow \mathbb{R}^2 \text{ be st}$$

$$T(1, 0) = (a, b) \text{ and } T(0, 1) = (c, d).$$

Then for any $(x_1, x_2) \in \mathbb{R}^2$

$$\begin{aligned} T(x_1, x_2) &= x_1 T(1, 0) + x_2 T(0, 1) \\ &= x_1 (a, b) + x_2 (c, d) \\ &= (ax_1 + cx_2, bx_1 + dx_2) \end{aligned}$$

3. (a). $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$, is defined by

(2)

$$T(x_1, x_2, x_3) = (x_1 - x_2 + 2x_3, 2x_1 + x_2, -x_1 - 2x_2 + 2x_3)$$

A vector $(a, b, c) \in \mathbb{R}^3$ is in the range of T if the system of linear eqns,

$$\begin{cases} x_1 - x_2 + 2x_3 = a \\ 2x_1 + x_2 = b \\ -x_1 - 2x_2 + 2x_3 = c \end{cases}$$

has a solution. In other words, $(a, b, c) \in \text{Range } T$ if the ~~aug~~ row-reduced form of the augmented matrix is consistent. i.e

if the row-reduced form of $\left(\begin{array}{ccc|c} 1 & -1 & 2 & a \\ 2 & 1 & 0 & b \\ -1 & -2 & 2 & c \end{array} \right)$ is consistent.

After a sequence of ~~row~~ elementary row-operations the above matrix reduces to,

$$(R:a) = \left(\begin{array}{ccc|c} 1 & 0 & 2/3 & \frac{b+a}{3} \\ 0 & 1 & -4/3 & \frac{b-2a}{3} \\ 0 & 0 & 0 & c+b-a \end{array} \right)$$

and clearly $(R:a)$ is consistent if $c+b-a=0$.

Hence $(a, b, c) \in \text{Range } T$ if $b+c-a=0$.

\Rightarrow vectors in $\text{Range } T$ are of the form $(b+c, b, c) = b(1, 1, 0) + c(1, 0, 1)$
 $\Rightarrow \text{Rank } T = 2$.

(b). If $(a, b, c) \in \mathbb{R}^3$ is in the null space of T (3)
then

$$T(a, b, c) = (0, 0, 0)$$

$$\Rightarrow \Rightarrow (a - b + 2c, 2a + b, -a - 2b + 2c) = (0, 0, 0)$$

$$\Rightarrow \begin{pmatrix} 1 & -1 & 2 \\ 2 & 1 & 0 \\ -1 & -2 & 2 \end{pmatrix} \begin{pmatrix} a \\ b \\ c \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}.$$

Row-reducing the coef matrix we get

$$\begin{pmatrix} 1 & 0 & 2/3 \\ 0 & 1 & -4/3 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} a \\ b \\ c \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

$$\Rightarrow \begin{aligned} a + 2/3 c &= 0 \\ b - 4/3 c &= 0. \end{aligned}$$

$$\Rightarrow (a, b, c) \in \text{Null space of } T \text{ if } a = -\frac{2}{3}c, b = \frac{4}{3}c$$

this implies that

$$\left(-\frac{2}{3}, \frac{4}{3}, 1\right) \mathbb{R} = \text{Null space of } T.$$

$$\Rightarrow \text{Nullity of } T = 1.$$