

Information-Theoretic Bell Inequalities

QCQI Phy631 Term Paper

Dhruva Sambrani

Local Realism

Locality

It is the belief that systems which are “far away” in space from our system will have no effect.

This assumption is the first assumption in Special Relativity too. This leads to a finite speed for causality to travel, which is the speed of light.

Philosophically too this seems to be a requirement of basic Physics, after all, what is the use of any of our laws if another particle on the other side of the universe can affect your particle.

Realism

This is the assumption that objects have properties whether or not said properties are measured. Is the moon there if you are not looking at it?

This too has philosophical implications. If no property of an object is real and is instead only an outcome of our measurement, what are we describing in Physics? Isn't the world then dependant on the observer and not the observed?

Some Mathematical formalism

Bayes' Theorem

$$p(a, b) = p(a|b)p(b) = p(b|a)p(a)$$

Information

$$\mathcal{I}(p) = -\log(p)$$

$$\mathcal{I}(p(a, b)) = -\log(p(a, b))$$

$$\mathcal{I}(p(b)) = -\log(p(b))$$

$$\mathcal{I}(p(a|b)) = -\log(p(a|b))$$

Then, from Bayes' Theorem,

$$\begin{aligned} -\log p(a, b) &= -\log(p(a|b)p(b)) = -\log(p(b|a)p(a)) \\ \implies \mathcal{I}(a, b) &= \mathcal{I}(a|b) + \mathcal{I}(b) = \mathcal{I}(b|a) + \mathcal{I}(a) \end{aligned}$$

Mean information

$$\mathcal{H}(A, B) = \sum_{a,b} p(a, b) \mathcal{I}(a, b)$$

This is the Shannon Entropy!

Similar to the Information Bayes' Theorem,

$$\mathcal{H}(A, B) = \mathcal{H}(A|B) + \mathcal{H}(B) = \mathcal{H}(B|A) + \mathcal{H}(A)$$

Mutual Information

$$\mathcal{I}(a; b) = \mathcal{I}(a) - \mathcal{I}(a|b) = \mathcal{I}(b) - \mathcal{I}(b|a) = \mathcal{I}(b; a)$$

$$\mathcal{H}(A; B) = \mathcal{H}(A) - \mathcal{H}(A|B) = \mathcal{H}(B) - \mathcal{H}(B|A) = \mathcal{H}(B; A)$$

Two Inequalities of Information Theory

$$\mathcal{H}(A|B) \leq \mathcal{H}(A) \leq \mathcal{H}(A, B)$$

The left-hand inequality means that removing a condition never decreases the information carried by a quantity. The right-hand inequality means that two quantities never carry less information than either quantity carries separately.

Bell's Inequality

Bell's Inequality

The simplest one

Setup

Consider two systems, \mathcal{A} and \mathcal{B} , with the set of measurable quantities A and A' , and, B and B' respectively.

A and A' (and by extension, B and B') may not, in general, commute, so cannot be measured simultaneously. Hence a run of the experiment will involve measuring **only one** of either A or A' .

Further, also assume that the systems \mathcal{A} and \mathcal{B} , are “far away” in space.

Properties of Observables

This proof of no hidden variable theory being able to explain Quantum Mechanics is one by contradiction.

Let us assume the observables(A and its friends) are all real and local. To recap, this means that the systems “actually” have a value for each property. That is \mathcal{A} is $|a, a'\rangle$ ¹(and similarly for \mathcal{B}).

Further, any measurements on \mathcal{A} should not effect \mathcal{B} and vice versa, as these systems are well separated and properties are local.

¹Abuse of notation - While normally, $|a, a'\rangle$ is used to denote a *simulataneous* eigenvector, here, we're using it to represent a state which “actually” has a and a' as values for A and A'

What we know

the

Though we can not measure A and A' separately, we know $p(a, a', b, b')$ from which follow the 4 pairwise probabilities in system \mathcal{A} and \mathcal{B} .

For example, $p(a, b) = \sum_{a', b'} p(a, a', b, b')$