

KNOW HOW TO USE YOUR DEVICE

Gliptozza

6mg/ml
PRE-FILLED PEN

You may have concerns about using an injectable medication for managing your type 2 diabetes. But it's important to realize the positive effect it may have on the management of your diabetes. And once you gain a little confidence in using them on your own, Gliptozza pen can become part of your daily routine.



Gliptozza should be taken daily exactly as prescribed by your Health Care Provider. It's best to get into a routine so you remember to take Gliptozza once a day, at the same time every day.

You should store it in a refrigerator at a temperature between 2°C to 8°C. Protect from light. Do not freeze.

Self-injecting with the Gliptozza pen

Inject your dose of Gliptozza under the skin (this is called a subcutaneous injection) in one of these injection sites: your stomach area (abdomen), upper leg (thigh), or upper arm, as instructed by your health care provider. Do not inject into a vein or muscle.

Make sure you read about and understand the proper care and storage of your Gliptozza pen. Refer leaflet enclosed in the pack.




KNOW HOW TO USE YOUR DEVICE


Gliptozza

6mg/ml
PRE-FILLED PEN

Technique to use Gliptozza Pen



Change (rotate) your injection site within the area you choose with each injection to reduce your risk of getting lumps under the skin (cutaneous amyloidosis). Do not use the same site for each injection.



- You may give an injection of Gliptozza and insulin in the same body area (such as your stomach area), but not right next to each other.

Do not take 2 doses of Gliptozza at the same time.

- If you take too much Gliptozza, call your healthcare provider right away.
- Taking too much Gliptozza may cause severe nausea, severe vomiting, and low blood sugar (hypoglycemia).
- Do not share your Gliptozza pen with other people, even if the needle has been changed. You may give other people a serious infection or get a serious infection from them.
- The Gliptozza pen you are using should be disposed as per local regulations 30 days after you start using it. Your dose of Gliptozza and other diabetes medicines may need to change because of:
- Change in level of physical activity or exercise, weight gain or loss, increased stress, illness, change in diet, or because of other medicines you take. So visit your doctor regularly as suggested by them to avoid any complication.
- Use New Needle for each dose