

NOTES ON LAMBDA CALCULUS

VAIBHAV KARVE

These notes were last updated September 13, 2018. They are notes taken from my reading of *Haskell Programming from First Principles* by *Chris Allen*, *Julie Moronuki*. I plan on expanding these notes further by reading the following at some unspecified time in the future:

- A tutorial introduction to the Lambda Calculus by *Raúl Rojas*.
- Introduction to Lambda Calculus by *Henk Barendregt* and *Erik Barendsen*.
- Proofs and Types by *Jean-Yves Girard*, *Paul Taylor* and *Yves Lafont*.

CONTENTS

1. Basics and definitions	1
2. Equivalences and reductions	1
3. Examples	3

1. BASICS AND DEFINITIONS

- (1) Lambda calculus has three basic components or *lambda terms* – expressions, variables and abstractions.
- (2) *Expressions* are variable names, abstractions, or combinations of other expression. *Variables* have no meaning or value, they are only names for potential inputs to functions. An *abstraction* is a function – it is a lambda term that has a head (a lambda) and a body and is applied to an argument. An *argument* is an input value.
- (3) Abstractions have two parts – a *head* and a *body*. The head of the function is a λ followed by a variable name. The body of the function is another expression. For example: $\lambda x..x^2$
Lambda abstractions are anonymous functions.
- (4) The variable named in the head is the *parameter* and *binds* all instances of that same variable in the body of the function. The dot (.) separates the parameters of the lambda from the function body.

2. EQUIVALENCES AND REDUCTIONS

- (1) *Alpha equivalence* states that $\lambda x..x$ is the same as $\lambda y..y$, that is, the variables x and y are not semantically meaningful except in their role in their single expressions.

- (2) *Beta reduction*: when applying a function to an argument, substitute the input expression for all instances of bound variables within the body of the abstraction.

$$(\lambda x.xx)3 = xx[x := 3] = 3\ 3$$

Hence, Beta reduction is the process of applying a lambda term to an argument, replacing the bound variables with the value of the argument, and eliminating the head.

$$\begin{aligned} (\lambda x.x)\lambda y.y &= x[x := (\lambda y.y)] \\ &= \lambda y.y \end{aligned}$$

- (3) Application in lambda calculus is left-associative.

$$\begin{aligned} (\lambda x.x)(\lambda y.y)z &= ((\lambda x.x)(\lambda y.y))z && \text{left-associativity} \\ &= (x[x := \lambda y.y])z && \text{beta reduction step 1} \\ &= (\lambda y.y)z && \text{beta reduction step 2} \\ &= y[y := z] && \text{beta reduction step 1} \\ &= z && \text{beta reduction step 2} \end{aligned}$$

- (4) Variables in the body that are not bound by the head are called *free variables*. For example, y is a free variable in the expression $\lambda x.xy$

$$(\lambda x.xy)z = xy[x := z] = zy$$

- (5) The alpha equivalence does not apply to free variables.

- (6) *Currying*: named after Haskell Curry is the shorthand notation of the type $\lambda xy...xy$ for multiple lambda functions $\lambda x.(\lambda y.xy)$.

$$\begin{aligned} \lambda xy.xy\ 1\ 2 &= \lambda x.(\lambda y.xy)\ 1\ 2 \\ &= (\lambda y.xy)[x := 1]\ 2 \\ &= (\lambda y.1y)\ 2 \\ &= (1y)\ [y := 2] \\ &= 1\ 2 \end{aligned}$$

or by using currying we perform the same calculation in fewer steps,

$$\begin{aligned} \lambda xy.xy\ 1\ 2 &= (\lambda y.xy)[x := 1]\ 2 \\ &= (\lambda y.1y)2 \\ &= (1y)[y := 2] \\ &= 1\ 2 \end{aligned}$$

- (7) A lambda term is in *beta normal form* when one cannot beta reduce (apply lambdas to arguments) its expressions any further. This corresponds to a fully evaluated function or fully executed program. The identity function $\lambda x.x$ is in normal form.
- (8) A *combinator* is a lambda term with no free variables. Combinators serve only to combine the arguments that are given. The following are combinators: $\lambda x.x$, $\lambda xy.x$, $\lambda xyz.xz(yz)$ and the following are not: $\lambda y.x$, $\lambda x.xz$. The point of combinators is that they can only combine the arguments they are given, without injecting any new values or random data.
- (9) A lambda term whose beta reduction never terminates is said to *diverge*. The lambda term *omega* defined as $(\lambda x.xx)(\lambda x.xx)$ diverges because

$$(\lambda x.xx)(\lambda x.xx) = (\lambda x.xx)(\lambda y.yy) = xx[x := \lambda y.yy] = (\lambda y.yy)(\lambda y.yy).$$

3. EXAMPLES

$$\begin{array}{lll}
(\lambda xy.xy)(\lambda z.a) \ 1 & (\lambda xyz.xz(yz))(\lambda x.z)(\lambda x.a) & (\lambda y.y)(\lambda x.xx)(\lambda z.zq) \\
= (\lambda y.(\lambda z.a)y)1 & = (\lambda xyb.xb(yb))(\lambda c.z)(\lambda d.a) & = (\lambda x.xx)(\lambda z.zq) \\
= (\lambda z.a)1 & = (\lambda yb.(\lambda c.z)b(yb))(\lambda d.a) & = (\lambda z.zq)(\lambda z.zq) \\
= a & = \lambda b.(\lambda c.z)b((\lambda d.a)b) & = (\lambda z.zq)(\lambda x.xq) \\
& = \lambda b.z((\lambda d.a)b) & = (\lambda x.xq)q \\
& = \lambda b.za & = qq \\
\\
(\lambda a.aa)(\lambda b.ba)c & (\lambda xyz.xz(yz))(\lambda mn.m)(\lambda p.p) & (\lambda xy.xxy)(\lambda x.xy)(\lambda x.xz) \\
= (\lambda d.dd)(\lambda b.ba)c & = (\lambda yz.(\lambda mn.m)z(yz))(\lambda p.p) & = (\lambda xy.xxy)(\lambda a.ay)(\lambda b.bz) \\
= (\lambda b.ba)(\lambda b.ba)c & = \lambda z.(\lambda mn.m)z((\lambda p.p)z) & = (\lambda y.(\lambda a.ay)(\lambda c.cy)y)(\lambda b.bz) \\
= (\lambda b.ba)(\lambda d.da)c & = \lambda z.(\lambda n.z)((\lambda p.p)z) & = (\lambda a.a(\lambda b.bz))(\lambda c.c(\lambda b.bz))(\lambda b.bz) \\
= ((\lambda d.da)a)c & = \lambda z.z & = (\lambda a.a(\lambda b.bz))(\lambda c.c(\lambda d.dz))(\lambda e.ez) \\
= aac & & = ((\lambda c.c(\lambda d.dz))(\lambda b.bz))(\lambda e.ez) \\
& & = ((\lambda b.bz)(\lambda d.dz))(\lambda e.ez) \\
& & = ((\lambda d.dz)z)(\lambda e.ez) \\
& & = (zz)(\lambda e.ez) \\
& & = yy(\lambda b.bz) \\
\\
(\lambda x.\lambda y.xyy)(\lambda a.a)b & (\lambda abc.cba)zz(\lambda wv.w) & \\
= (\lambda y.(\lambda a.a)yy)b & = (\lambda bc.cbz)z(\lambda wv.w) & \\
= (\lambda a.a)bb & = (\lambda c.czz)(\lambda wv.w) & \\
= bb & = (\lambda wv.w)zz & \\
& = (\lambda v.z)z & \\
& = z &
\end{array}$$