1

11.14-4

EE23BTECH11048-Ponugumati Venkata Chanakya*

QUESTION: Which of the following functions of time represent (a) simple harmonic, (b) periodic but not simple harmonic, and (c) non-periodic motion? Give period for each case of periodic motion (ω is any positive constant):

- 1) $\sin(\omega t) \cos(\omega t)$
- 2) $\sin^3(\omega t)$
- 3) $3\cos\left(\frac{\pi}{4}-2\omega t\right)$
- 4) $\cos(\omega t) + \cos(3\omega t) + \cos(5\omega t)$
- 5) $\exp(-\omega^2 t^2)$
- 6) $1 + \omega t + \omega^2 t^2$

Solution::

1) Periodic function:

$$x(t+T) = x(t) \quad \forall x \in \mathbb{R} \tag{1}$$

where min T s.t T > 0 is time period

2) SHM:

For a function to be in shm it must satisfy

$$\frac{d^2x(t)}{dt^2} = -(2\pi f_0)^2 x(t) \tag{2}$$

Variable	Description	formula	
x(t)	Displacemen wrt mean position	none	
ω	Angular frequncy	$2\pi f$	
T	Time period	$\frac{1}{f}$	
φ	phase angle	none	
	TABLE 0	•	

INPUT PARAMETERS

1) $\sin(2\pi ft) - \cos(2\pi ft)$

The function can be rewritten as:

$$= \sin(2\pi f t) - \sin\left(\frac{\pi}{2} - 2\pi f t\right) \tag{3}$$

$$= 2\cos\left(\frac{\pi}{4}\right)\sin\left(2\pi ft - \frac{\pi}{4}\right) \tag{4}$$

$$=\sqrt{2}\sin\left(2\pi ft - \frac{\pi}{4}\right) \tag{5}$$

$$\frac{d^2(\sin(2\pi ft) - \cos(2\pi ft))}{dt^2} = -(2\pi f)^2(\sin(2\pi ft) - \cos(2\pi ft))$$
 (6)

$$\frac{d^2x(t)}{dt^2} = -(2\pi f)^2 x(t) \tag{7}$$

(8)

 \therefore SHM, T is $\frac{1}{f}$ and ϕ is $\left(\frac{-\pi}{4}\right)$ or $\left(\frac{7\pi}{4}\right)$

$$\sin\left(2\pi f\left(t+\frac{1}{f}\right)\right) - \cos\left(2\pi f\left(t+\frac{1}{f}\right)\right) = \sin(2\pi ft) - \cos(2\pi ft) \tag{9}$$

Graph of function is shown in (Fig. 1)

(3) $3\cos\left(\frac{\pi}{4}-4\pi ft\right)$

This function can be rewritten as

$$=3\cos\left(4\pi ft - \frac{\pi}{4}\right)\tag{10}$$

$$\frac{d^2\left(3\cos\left(\frac{\pi}{4} - 4\pi ft\right)\right)}{dt^2} = -3(4\pi f)^2\left(\cos\frac{\pi}{4} - 4\pi ft\right) \tag{11}$$

$$\frac{d^2x(t)}{dt^2} = -(4\pi f)^2 x(t) \tag{12}$$

 \therefore SHM, T is $\frac{1}{2f}$ and ϕ is $\left(\frac{-\pi}{4}\right) or \left(\frac{7\pi}{4}\right)$

$$3\cos\left(\frac{\pi}{4} - 4\pi f\left(t + \frac{1}{2f}\right)\right) = 3\cos\left(\frac{\pi}{4} - 4\pi f t\right) \tag{13}$$

(14)

Graph of function is shown in (Fig. 1)

(4) $\cos(2\pi ft) + \cos(6\pi ft) + \cos(10\pi ft)$

This function can be rewritten as

$$= \cos(2\pi ft) + \cos(10\pi ft) + \cos(6\pi ft) \tag{15}$$

$$= 2\cos\left(\frac{2\pi ft + 10\pi ft}{2}\right)\cos\left(\frac{10\pi ft - 2\pi ft}{2}\right) + \cos(6\pi ft) \quad (16)$$

$$= 2\cos(6\pi ft)\cos(2\pi ft) + \cos(6\pi ft) \tag{17}$$

$$= \cos(6\pi f t)(1 + 2\cos(2\pi f t)) \tag{18}$$

$$\frac{d^2\cos(2\pi ft) + \cos(6\pi ft) + \cos(10\pi ft)}{dt^2} = (2\pi f)^2\cos(2\pi ft) + (6\pi f)^2\cos(6\pi ft) + (10\pi f)^2\cos(10\pi ft)$$
(19)

$$\frac{d^2x(t)}{dt^2} \neq -(2\pi f)^2 x(t) \tag{20}$$

Period of $\cos(6\pi ft)$ is $\frac{1}{3f}$ Period of $1 + 2\cos(2\pi ft)$ is $\frac{1}{f}$ Lcm is $\frac{1}{f}$ \therefore SHM,T is $\frac{1}{f}$

$$\cos\left(2\pi f\left(t+\frac{1}{f}\right)\right) + \cos\left(6\pi f\left(t+\frac{1}{f}\right)\right) + \cos\left(10\pi f\left(t+\frac{1}{f}\right)\right) = \cos(2\pi ft) + \cos(6\pi ft) + \cos(10\pi ft)$$
(21)

Graph of function is shown in (Fig. 1)

(5) $\exp(-(2\pi f)^2 t^2)$

As
$$T \to \infty$$
 (22)

$$\exp\left(-(2\pi f)^2 t^2\right) \to \infty \tag{23}$$

$$\frac{d^2(\exp\left(-(2\pi f)^2 t^2\right)))}{dt^2} = 2(2\pi f t)^2 \exp\left(-(2\pi f)^2 t^2\right) + 2(2\pi f t)^4 \exp\left(-(2\pi f)^2 t^2\right)$$
(24)

$$\frac{d^2x(t)}{dt^2} \neq -(2\pi f)^2 x(t)$$
 (25)

.. This never repeats and non periodic Graph of function is shown in (Fig. 1)

(6)
$$1 + 2\pi f t + (2\pi f)^2 t^2$$

As
$$T \to \infty$$
 (26)

$$1 + 2\pi f t + (2\pi f)^2 t^2 \to \infty \tag{27}$$

$$\frac{d^2(1+2\pi ft+(2\pi f)^2t^2)}{dt^2}=2(2\pi f)^2\tag{28}$$

$$\frac{d^2x(t)}{dt^2} \neq -(2\pi f)^2 x(t)$$
 (29)

∴ This never repeats and non periodic Graph of function is shown in (Fig. 1)

TABLE 1 Summary

	Function	Periodic	Simple harmonic motion	Non Periodic	Т	φ
(a)	$\sin(2\pi ft) - \cos(2\pi ft)$	Yes	Yes	No	$\frac{1}{f}$	$\left(\frac{-\pi}{4}\right)$
(b)	$\sin^3(2\pi ft)$	Yes	No	No	$\frac{1}{f}$	_
(c)	$3\cos\left(\frac{\pi}{4}-4\pi ft\right)$	Yes	Yes	No	$\frac{1}{2f}$	$\left(\frac{-\pi}{4}\right)$
(d)	$\cos(2\pi ft) + \cos(6\pi ft) + \cos(10\pi ft)$	Yes	No	No	\overline{f}	_
(e)	$\exp\left(-(2\pi f t)^2\right)$	No	No	Yes	_	_
(f)	$1 + (2\pi f)t + (2\pi f)^2 t^2$	No	No	Yes	_	-




Fig. 1. $\sin(2\pi ft) - \cos(2\pi ft)$




Fig. 1. $\sin^3(2\pi ft)$




Fig. 1. $3\cos\left(\frac{\pi}{4} - 4\pi ft\right)$




Fig. 1. $\cos(2\pi f t) + \cos(6\pi f t) + \cos(10\pi f t)$




Fig. 1. $exp^{(-(2\pi ft)^2)}$




Fig. 1. $1 + 2\pi f t + (2\pi f t)^2$