

Little Red Dot Density and Temperature Prediction Using Triplet Helium Strengths

AKSHAJ VYAS¹ AND IAN BISHOP²

¹*Lake Travis High School*

²*Basis Cedar Park School*

(Dated: July 2026)

1. INTRODUCTION

1.1. Background

Compared to previous telescopes, the James Webb Space Telescope (JWST) demonstrated a higher resolution and greater observational depth at infrared wavelengths than any previous space-based or earth-based telescope. This led to the discovery of Little Red Dots (LRDs) within JWST deep-field images from the CEERS survey (S. L. Finkelstein et al. 2025). LRDs are present in all JWST deep field images, especially at redshifts of $4 < z < 9$ (V. Kokorev et al. 2025). LRDs were deemed noteworthy due to a unique "v-shaped" spectral distribution, with opposite rest-UV and rest-optical slopes (J. Matthee et al. 2024) (D. D. Kocevski et al. 2025). Further spectroscopy revealed the presence of unusual Balmer line broadening within the spectral distribution (D. D. Kocevski et al. 2025) (A. J. Taylor et al. 2025).

LRDs were originally hypothesized to be large early-universe galaxies, though a contradiction arose, where the predicted stellar masses within the galaxies were unexpectedly high compared to prior observations (I. Labbé et al. 2023). Further spectroscopic investigations demonstrated that the spectral distributions of LRDs were dominated by supermassive black hole (SMBH) emission instead of stellar emission (D. D. Kocevski et al. 2025) (A. J. Taylor et al. 2025). The most recent studies have explained the Balmer line broadening by introducing the idea a dense cocoon of excited gas surrounding the SMBH (R. P. Naidu et al. 2025) (V. Rusakov et al. 2026).

Previous work has focused on Balmer emission lines to estimate the physical properties of LRDs. This work substitutes Balmer emission lines with Helium emission lines to predict the temperature and density of the hypothesized cocooning gas enshrouding LRDs. Helium emission lines show promise in being able to predict physical properties, as they exhibit similar broadening and absorption features as Balmer lines, indicating a direct relationship between the cocooning gas and the Helium flux (I. Juodbalis et al. 2024) (B. . Wang et al. 2025) (V. Kokorev et al. 2025) (A. de Graaff et al. 2025).

1.2. Goals

To estimate the temperature and density of LRDs, we perform spectral analysis of four critical emission lines: He I 10830Å, H I $P\alpha\gamma$, He I 5876Å, and He I 7065Å. Additionally, we lay the foundation for future simulations which can provide significantly more accurate estimates of physical characteristics, through more precise models. Understanding the temperature and density of LRDs is critical to determining the structure of these objects, and could eventually result in a conclusive theory for LRDs.

1.3. Helium

The metastable He I helium line 10830Å emerges when helium transitions from $1s2p^3P_J^o$ to $1s2s^3S_1$. The He I 10830Å metastable transition is unique as its lower metastable level, known as the He I 2^3S_1 state, is substantially more stable than other non-ground state helium. This is due to the decay being forbidden by quantum mechanics, leading to an extremely low probability of transition. When isolated, a helium atom can remain in the 2^3S_1 state for about $7.8 \cdot 10^3 s$ (Y. R. Sun & S.-M. Hu 2020). This property of metastable helium creates a cycle between emission and absorption of free electrons, causing a strong correlation between the

strength of the emission line and electron density. We take advantage of this property to accurately estimate the density of LRDs.

We compare the He I 10830Å line to the H I $Pa\gamma$ line due to its consistency among the LRD sample and close proximity to 10830Å, which eliminates the requirement for dust correction. We also use H I $Pa\gamma$ due to its relatively low sensitivity to density changes compared to H I 10830Å, making it a good indicator of correlation between H I 10830Å and density. (E. Aver et al. 2015)

As for the He I 5876Å and 7065Å, these emerge when Helium transitions to 2^3S from the states 3^3D and 3^3S , respectively. He I 7065Å was chosen as it is highly sensitive to electron density. This is because the metastable Helium is easily excited to the 3^3S state, leading to the population of metastable Helium being intertwined with the population of 3^3S Helium. This leads to both He I 7065Å and He I 10832Å being correlated with electron density. He I 5876 was chosen as it was the most common line among the sampled LRDs.




Figure 1. Figure 3 from D. A. Berg et al. (2025) describes the transition between every Helium state, and particularly emphasizes the longevity of the metastable state.

2. METHODS

2.1. Data

The 1-dimensional extractions used for this work are courtesy of the OCEANS collaboration (K. Davis et al. 2026). The extractions utilized the NIRSpec instrument from the JWST, probing the rest-optical observed-infrared light from the LRD. We configured the telescope to the high-resolution G235H and G395H gratings, both of which covered ranges in the single-digit microns. For our metastable helium analysis, we obtained precalculated narrow and broad fluxes for the He I 10830Å line from I. Juodbalis et al. (2024), B. Wang et al. (2025), V. Kokorev et al. (2025), and A. de Graaff et al. (2025).

Before we performed spectral analysis, we cleaned the data to prepare it for flux measurement. First, we converted the units of μm to Å and Jy (or μJy) to $\text{erg} \cdot \text{cm}^{-2} \text{s}^{-1} \text{Å}^{-1}$. This conversion leads to LiME providing integrated flux measurements in $\text{erg} \cdot \text{cm}^{-2} \text{s}^{-1}$. Next, we adjusted the wavelength for redshift

using the formula $\lambda_{rest} = \frac{\lambda_{obs}}{1+z}$, where z represents the redshift. The wavelength was corrected utilizing the Python library NumPy (C. R. Harris et al. 2020). Although the single-digit micron range exceeds the wavelength range of visible and near-infrared light, correcting for redshift reveals the appropriate emission lines.

2.2. Flux Measurement

We use the Python library LiME (Line Measuring library) to measure fluxes V. Fernández et al. (2024). We measured narrow fluxes from the observed data for He I 5876Å, He I 7065Å, and H I 10941Å. Because He I 10830Å had a complex line shape with broad and narrow emission superimposed with absorption, we used pre-aggregated tables from prior research papers (I. Juodbalis et al. 2024; B. . Wang et al. 2025; V. Kokorev et al. 2025; A. de Graaff et al. 2025) to avoid repeating the complicated work.

Next, we fit a continuum to the spectrum using a cubic equation. To eliminate emission lines from the continuum, we iteratively refit the cubic while clipping emission lines past a σ threshold above the continuum. As the iterations progress, we gradually reduce the threshold from 3σ to 1σ . Then, we performed line detection, where we utilized a neighbor comparison algorithm to detect all extrema relative to the continuum, and compared the detected extrema to LiME's internal database of known spectral lines. If a spectra excluded any of the lines, it was removed as a non-detection. The continuum fitting and line detection were implemented by LiME. However, tuning the parameters to the values mentioned was essential in order to achieve accurate line predictions. The accuracy of the line predictions was determined by visually inspecting the plots to ensure most peaks and valleys were mapped to an appropriate emission line.

Next, we measured fluxes through profile fitting. For the He I 5876Å, 7065Å, and H I Pa γ lines, the flux was best modeled by a Gaussian profile without broadening or blending. The Gaussian was fit through a least squares regression. We measured the flux by integrating over this Gaussian, taking into account the continuum as the floor for integration. The criteria for a false positive was a 3σ detection for Pa γ and a 2σ detection for other lines. Distributions which failed to meet this threshold were removed as non-detections.

For He I 10830Å, the fluxes were loaded from previous spectroscopic papers. While Wang et al precalculated the total flux, every other source provided only the narrow, broad and absorption flux. To find the total flux, the narrow and broad flux were added together, while the absorption was reintroduced into the line profile. We calculated the error based on formulas detailed in the next section.

Finally, we calculated $\frac{F_{5876\text{\AA}}}{F_{7065\text{\AA}}}$, as well as $\frac{F_{10830\text{\AA}}}{F_{10941\text{\AA}}}$, and recorded these values for future comparison. The values were recorded and stored in a .csv file using the Python library pandas (T. pandas development team 2020).

Table 1. Flux value table of each LRD with both the He I 5876Å and 7065Å line. Flux was measured in units of $10^{19} \cdot \text{erg} \cdot \text{cm}^{-2} \cdot \text{s}^{-1}$

LRD	He I 7065Å flux	5876Å flux
OCEANS 20504 P4	3.67 ± 0.72	8.49 ± 0.10
OCEANS 20504 P1	8.51 ± 1.14	4.79 ± 0.95
OCEANS 169045 P4	7.52 ± 0.68	3.84 ± 0.57
OCEANS 102364 P5	2.98 ± 0.83	2.96 ± 0.53

Table 2. Flux value table of each LRD with both the He I 10830Å and $Pa\gamma$ lines. Flux was measured in units of $10^{18} \cdot \text{erg} \cdot \text{cm}^{-2} \cdot \text{s}^{-1}$

LRD	He I 10830Å flux	$Pa\gamma$ flux
JADES 28074 G235M	$82.11^{+1.303}_{-1.400}$	22.1 ± 1.27
JADES 28074 G395M	$82.11^{+1.303}_{-1.400}$	21.2 ± 1.69
RUBIES 40579 G395M	70^{+4}_{-3}	40.5 ± 4.23
RUBIES 154183 G395M	$2.7^{+5.8}_{-5.8}$	2.74 ± 1.44
GLIMPSE 17775 G395M	$24.58^{+2.334}_{-2.334}$	5.16 ± 0.29

2.3. Flux Error Calculation

The error propagation is derived from the LiME measurement, and represents the error of the fit. Flux error was calculated simultaneously with flux values. Whenever operations were performed on flux values, we performed the corresponding quadrature operation on error. For addition and subtraction, we use the following formula:

$$\sigma_z = \sqrt{\sigma_x^2 + \sigma_y^2}$$

This formula was specifically used when adding the broad and narrow flux to calculate the total positive and negative error for He I 10830Å:

$$\sigma_{\text{He, total}} = \sqrt{\sigma_{\text{He, narrow}}^2 + \sigma_{\text{He, broad}}^2}$$

For error derived from the multiplication or division of flux, we instead add the relative errors in quadrature:

$$\frac{\sigma_z}{z} = \sqrt{\left(\frac{\sigma_x}{x}\right)^2 + \left(\frac{\sigma_y}{y}\right)^2}$$

We utilized this formula when calculating flux ratios between different flux measurements to calculate the total error of our ratios:

$$\sigma_R = R \sqrt{\left(\frac{\sigma_{F_1}}{F_1}\right)^2 + \left(\frac{\sigma_{F_2}}{F_2}\right)^2}$$

2.4. Dust Correction

We shifted the line fluxes to reverse the effects of dust extinction on the emission spectra. This applies only to the 5876Å and 7065Å emission lines. For the other ratio, the relative effect of dust extinction is minimal, since the wavelengths of He I 10830Å and H I $Pa\gamma$ are adjacent.

We utilized the `astropy.dust_extinction` library in Python to accomplish correction (K. D. Gordon 2024). We chose a model called G24_SMC Avg, representing conditions in the SMC (Small Magellanic Cloud) (K. D. Gordon et al. 2024). This model was chosen as it was best representative of the low-metallicity conditions in the early universe. This provided a rough estimate for conditions in LRDs. The G24_SMC Avg model requires A_V , the total visual extinction, to be passed in for a correct model. We calculated this value using Balmer decrements (defined as $H\alpha / H\beta$) and a formula derived from E. J. Nelson et al. (2016).

The color excess was calculated using this formula:

$$E(H\beta - H\alpha) = 2.5 \cdot \log\left(\frac{(H\alpha/H\beta)_{\text{obs}}}{(H\alpha/H\beta)_{\text{int}}}\right)$$

Then, we multiplied the color excess by the reddening curve to find the total visual extinction:

$$A_V = \frac{E(H\beta - H\alpha)}{k(\lambda_{H\beta}) - k(\lambda_{H\alpha})}$$

We passed A_V into the model to obtain a correction factor for each flux value, which we divide the flux by to obtain the corrected values.

We used the Python library `pyneb` in order to generate theoretical predictions of emissivity ratios from density and temperature (V. Luridiana et al. 2015). To model a wide variety of LRD conditions, we chose a range of density logarithmically from 10^5 to 10^{13} cm^{-3} , and a range of temperature linearly from 5000 K and 25000 K . For He I 10830Å and H I $Pa\gamma$, the range of density was adjusted to 10^9 to 10^{15} . The ranges were both generated at resolutions of 100 steps, and were capped at these particular values since `pyneb` would lose precision and throw an error for values beyond this range. We computed the flux of He I 7065Å and 5876Å for each combination of temperature and density. Then, we calculated the ratio of these two values and stored them in a matrix.

Finally, we calculated the difference in flux between each place in the matrix and each observed flux value, recording the minimum difference for each observed flux. This allows us to find the temperature and density prediction closest to the observed flux.

Table 3. This is a temperature/density prediction table of each LRD, utilizing the 5876Å and 7065Å ratio.

LRD	Temperature (K)	Density cm^{-3}
OCEANS 20504 P4	9108	$1 \cdot 10^{13}$
OCEANS 20504 P1	11840	$2.29 \cdot 10^7$
OCEANS 169045 P4	9108	$1 \cdot 10^{13}$
OCEANS 102364 P5	9108	$1 \cdot 10^{13}$

2.6. Abundance Calculation

Using the predictions given by `pyneb`, we are comparing two emission lines at a one-to-one ratio without acknowledging the relative prevalence of different element types. This issue does not apply to the ratio between the He I 5876Å and 7065Å line though, as they both derive from the same element, Helium. For this reason, we will focus on the ratio between the He I 10830Å and H I $Pa\gamma$ lines.

In order to find the ratio between the flux of two emission lines, we must take into consideration two main components: the number of emitting ions and the emission efficiency of each ion. This can be seen in the following equation from E. Aver et al. (2013), which has been adapted for the He I 10830Å and H I 10941Å lines:

$$\frac{F(\text{He I } \lambda 10830)}{F(\text{Pa } \gamma)} \approx \frac{n(\text{He}^+)}{n(\text{H}^+)} \frac{\epsilon(\text{He I } \lambda 10830)}{\epsilon(\text{Pa } \gamma)}$$

Although there are other observational factors that affect the measured fluxes of these lines, they are removed from the equation because these factors affect the He I 10830Å and H I $Pa\gamma$ lines similarly, and cancel once the ratio is calculated.

`pyneb` only gives us the emission portion of the above equation:

$$R_{\text{PyNeb}} = \frac{\epsilon(\text{He I } \lambda 10830)}{\epsilon(\text{Pa } \gamma)}$$

We still must account for the abundance of He^+ and H^+ to properly predict the flux ratio. To do this, we must measure the flux of a different helium line and hydrogen line. We found that the He I 7065 and H α lines were the most consistently simultaneously observed lines in systems with He I 10830Å emission, measuring the narrow component of each for consistency. Rearranging the above flux ratio equation and substituting these lines, we get the following:

$$\frac{n(\text{He}^+)}{n(\text{H}^+)} \approx \frac{F(\text{He I } \lambda 7065)}{F(\text{H } \alpha)} \frac{\epsilon(\text{H } \alpha)}{\epsilon(\text{He I } \lambda 7065)}$$

However, we encounter an issue where in order to calculate the emissivity needed to find the abundance, we need a given temperature and density, but in order to get that temperature and density, we need to calculate

the flux ratios that require the abundance ratio. This creates a circular loop where each component relies on the other.

For simplicity, we assume the abundance ratio here is approximately 0.01. We multiply each element in the Pyneb prediction matrix by this value to get a more reasonable result.

3. RESULTS

All plots featured in this section were created using `matplotlib` (J. D. Hunter 2007).

3.1. Metastable Flux

Analysis of the spectra for each of the 4 LRDs revealed that the $Pa\gamma$ line was prominent in 4 of the 5 observations. While the metastable He I 10941Å line wasn't measured directly from the spectra, it was strong in all 5 observations. It is important to note, however, that these lines were still relatively weak compared to other major features of the spectra.

The GLIMPSE 17775 G395M, JADES 28074 G235M, and JADES 28074 G395M observations displayed a strong Gaussian fit for the $Pa\gamma$ line. The RUBIES 40579 G395M and RUBIES 154183 G395M observations had more ambiguous results, with RUBIES 40579 showing signs of a blending, and RUBIES 154183 showing a detection close to the continuum.




Figure 2. The fit of $Pa\gamma$ in GLIMPSE 17775 g395m. This is an example of a well-fit distribution. JADES 28074 g235m and JADES 28074 g395m yielded similar fits.




Figure 3. The fit of $Pa\gamma$ in RUBIES 40579 g395m. This shows the result of a blended line, which is less favorable.




Figure 4. The fit of $Pa\gamma$ in RUBIES 154183 g395m. Here we see the fit only barely above the continuum.

Despite JADES 28074 g295m and JADES 28074 g395m being different observations, their flux measurements are relatively consistent.

Error between LRDs varied heavily, ranging between ± 2.5 in GLIMPSE 17775 to $^{+0.35}_{-0.34}$ in RUBIES 40579.

Analysis of these LRDs' ratio $\frac{F_{He I 10830 \text{ Å}}}{F_{Pa\gamma}}$ reveal a mean of 3.01, with a standard deviation of ~ 1.42 . We also find that a median ratio of ~ 3.71 is exhibited.

Originally, we set the range for our Pyneb density predictions to be from 10^5 to 10^{13} .




Figure 5. Flux ratio prediction matrix generated by `pyneb`, with density scaled logarithmically. Both the lower density and the higher density regions show similarly low flux ratio values.




Figure 6. Graph showing which density lines the flux ratios are best approximated by. Density lines are chosen based on a logarithmic scale similar to that of the `pyneb` matrix. Notably, the measured flux ratios fall near the lower electron density lines.

This, however, creates similarly low flux ratios for both the extreme high-density regions in the prediction matrix and the lower-density regions. This allows for the same low flux ratio LRDs to be misidentified as being low-density, which contradicts the results seen for our He I 7065Å and He I 5876Å lines (discussed in the next subsection). We can see this lower density misrepresentation in Figure 6, where the LRDs fall along the lower-density lines. To accommodate for this, we change the density range for which we create our prediction matrix, now using 10^9 to 10^{15} to eliminate the lower regions with low flux ratio.




Figure 7. Flux ratio prediction matrix generated by Pyneb, with density scaled logarithmically. Here we see that now only the higher density region has areas of low flux ratio.




Figure 8. Graph showing which density lines the flux ratios are best approximated by. Density lines are chosen based on a logarithmic scale similar to that of the Pyneb matrix. Here, the LRDs now reveal themselves to be closer to the extreme high-density lines.

Using this adjusted density range, we now find that the results become more in line with those seen for the He I 7065Å and 5876Å lines (discussed in the next subsection), where the LRDs exhibit signs of extremely high density.

3.2. Other Flux

The spectra of the LRDs revealed that He I 5876Å and 7065Å were relatively weak compared to the major features of the spectra, such as Balmer breaks. Additionally, only 30% of the observed LRDs exhibited statistically significant observations of both fluxes. However, He I 5876Å was detected more commonly than He I 7065Å.

Despite their small presence within the spectra, most emission lines were well-fit by a Gaussian. However, many lines were ambiguous detections, often located only marginally above the continuum. The presence of poor fits was reflected in the calculated flux values. For instance, the flux error peaked at 33% for OCEANS 102364, with a sigma confidence of only 3.03σ .




Figure 9. The fit of He I 5876Å in OCEANS 169045. This is an example of a well-fit distribution.




Figure 10. The fit of He I 7065Å in OCEANS 102364. This is an example of a poor fit, as evidenced by the peak of the emission line being barely above the surrounding continuum.

Another intriguing observation was the variability between different observations of OCEANS 20504. For instance, the P4¹ observation, exhibited the least significant error and was situated considerably below the other data. However, the other observation of OCEANS 020504, the P1 observation, was closer to the other data points and had a prominent error.

The ratio $\frac{F_{5876\text{\AA}}}{F_{7065\text{\AA}}}$ exhibited a median of around 1.39, with a standard deviation of 0.62. The values ranged from as low as 0.43 to as high as 1.96. All flux ratios except OCEANS 20504's P4 observation far exceeded Pyneb's emissivity predictions. Pyneb's emissivity predictions exhibited a maximum of 1, even at densities of 10^{11}cm^{-3} . Consequently, this indicates the presence of extremely dense areas $> 10^{13}\text{cm}^{-3}$.

Considering temperature predictions, the average predicted temperature was 10521K , close to the hypothesized value of $\sim 10000\text{K}$ (R. P. Naidu et al. 2025). However, temperature predictions and density predictions often varied significantly when modifying the range of temperature. In order to prevent such inconsistencies, we manually limited the range of predicted temperature to a window of 3000K around 10000K , which provided more consistent density predictions and are reflective of the temperatures of LRDs implied by other observations.

3.2.1. Flux Corrections from Dust Extinction

As a consequence of dust correction, all flux values significantly decreased, with the average moving from 1.29 to 0.78. However, one outlier continued to persist: OCEANS 169045 continued to exhibit a flux ratio of 1.65 despite dust correction. Despite this anomaly, the predictions of `pyneb` were overall more accurate, causing the density predictions to be situated around $10^{11} - 10^{12}$ for most LRDs. AdTechnical Context: In software development, throw an error or throw an exception is the correct, professional way to describe triggering a program failure. ditionally, the average temperature predictions reduced slightly to 9010K .




Figure 11. A plot of the matrix generated by Pyneb.

¹ P4 (short for pointing 4) denotes the number of the observation.




Figure 12. A plot of LRDs and their temperature and emissivity ratios compared to different lines of constant density.




Figure 13. A plot of LRDs and their temperature and emissivity ratios compared to different lines of constant density. This time, the emissivity ratios are adjusted for dust correction.

4. DISCUSSION

4.1. Importance of Dust Correction

As mentioned in the Results section, the flux ratios between 7065Å and 5876Å were excessively high without dust correction, with the emission of 7065Å displaying two times the flux as 5876Å in one scenario. However, the abnormal flux values is attributable to the fact that dust extinction amplifies the longer wavelength, 7065Å, stronger than the shorter wavelength, 5876Å. Correcting this dust extinction reduces the flux ratios and significantly increases concordance with theoretical predictions. Interestingly, dust correction does not completely resolve the problem of inaccurate flux, as several outliers continue to persist past Pyneb’s prediction bounds.

Furthermore, within our results for the ratios between the metastable He I 10830Å line and $Pa\gamma$ line, we see that even when dust extinction is no longer a factor affecting the ratios, our flux ratios are still pushing the boundaries of what Pyneb is able to predict. The primordial helium-to-hydrogen abundance is 0.08 (T. Hsyu et al. 2020). The abundance ratio used in this study was set as 0.01, which is a much more conservative value. Even with this conservative estimate though, we see that our LRDs fall near the bottom of our density lines, nearly falling below Pyneb’s predicted ratios. These results, combined with the results of the dust corrected $\frac{F_{5876\text{\AA}}}{F_{7065\text{\AA}}}$ ratios allow us to reasonably conclude that dust correction ultimately plays a minimal role in explaining why these ratios seem to go beyond what Pyneb can produce.

4.2. Limitations of Models

To reiterate, LRDs are hypothesized to contain SMBHs surrounded by a highly dense region of gas. The density estimates reveal the gas exhibits a density of at least 10^{10}cm^{-3} , strongly evidencing the presence of an external dense medium. This medium is likely to be optically thick and have strong collisional effects, due to the high density observed.

Two main drawbacks exist when using Pyneb to measure LRD Helium fluxes. First, Helium is treated as a recombination line within `pyneb`, where collisions are assumed to have minimal effect on emissivity. Second, the high density implies an optically thick region, while Pyneb assumes an optically thin medium. One option to resolve these assumptions, and therefore reconcile agreement with experimental data, is to augment radiative transfer code to analyze collisional effects. The library `CLOUDY` in Python could simulate more representative physics and will be the subject of future work.

4.3. Low Detection Rate

Another unique observation from this data is the low detection rate among the LRDs for this ratio. This is in part due to the rarity of the 7065Å line, which acts as the bottleneck for detection. Increased observational depth could increase the significance of the detections and recover lines in more sources. Additionally,

extending the wavelength range of observation could additionally reveal Helium lines in sources where there was no detection.

5. CONCLUSIONS

The goal of this study was to probe the density and temperature of LRDs using Helium emission spectra. The findings indicate a moderately effective probe, capable of restricting most densities to a range of $\geq 10^{11} - 10^{12} \text{ cm}^{-3}$. However, specific density amounts varied widely among different LRDs, with some vastly exceeding predictions. The application of dust correction was critical to arriving at reasonable results, as values significantly exceeded the theoretical predictions when it was not applied. As for temperatures, most LRDs were well-fit near the consensus of 10000 K , although the mean slightly decreased once dust correction was applied. The high densities indicate the presence of an enshrouded supermassive black hole within the LRD, largely reaffirming the existing theory.

However, many improvements to the data collection techniques exist, such as increased observational depth and expanded wavelength coverage. The comparison of the observed flux ratio to theoretical Pyneb predictions imply that more advanced modeling techniques are needed to reproduce the observed high ratios. One software tool which can accomplish this is the Python library CLOUDY, which acknowledges the optical depth of the measurement alongside collisional effects within the cocooning gas.

6. ACKNOWLEDGEMENTS

We would like to acknowledge Kelcey Davis due to her extensive feedback during the writing process, alongside her assistance in debugging our codebase. This work was completed at a research internship for the Institute for Computing in Research.

REFERENCES

- Aver, E., Olive, K. A., Porter, R., & Skillman, E. D. 2013, *Journal of Cosmology and Astroparticle Physics*, 2013, 017017, doi: [10.1088/1475-7516/2013/11/017](https://doi.org/10.1088/1475-7516/2013/11/017)
- Aver, E., Olive, K. A., & Skillman, E. D. 2015, *Journal of Cosmology and Astroparticle Physics*, 2015, 011011, doi: [10.1088/1475-7516/2015/07/011](https://doi.org/10.1088/1475-7516/2015/07/011)
- Berg, D. A., Sanders, R. L., Shapley, A. E., et al. 2025, *The Astrophysical Journal*, 996, 68, doi: [10.3847/1538-4357/ae18db](https://doi.org/10.3847/1538-4357/ae18db)
- Davis, K., Brooks, M., Simons, R. C., et al. 2026, *OCEANS of Absorption: High-resolution NIRSpec Spectroscopy Reveals Diverse Balmer-line Absorption in Little Red Dots*, <https://arxiv.org/abs/2606.00258>
- de Graaff, A., Rix, H.-W., Naidu, R. P., et al. 2025, *Astronomy & Astrophysics*, 701, A168, doi: [10.1051/0004-6361/202554681](https://doi.org/10.1051/0004-6361/202554681)
- Fernández, V., Amorín, R., Firpo, V., & Morisset, C. 2024, *A&A*, 688, A69, doi: [10.1051/0004-6361/202449224](https://doi.org/10.1051/0004-6361/202449224)
- Finkelstein, S. L., Bagley, M. B., Arrabal Haro, P., et al. 2025, *The Astrophysical Journal Letters*, 983, L4, doi: [10.3847/2041-8213/adbbd3](https://doi.org/10.3847/2041-8213/adbbd3)
- Gordon, K. D. 2024, *Journal of Open Source Software*, 9, 7023, doi: [10.21105/joss.07023](https://doi.org/10.21105/joss.07023)
- Gordon, K. D., Fitzpatrick, E. L., Massa, D., et al. 2024, *ApJ*, 970, 51, doi: [10.3847/1538-4357/ad4be1](https://doi.org/10.3847/1538-4357/ad4be1)
- Harris, C. R., Millman, K. J., van der Walt, S. J., et al. 2020, *Nature*, 585, 357, doi: [10.1038/s41586-020-2649-2](https://doi.org/10.1038/s41586-020-2649-2)
- Hsyu, T., Cooke, R. J., Prochaska, J. X., & Bolte, M. 2020, *The Astrophysical Journal*, 896, 77, doi: [10.3847/1538-4357/ab91af](https://doi.org/10.3847/1538-4357/ab91af)
- Hunter, J. D. 2007, *Computing in Science & Engineering*, 9, 90, doi: [10.1109/MCSE.2007.55](https://doi.org/10.1109/MCSE.2007.55)
- Juodbalis, I., Ji, X., Maiolino, R., et al. 2024, *Monthly Notices of the Royal Astronomical Society*, 535, 853873, doi: [10.1093/mnras/stae2367](https://doi.org/10.1093/mnras/stae2367)
- Kocevski, D. D., Finkelstein, S. L., Barro, G., et al. 2025, *The Astrophysical Journal*, 986, 126, doi: [10.3847/1538-4357/adbc7d](https://doi.org/10.3847/1538-4357/adbc7d)
- Kokorev, V., Chisholm, J., Naidu, R. P., et al. 2025, *The Deepest GLIMPSE of a Dense Gas Cocoon Enshrouding a Little Red Dot*, <https://arxiv.org/abs/2511.07515>
- Labbé, I., van Dokkum, P., Nelson, E., et al. 2023, *Nature*, 616, 266, doi: [10.1038/s41586-023-05786-2](https://doi.org/10.1038/s41586-023-05786-2)

- Luridiana, V., Morisset, C., & Shaw, R. A. 2015, A&A, 573, A42, doi: [10.1051/0004-6361/201323152](https://doi.org/10.1051/0004-6361/201323152)
- Matthee, J., Naidu, R. P., Brammer, G., et al. 2024, The Astrophysical Journal, 963, 129, doi: [10.3847/1538-4357/ad2345](https://doi.org/10.3847/1538-4357/ad2345)
- Naidu, R. P., Matthee, J., Katz, H., et al. 2025, arXiv e-prints, arXiv:2503.16596, doi: [10.48550/arXiv.2503.16596](https://doi.org/10.48550/arXiv.2503.16596)
- Naidu, R. P., Matthee, J., Katz, H., et al. 2025, A ”Black Hole Star” Reveals the Remarkable Gas-Enshrouded Hearts of the Little Red Dots, <https://arxiv.org/abs/2503.16596>
- Nelson, E. J., van Dokkum, P. G., Momcheva, I. G., et al. 2016, ApJL, 817, L9, doi: [10.3847/2041-8205/817/1/L9](https://doi.org/10.3847/2041-8205/817/1/L9)
- pandas development team, T. 2020, pandas-dev/pandas: Pandas, latest Zenodo, doi: [10.5281/zenodo.3509134](https://doi.org/10.5281/zenodo.3509134)
- Rusakov, V., Watson, D., Nikopoulos, G. P., et al. 2026, Nature, 649, 574, doi: [10.1038/s41586-025-09900-4](https://doi.org/10.1038/s41586-025-09900-4)
- Sun, Y. R., & Hu, S.-M. 2020, Natl. Sci. Rev., 7, 1818
- Taylor, A. J., Finkelstein, S. L., Kocevski, D. D., et al. 2025, The Astrophysical Journal, 986, 165, doi: [10.3847/1538-4357/add15b](https://doi.org/10.3847/1538-4357/add15b)
- Wang, B. ., de Graaff, A., Davies, R. L., et al. 2025, The Astrophysical Journal, 984, 121, doi: [10.3847/1538-4357/adclca](https://doi.org/10.3847/1538-4357/adclca)