

Eigenvalues and inverse problems

The sharp spectral-matching constant for normal matrices

Difficulty: extreme **Importance:** interesting to the community**Status:** Open

Rating rationale: Extreme reflects a longstanding sharp-constant problem beyond the false constant-one conjecture; community impact is eigenvalue matching and perturbation theory for normal matrices.

Supporting completion barriers and stationary model — 14 September 2026

Sidney Holden (Center for Computational Biology, Flatiron Institute, Simons Foundation). The [report](#), Sections 2–7, proves odd-polygon completion barriers and solves a stationary two-line variational model. The formal quotient near 1.0373054 is not a finite normal-matrix lower bound. The inherited order-193 rational certificate was rerun by both exact positivity methods. Neither this continuation nor that certificate determines the unrestricted sharp constant; status stays Open.

See the [independent informal Codex AI-agent review](#) for the exact audited scope and checks, and the [submission record](#) for verified affiliation, original materials and reproduction details. No full resolution, external human peer review, formal verification or historical novelty is claimed. No Lean verification was performed.

Reviewed continuation — 13 September 2026

Sidney Holden (Center for Computational Biology, Flatiron Institute, Simons Foundation) supplies an exact rational seven-dimensional normal pair proving $C_{\text{normal}} > 1.03077$; see the [continuation report](#), Section 3, Theorem 3.1. A Hall obstruction supplies the matching lower bound and exact positive-definiteness certifies the full operator-norm upper bound. The report also proves restricted dimension reductions in Sections 6–8.

Status remains **Open**: this certified lower bound and the restricted reductions do not determine the exact universal constant or a matching upper bound. No global extremality or finite-dimensional stabilization is asserted.

The [independent informal Codex AI-agent review](#) records its accepted scope and checks. [Submission, original package, reproduction and verified affiliation](#). This is not external human peer review or formal verification; no Lean verification was performed.

Supporting subclass lemma — 12 September 2026

Sidney Holden (Center for Computational Biology, Flatiron Institute, Simons Foundation) supplies [a proof of the sharp constant-one bound](#) when at least one normal input has at most two distinct eigenvalues, using [Theorem 1](#). This passed an [independent informal Codex AI-agent audit](#). It gives no new global bound or value for C_{normal} ; status remains Open. [Submission and verified affiliation](#). No novelty, external human peer review or formal verification is claimed.

Problem statement

For normal matrices $A, B \in \mathbb{C}^{n \times n}$, list their eigenvalues $\alpha_1, \dots, \alpha_n$ and $\beta_1, \dots, \beta_n$ with algebraic multiplicity. Define the bottleneck matching distance

$$d_{\infty}(A, B) = \min_{\pi \in S_n} \max_{1 \leq j \leq n} |\alpha_j - \beta_{\pi(j)}|,$$

where S_n is the permutation group. Determine the exact value of

$$C_{\text{normal}} = \sup_{n \geq 2} \sup_{\substack{A, B \in \mathbb{C}^{n \times n} \\ A \neq B}} \sup_{\text{normal}} \frac{d_{\infty}(A, B)}{\|A - B\|_2}.$$

Here normal means $AA^* = A^*A$, and $\|\cdot\|_2$ is the spectral norm. The target is a single sharp constant valid in every dimension. Equivalently, find the smallest C for which $d_{\infty}(A, B) \leq C\|A - B\|_2$ for all such pairs, with matching sharpness evidence. The conjectured value $C = 1$ has already been refuted; it is not the question being posed.

Why it matters

This is a sharp conditioning question for the complete spectrum of a normal matrix. Matching counts repeated eigenvalues, so a Hausdorff-distance estimate that ignores multiplicity does not answer it.

References

R. Bhatia, *Perturbation Bounds for Matrix Eigenvalues*, SIAM, 2007, Chapter IX, pp. 154–155; the question is restated as Problem 27 in X. Zhan, *Open problems in matrix theory*, §21, pp. 12–13. P. Šemrl, *Geometrical techniques in linear algebra*, ICTP lecture notes, 2009, p. 10, Theorem 5 and the following open question. J. A. Holbrook, *Spectral variation of normal matrices*, LAA 174 (1992), 131–144.

Earlier status check — 2026-09-08

The cited sources distinguish the unresolved sharp constant from Holbrook’s counterexample to one. They give $1 < C_{\text{normal}} < 3$. A. Parusiński and A. Rainer, *Eigenvalue stability of Hermitian and normal matrices*, v1 (2026), Proposition 3.4, still states the classical universal bound with $1 < C < 3$; it does not determine the sharp constant. Searches for *sharp spectral variation constant normal matrices*, *spectral variation best possible constant*, and 2025/2026 follow-ups found no exact value. The explicit open-question sources are historical; the recent bound is context, not an explicit reaffirmation of openness. This is a bounded literature check.

Audit update — 2026-09-10

The 2026 *spectral perturbation manuscript*, Proposition 3.4, still uses a universal constant strictly between 1 and 3 rather than determining the optimum. Sharp-normal-spectral-matching searches found no exact value. This supports the retained open verdict alongside the historical lower-bound examples cited above; the old constant-one statement is already false and is not the target here.