

IE-28: Rigorous Extensions and Exact Certificates

All-stage local existence, an explicit node family,
and a limitation of the eigenpolynomial construction

Prepared by ChatGPT

12 September 2026

Abstract

For the collocation integration matrix on distinct positive nodes, the target is one positive diagonal D for which $I - D^{-1}A$ is nilpotent. We give a self-contained all-stage existence theorem for nodes sufficiently close to a common positive point. The proof regularizes a determinant at coincident nodes, constructs an algebraic confluent seed by a triangular recursion, and applies the implicit function theorem with an explicitly nonsingular Jacobian. We also characterize the node sets admitting a scalar diagonal: they are scaled zeros of the ordinary Laguerre polynomial. Exact integer interval calculations prove twelve additional point or uniform-neighborhood assertions in dimensions four through seven. A further exact sign certificate shows that, at eleven stages near confluence, no eigenpolynomial with all nonnegative coefficients can produce the required diagonal, although a positive diagonal does exist. The arbitrary-node, arbitrary-stage assertion is *not proved or disproved* here. The earlier complete two- and three-stage arguments are retained.

Status of the requested full solution. No full solution was obtained. In particular, the local all-stage theorem below must not be read as a theorem for every admissible node set. This archive contains additional rigorous results and checked computations, not a claimed resolution of IE-28. No claim of priority or independent peer review is made.

What is established

Quantifiers	Result and evidence
Every node pair or triple	Complete analytic proofs, Sections 2 and the preserved prior writeup.
Every stage count; nodes near a common positive point	Analytic existence proof, Theorem 5.3. The neighborhood depends on the stage count.
Every stage count; scaled Laguerre nodes	Explicit scalar diagonal, Theorem 3.2.
Six node sets and six node neighborhoods, $4 \leq n \leq 7$	Exact integer interval certificates, Section 7.
Eleven stages near confluence	Exclusion of a <i>stronger ansatz</i> , not of IE-28, Theorem 6.2.
Every admissible node set at every stage count	Not established.

Contents

1	The exact target and useful reductions	3
2	Complete small-stage cases	4
2.1	Two stages	4
2.2	Three stages	4
3	An explicit family at every stage count	5
4	Removing the singularity at coincident nodes	6
5	All-stage existence near a common positive point	7
6	An exact obstruction to a stronger ansatz	9
7	Exact certificates away from confluence	10
7.1	What the verifier checks	11
7.2	Certified node neighborhoods	11
8	Continuation, checks, and the remaining global gap	11
8.1	Numerical continuation is explicitly not a proof of global existence	11
8.2	Independent checks included	12
8.3	What would still be needed for a full solution	12
9	Reproduction and archive map	13

1 The exact target and useful reductions

Let $n \geq 2$ and $0 < c_1 < \dots < c_n \leq 1$. Define

$$\ell_j(t) = \prod_{k \neq j} \frac{t - c_k}{c_j - c_k}, \quad A_{ij} = \int_0^{c_i} \ell_j(t) dt. \quad (1)$$

The target is

$$D = \text{diag}(d_1, \dots, d_n), \quad d_i > 0, \quad (I - D^{-1}A)^n = 0. \quad (2)$$

It is the fixed-diagonal stiff-limit problem stated in IE-28 [1] and in the conjecture of van der Houwen and de Swart [2, Section 3.2.1]. The MIN-SR-S treatment [3, Section 2.2.3] does not supply a general existence theorem. Nilpotence of $A - D$, a triangular preconditioner, or a sequence of different diagonal sweep matrices would not solve (2). No ordering of the d_i is required.

Put $C = \text{diag}(c_i)$ and $V_{ij} = c_i^{j-1}$. Interpolating monomials and integrating exactly gives

$$A = CV \text{diag}(1, 1/2, \dots, 1/n) V^{-1}, \quad \det A = \frac{\prod_i c_i}{n!} > 0. \quad (3)$$

All eigenvalues of $D^{-1}A$ are one if and only if all eigenvalues of its inverse $A^{-1}D$ are one. Cayley–Hamilton then gives (2). Equivalently, with $G = D^{-1}$,

$$\det(I - \lambda AG) = (1 - \lambda)^n. \quad (4)$$

Here AG and GA are similar. Coefficient identities, rather than numerical eigenvalues near a defective multiple eigenvalue, are used throughout.

For later computation let

$$w_i = \left(\prod_{j \neq i} (c_i - c_j) \right)^{-1}, \quad S = \text{diag}(c_i/w_i).$$

Direct differentiation of the interpolation formula gives

$$A^{-1} = SMS^{-1}, \quad M_{ij} = \begin{cases} (c_i - c_j)^{-1}, & i \neq j, \\ c_i^{-1} + \sum_{j \neq i} (c_i - c_j)^{-1}, & i = j. \end{cases} \quad (5)$$

Indeed, if $W_{ij} = c_i^j$ and $W'_{ij} = jc_i^{j-1}$, then $A^{-1} = W'W^{-1}$. Equivalently it maps values of a polynomial q of degree at most n , with $q(0) = 0$, to the values of q' . Applying this to the cardinal polynomials $t\ell_j(t)/c_j$ proves (5). Diagonal matrices commute with S , so $A^{-1}D$ is similar to MD .

Two immediate necessary identities are

$$\prod_i d_i = \frac{\prod_i c_i}{n!}, \quad \text{tr}(MD) = \sum_i \frac{d_i}{c_i} + \sum_{i < j} \frac{d_i - d_j}{c_i - c_j} = n. \quad (6)$$

They are not sufficient in general.

Lemma 1.1 (Eigenpolynomial). *If p is a nonzero real polynomial of degree at most n , $p(0) = 0$, and $p'(c_i) \neq 0$, then $d_i = p(c_i)/p'(c_i)$ makes one an eigenvalue of $A^{-1}D$, provided the vector of sampled values of p is nonzero.*

Proof. The polynomial-value identity is $A(p'(c_i))_i = (p(c_i))_i$. Thus $D^{-1}A$ fixes the nonzero vector $(p'(c_i))_i$. The stated sampled vector is nonzero whenever all $p(c_i) > 0$, as in the applications below. $\square$

Scaling causes no loss: $A(sc) = sA(c)$ for $s > 0$, so $D \mapsto sD$ preserves the target when the nodes are scaled by s .

2 Complete small-stage cases

This section summarizes the complete constructions from the preserved `prior_work/writeup.pdf`; their short proofs are included here.

2.1 Two stages

Let $c_1 = a < c_2 = b$ and $r = \sqrt{2ab}$. Set

$$d_i = \frac{c_i(c_i + r)}{2c_i + r}, \quad i = 1, 2. \quad (7)$$

The polynomial $p(t) = t(t + r)$ gives eigenvalue one by Lemma 1.1. Moreover

$$\frac{2d_1d_2}{ab} = 1 \iff 2(a + r)(b + r) = (2a + r)(2b + r) \iff r^2 = 2ab.$$

Thus $\det(A^{-1}D) = 1$, so its other eigenvalue is also one. Positivity is immediate, and Cayley–Hamilton proves the target.

2.2 Three stages

For $r > 0$ and $0 \leq \theta \leq 1$, use

$$p(t) = t(t + r)(\theta t + r), \quad d_i(r, \theta) = \left(\frac{1}{c_i} + \frac{1}{c_i + r} + \frac{\theta}{\theta c_i + r} \right)^{-1}. \quad (8)$$

There is always an eigenvalue one. Write

$$H_i(r, \theta) = 1 + \frac{c_i}{c_i + r} + \frac{\theta c_i}{\theta c_i + r}, \quad G(r, \theta) = \det(MD) = 6 \prod_i H_i(r, \theta)^{-1}.$$

For each fixed θ , G is strictly increasing in r and tends to 6 as $r \rightarrow \infty$. Its limit as $r \downarrow 0$ is $6/27$ for $\theta > 0$, and $6/8$ for $\theta = 0$. Hence there is a unique $r(\theta) > 0$ with $G = 1$.

This root is continuous on $[0, 1]$. To justify continuity at the corner $(r, \theta) = (0, 0)$, take any r small enough that $6 \prod_i (1 + c_i/(c_i + r))^{-1} < 1$; then $G(r, \theta) < 1$ for every θ . There is also a uniform upper bracket: at $r = 4c_3$ every $H_i \leq 7/5$, so $G \geq 6/(7/5)^3 > 1$. Consequently the roots lie in a common compact subinterval of $(0, 4c_3)$; strict monotonicity and continuity give the claim.

Set $T(\theta) = \text{tr}(MD(r(\theta), \theta)) - 3$ and let $\sigma_1 = c_1 + c_2 + c_3$, $\sigma_2 = c_1c_2 + c_1c_3 + c_2c_3$, $\sigma_3 = c_1c_2c_3$. At $\theta = 0$, substitution in (6) and clearing denominators yields

$$T(0) = \frac{r(3r^2 + 3\sigma_1r + 2\sigma_2)}{\prod_i (2c_i + r)} > 0. \quad (9)$$

At $\theta = 1$, write $P_3 = \prod_i (3c_i + r)$ and

$$F_3 = 6 \prod_i (c_i + r) - P_3 = 5r^3 + 3\sigma_1 r^2 - 3\sigma_2 r - 21\sigma_3.$$

The determinant constraint is $F_3 = 0$, and direct simplification gives

$$P_3 T(1) = F_3 - 2(r^3 + 3\sigma_3) < 0. \quad (10)$$

Both identities are verified as formal rational identities by the supplied symbolic test. The intermediate value theorem supplies θ_* with $T = 0$. At this point $A^{-1}D$ has an eigenvalue one, trace three, and determinant one. Its other two eigenvalues have sum two and product one, hence are both one. This proves (2) for every node triple.

Finally the denominator in (8) is strictly between $1/c_i$ and $3/c_i$, and it strictly decreases as c_i increases. Therefore

$$c_i/3 < d_i < c_i, \quad d_1 < d_2 < d_3.$$

The use of trace, determinant, and one known eigenvalue in this last spectral step is special to three stages; it does not settle four or more stages.

3 An explicit family at every stage count

Lemma 3.1 (Characteristic polynomial from the node polynomial). *Let $\pi(t) = \prod_{i=1}^n (t - c_i)$. Then*

$$\det(zI - A) = \frac{1}{n!} \sum_{j=0}^n \pi^{(j)}(0) z^j = \sum_{k=0}^n (-1)^k \frac{(n-k)!}{n!} e_k(c) z^{n-k}, \quad (11)$$

where e_k is the k th elementary symmetric polynomial.

Proof. Identify sampled vectors with polynomials of degree at most $n-1$. Integration followed by reduction modulo π represents A . In the basis $1, t, t^2/2!, \dots, t^{n-1}/(n-1)!$, integration maps each basis vector other than the last to the next one. For the last vector it gives $t^n/n!$, which, modulo π , is

$$-\sum_{j=0}^{n-1} \frac{\pi^{(j)}(0)}{n!} \frac{t^j}{j!}.$$

The representing matrix is a companion matrix with ones on the first subdiagonal and these coefficients in its last column. Its characteristic polynomial is the first expression in (11). The second follows by expanding π . $\square$

Define the ordinary Laguerre polynomial by

$$L_n(x) = \sum_{j=0}^n (-1)^j \binom{n}{j} \frac{x^j}{j!}.$$

Theorem 3.2 (Scalar diagonals are exactly the scaled Laguerre case). *Let $x_1 < \dots < x_n$ be the zeros of L_n . They are positive and simple. For any $\delta > 0$ with $\delta x_n \leq 1$, the nodes $c_i = \delta x_i$ satisfy (2) with*

$$D = \delta I. \quad (12)$$

Conversely, if a node set admits a scalar positive diagonal $D = \delta I$ satisfying (2), its node polynomial is $\pi(t) = (-\delta)^n n! L_n(t/\delta)$.

Proof. For completeness, the Rodrigues identity

$$e^{-x}L_n(x) = \frac{1}{n!} \frac{d^n}{dx^n}(e^{-x}x^n)$$

follows by the product rule. Integrating by parts n times shows that L_n is orthogonal to every polynomial of degree less than n against e^{-x} on $(0, \infty)$. The boundary terms vanish. If L_n had fewer than n sign-changing zeros there, multiply the factors corresponding to those zeros to obtain a polynomial of degree less than n whose product with L_n has one sign and is not zero almost everywhere. Its integral cannot vanish, contradicting orthogonality. Thus all n zeros are positive and simple.

For the scaled zeros, the monic node polynomial is $\pi(t) = (-\delta)^n n! L_n(t/\delta)$, so

$$\frac{\pi^{(j)}(0)}{n!} = (-1)^{n+j} \delta^{n-j} \binom{n}{j}.$$

Lemma 3.1 gives $\det(zI - A) = (z - \delta)^n$. Cayley–Hamilton proves (12). Conversely the scalar target forces that characteristic polynomial, so (11) determines every Taylor coefficient of π and gives the stated Laguerre expression. $\square$

This is an exact all-stage class, not a reduction of arbitrary nodes to this class: scaling alone cannot turn an arbitrary node set into Laguerre zeros.

4 Removing the singularity at coincident nodes

For a polynomial g of degree at most $n - 1$ define, initially at distinct nodes,

$$\Phi(\lambda; c, g) = \frac{\det[jc_i^{j-1} - \lambda g(c_i)c_i^j]_{i,j=1}^n}{n! \Delta(c)}, \quad \Delta(c) = \prod_{i < j} (c_j - c_i). \quad (13)$$

The numerator is alternating in the node variables. In the polynomial ring whose other indeterminates are λ and the coefficients of g , it is divisible by each $c_j - c_i$, hence by their product. Consequently Φ is a polynomial and extends across all node collisions.

Lemma 4.1. *At distinct positive nodes, if $G = \text{diag}(g(c_i))$, then*

$$\Phi(\lambda; c, g) = \det(I - \lambda AG). \quad (14)$$

Thus $g(c_i) > 0$ and $\Phi = (1 - \lambda)^n$ give a solution $D = \text{diag}(1/g(c_i))$ of IE-28.

Proof. Use W, W' from Section 1. The numerator is $\det(W' - \lambda GW)$. Since $A^{-1} = W'W^{-1}$ and $\det W' = n! \Delta(c)$, its ratio to $\det W'$ is $\det(A^{-1} - \lambda G)/\det A^{-1}$, which is (14). $\square$

A division-free formula is useful both as a check and as a numerical implementation. Let $h_m(x_1, \dots, x_r)$ denote the complete homogeneous symmetric polynomial, with $h_0 = 1$ and $h_m = 0$ for $m < 0$. The elementary monomial divided-difference identity is

$$[x_1, \dots, x_{k+1}]t^m = h_{m-k}(x_1, \dots, x_{k+1}).$$

For $g(t) = \sum_{r=0}^{n-1} g_r t^r$, define, for $0 \leq k \leq n-1$, $1 \leq j \leq n$,

$$\begin{aligned} (B_0)_{k,j-1} &= j h_{j-1-k}(c_1, \dots, c_{k+1}), \\ (B_1)_{k,j-1} &= \sum_{r=0}^{n-1} g_r h_{r+j-k}(c_1, \dots, c_{k+1}). \end{aligned} \quad (15)$$

Changing evaluation rows into successive divided-difference rows divides their determinant by $\Delta(c)$. Hence

$$\Phi(\lambda; c, g) = \det(B_0 - \lambda B_1)/n!, \quad \det B_0 = n!. \quad (16)$$

Unlike the defining quotient, this formula can be evaluated at coincident nodes without subtracting nearly equal nodes or dividing by node gaps.

5 All-stage existence near a common positive point

Parameterize g by $a = (a_1, \dots, a_n)$ through

$$H_a(z) = \sum_{j=1}^n a_j z^j, \quad g_a(t) = - \sum_{j=1}^n j a_j (1-t)^{j-1}. \quad (17)$$

In particular $g_a(1) = -a_1$ and $\int_1^t g_a(s) ds = H_a(1-t)$.

Lemma 5.1 (Confluent determinant). *At $c_1 = \dots = c_n = 1$, the polynomial extension satisfies*

$$\begin{aligned} \Phi(\lambda; \mathbf{1}, g_a) &= [z^n] \frac{\exp(\lambda H_a(z))}{1-z} = 1 + \sum_{k=1}^n f_k(a) \lambda^k, \\ f_k(a) &= \frac{1}{k!} [z^n] \frac{H_a(z)^k}{1-z}. \end{aligned} \quad (18)$$

Only finite Taylor coefficients are used in this expression.

Proof. The confluent evaluation determinant in (13) has rows consisting of the Taylor coefficients of $q' - \lambda g_a q$ of orders $0, \dots, n-1$ at $t = 1$, with q ranging over $t, t^2, \dots, t^n$. Write $s = t - 1$ and $q(t) = \sum_{m=0}^n y_m s^m$. The constraint $q(0) = 0$ is

$$\sum_{m=0}^n (-1)^m y_m = 0. \quad (19)$$

For an arbitrary y_0 , the equations that the indicated residual coefficients vanish form a triangular recursion for $y_1, \dots, y_n$, with pivots $1, 2, \dots, n$. Its solution is

$$y_m = y_0 [s^m] \exp(\lambda H_a(-s)), \quad 0 \leq m \leq n.$$

Eliminating $y_1, \dots, y_n$ from these equations and the boundary equation leaves y_0 multiplied by

$$\sum_{m=0}^n [z^m] \exp(\lambda H_a(z)).$$

To see that this is an identity of determinants, not just an equivalence of zero sets, order the augmented equations with (19) first and the residual equations next. Their determinant is $n!$

times the displayed factor by the triangular elimination. Eliminating y_0 using its unit coefficient in (19) gives precisely the restricted residual matrix. The change from coefficients in $t, t^2, \dots, t^n$ to $y_1, \dots, y_n$ has determinant one (its entries are $\binom{j}{k}$). Division by $n!$ therefore gives the normalized determinant in (13). Summing the Taylor coefficients is coefficient extraction against $(1 - z)^{-1}$, proving (18). $\square$

Lemma 5.2 (Triangular algebraic seed). *There is a unique $a^* \in \mathbb{R}^n$ with $a_1^* < 0$ such that $f_k(a^*) = (-1)^k \binom{n}{k}$ for $k = 1, \dots, n$. It is given by*

$$\begin{aligned} r_n &= (n!)^{1/n}, & a_1^* &= -r_n, \\ a_j^* &= \frac{(k-1)!}{(a_1^*)^{k-1}} \left((-1)^k \binom{n}{k} - f_k(a_1^*, \dots, a_{j-1}^*, 0, \dots, 0) \right), & k &= n - j + 1, \quad 2 \leq j \leq n. \end{aligned} \quad (20)$$

The Jacobian of $a \mapsto (f_n, f_{n-1}, \dots, f_1)$ at this seed is nonsingular, with determinant

$$\frac{(a_1^*)^{n(n-1)/2}}{\prod_{m=0}^{n-1} m!} \neq 0. \quad (21)$$

Proof. Since H_a has no constant term, $f_n = a_1^n/n!$, and its required value fixes the unique negative $a_1 = -r_n$. For $k = n - j + 1$, the terms of H_a^k of degree at most n involve no coefficient beyond a_j . A term containing a_j can have degree at most n only when it contains $a_j z^j$ exactly once and $a_1 z$ in all other factors. For $j \geq 2$ its coefficient in f_k is therefore $a_1^{k-1}/(k-1)!$, nonzero when $a_1 \neq 0$. This proves the recursion. In the reversed row order, the Jacobian is triangular, and its diagonal is

$$\frac{a_1^{n-1}}{(n-1)!}, \frac{a_1^{n-2}}{(n-2)!}, \dots, a_1, 1.$$

Their product proves (21). The first diagonal entry comes by differentiating f_n and is consistent with this expression. $\square$

Theorem 5.3 (All-stage clustered-node existence). *For every integer $n \geq 2$, there exists $\varepsilon_n > 0$ such that all distinct nodes with $1 - \varepsilon_n < c_1 < \dots < c_n \leq 1$ admit a positive diagonal satisfying (2). There is a real-analytic branch, in polynomial coordinates, with*

$$d_i \longrightarrow (n!)^{-1/n} \quad (c \longrightarrow \mathbf{1}). \quad (22)$$

More generally, nodes sufficiently close to any common $c_ > 0$ admit such a diagonal, with limiting entry $c_*(n!)^{-1/n}$.*

Proof. Let $\mathcal{F}(a, c)$ be the vector of coefficients of $\lambda^n, \dots, \lambda$ in $\Phi(\lambda; c, g_a) - (1 - \lambda)^n$. By Section 4, this is a polynomial map in (a, c) , including at $c = \mathbf{1}$. Lemmas 5.1 and 5.2 give $\mathcal{F}(a^*, \mathbf{1}) = 0$ and an invertible derivative with respect to a . The real-analytic implicit function theorem yields a solution $a(c)$ for all c in some neighborhood of $\mathbf{1}$.

At the center, $g_{a^*}(1) = r_n > 0$. By continuity, after shrinking that neighborhood, $g_{a(c)}(c_i) > 0$ for every i . Set $d_i = 1/g_{a(c)}(c_i)$. Lemma 4.1 proves (2) at all distinct positive nodes in the neighborhood. It also gives the limit (22). Scaling from Section 1 proves the assertion at a common point c_* . $\square$

Remark 5.4 (What locality means). The theorem gives a positive neighborhood for each fixed n . It does not bound ε_n uniformly in n , does not claim that $\varepsilon_n = 1$, and does not prove that this branch extends through the entire ordered-node domain. The coincident-node matrix A itself is not used: the implicit function theorem is applied to its polynomial determinant regularization.

6 An exact obstruction to a stronger ansatz

A tempting route is to choose

$$p(t) = t \prod_{j=1}^{n-1} (t + \alpha_j), \quad \alpha_j > 0, \quad d_i = p(c_i)/p'(c_i). \quad (23)$$

It ensures positivity and $c_i/n < d_i < c_i$, but those extra restrictions were not required by IE-28. The following result prevents promoting this route to an unrestricted all-stage proof.

Define the confluent normalized polynomial

$$p_*(t) = \sum_{m=0}^n \left([z^m] e^{H_{a^*}(z)} \right) (1-t)^m. \quad (24)$$

Then $p_*(1) = 1$, and $p_*(0) = \Phi(1; \mathbf{1}, g_{a^*}) = 0$. It is the unique polynomial of degree at most n with value one at 1 whose residual $p'_* - g_{a^*} p_*$ has zero Taylor coefficients of orders $0, \dots, n-1$ there.

Proposition 6.1 (Exact integer sign certificate). *For $n = 11$, the coefficient $\beta_{11} = [t^{11}]p_*(t)$ satisfies*

$$-1.664575 \cdot 10^{-10} < \beta_{11} < -1.664574 \cdot 10^{-10} < 0. \quad (25)$$

These inequalities are verified with integer arithmetic, not inferred from floating-point roots.

Certificate and its mathematical justification. The file `cluster/certify_seed.py` brackets $r = (11!)^{1/11}$ between consecutive multiples of 10^{-100} by comparing integer eleventh powers. The strict bracket is checked exactly. It propagates this bracket through (20) and the truncated exponential in (24), using outward-rounded dyadic intervals at 512 bits. All sums, products, reciprocals, and factorial divisions enclose the exact real operations. The denominators in (20) are nonzero powers of the strictly negative a_1 . Consequently the resulting interval encloses the exact algebraic coefficient $(-1)^{11} [z^{11}] e^{H_{a^*}(z)}$. Its two endpoints are checked, by integer cross multiplication, to lie strictly between the rational endpoints in (25). The full rational endpoints and the root bracket are saved in `cluster/seed11_certificate.json`. This finite verification uses Python's standard library only. It is a reproducible computer-assisted sign proof; the decimal display printed by the program is not used by the proof. $\square$

Theorem 6.2 (No nonnegative-coefficient eigenpolynomial near confluence). *For $n = 11$ there exists $\eta > 0$ such that, for any distinct nodes $1 - \eta < c_1 < \dots < c_{11} \leq 1$, no polynomial p with*

$$p(0) = 0, \quad p \neq 0, \quad \deg p \leq 11, \quad \text{all coefficients of } p \text{ nonnegative}$$

can yield a solution of (2) by $d_i = p(c_i)/p'(c_i)$. In particular, (23) cannot solve these node sets. Nevertheless, for a sufficiently small such η , positive diagonals do exist by Theorem 5.3.

Proof. Suppose to the contrary that there are node vectors $c^{(m)} \rightarrow \mathbf{1}$ and such polynomials p_m producing solutions. Normalize $p_m(1) = 1$, which does not change p_m/p'_m . Their coefficients form a compact simplex. Along a subsequence they converge coefficientwise to a polynomial p with nonnegative coefficients, $p(0) = 0$, and $p(1) = 1$.

On a fixed real neighborhood of 1, the p_m are uniformly bounded away from zero; for example $p_m(t) \geq 2^{-11}$ for $1/2 \leq t \leq 1$. Their derivatives of all fixed orders converge, so $g_m = p'_m/p_m$ and their

fixed-order derivatives converge uniformly on a smaller neighborhood. Let $\tilde{g}_m$ be the polynomial of degree at most ten interpolating g_m at $c^{(m)}$. Divided differences converge to derivatives at a repeated node; this follows directly from the integral form of a divided difference and the uniform derivative convergence. Thus $\tilde{g}_m$ converges, coefficientwise, to the degree-ten Taylor polynomial of $g = p'/p$ at 1. Call this limit g_0 .

The solution assumption and Lemma 4.1 give $\Phi(\lambda; c^{(m)}, \tilde{g}_m) = (1 - \lambda)^{11}$. Polynomial continuity gives the same identity at $(1, g_0)$. Moreover $g_0(1) = p'(1) > 0$; in fact $1 \leq p'(1) \leq 11$ from the normalized nonnegative coefficients. By the uniqueness in Lemma 5.2, $g_0 = g_{a^*}$. The residual $p' - g_0 p$ therefore vanishes to order at least 11 at 1. Its Taylor recursion, together with $p(1) = 1$, forces $p = p_*$ from (24). This contradicts the negative leading coefficient in Proposition 6.1. $\square$

The theorem does not say that a numerical solver failing to find negative roots has disproved IE-28. It proves failure of a specified auxiliary construction and separates that failure from the positive-diagonal question.

7 Exact certificates away from confluence

The analytical local theorem does not identify its neighborhood size. Independently, the archive supplies exact certificates for specified node sets and explicit node boxes. No assertion in this section is based only on a small residual.

For $M(c)$ from (5), write

$$F_k(d, c) = \sum_{\substack{S \subseteq \{1, \dots, n\} \\ |S|=k}} \det M(c)[S, S] \prod_{i \in S} d_i - \binom{n}{k}, \quad 1 \leq k \leq n. \quad (26)$$

Then $F = 0$ is exactly the characteristic-coefficient system for MD having all eigenvalues one. The Jacobian used by the verifier is the derivative of this displayed polynomial with respect to d .

Theorem 7.1 (Uniform contraction certificate). *Let $\mathcal{C}$ be a closed node box consisting only of separated positive nodes with last node at most one, and let $X = \{d : \|d - d^0\|_\infty \leq r\}$ have strictly positive coordinates. Let R be a fixed real $n \times n$ matrix. Suppose rigorous bounds satisfy*

$$\begin{aligned} \|RF(d^0, c)\|_\infty &\leq \xi & (c \in \mathcal{C}), \\ \|I - RF_d(d, c)\|_\infty &\leq q < 1 & (d \in X, c \in \mathcal{C}), \\ \xi + qr &< r. \end{aligned} \quad (27)$$

For every $c \in \mathcal{C}$ there exists exactly one zero of $F(\cdot, c)$ in X . It gives a positive diagonal solving (2).

Proof. For each fixed c , define $T_c(d) = d - RF(d, c)$. The derivative bound and the mean-value integral on the convex box X make T_c a contraction with constant at most q . The first and third bounds show $T_c(X) \subset X$. Banach's fixed-point theorem gives a unique fixed point in X . For any one $d \in X$, $\|I - RF_d(d, c)\| < 1$ implies that $RF_d(d, c)$ is invertible, so the square matrix R is invertible. Fixed points of T_c are therefore exactly zeros of $F(\cdot, c)$. Positivity is part of the hypothesis on X , and the coefficient characterization proves the target. Uniqueness is only asserted *inside this diagonal box*. $\square$

7.1 What the verifier checks

All centers, radii, and preconditioner entries in the JSON certificates are exact rational numbers (decimal strings denote exact terminating rationals). The verifier uses intervals $[L/2^b, U/2^b]$ with integer endpoints. Addition, multiplication, and reciprocals are rounded outward using integer floor and ceiling division. Principal determinants are enclosed by division-free subset dynamic programming, not by a floating-point factorization. At $b = 320$, it obtains integer upper bounds E, Q for $2^b \xi, 2^b q$. For $r = r_N/r_D$, its final self-mapping test is precisely

$$Er_D + Qr_N < 2^b r_N,$$

and its contraction test is $Q < 2^b$. Both are integer comparisons. The positivity of the diagonal box, separation of every node interval, and the upper node bound are also checked. A passing result is an application of Theorem 7.1; the approximate candidate generator is outside the trusted verification path.

7.2 Certified node neighborhoods

In the following table the last node is fixed at one; every other node may vary independently by $\pm\delta$ about its listed center. The diagonal box is centered at the rational vector in the certificate file and has radius r . The displayed q, ξ bounds have been rounded *upward* for readability; the actual acceptance tests use their exact rational versions.

n	Node center	δ	r	q upper	ξ upper
4	(.1, .3, .6, 1)	10^{-5}	10^{-3}	.222	.000600
4	$c_i = i/4$	10^{-4}	10^{-2}	.475	.00436
5	$c_i = i/5$	10^{-6}	10^{-3}	.139	.000108
6	$c_i = i/6$	10^{-6}	10^{-3}	.346	.000252
7	$c_i = i/7$	10^{-7}	10^{-3}	.771	.0000622
5	(.01, .02, .03, .04, 1)	10^{-15}	10^{-8}	.228	$9.0 \cdot 10^{-10}$

All six boxes pass. There is also a separate, much tighter, point certificate at each of these centers, using $r = 10^{-30}$. Thus twelve certificates pass in total. The separated five-node example needs a small box under the straightforward interval bounds used here; no claim of a maximal neighborhood is made. Scaling the nodes and diagonal extends each normalized assertion to its corresponding positive scaling family.

8 Continuation, checks, and the remaining global gap

8.1 Numerical continuation is explicitly not a proof of global existence

The regularization supplies a natural numerical path

$$c_i(s) = 1 - s(1 - c_i), \quad 0 \leq s \leq 1,$$

starting at the exact algebraic seed. The supplied program performs damped Newton corrections while keeping all sampled $g_a(c_i(s))$ positive. For improved conditioning it uses $z_i = 1 - c_i$ and the

polynomial basis $q_j(t) = (1 - t)^j - 1$, which automatically satisfies the boundary condition. In this basis the divided-difference derivative matrix is

$$T_{k,j-1} = -j h_{j-1-k}(z_1, \dots, z_{k+1}),$$

and the derivative of the $g_a q_j$ matrix with respect to a_{r+1} is

$$-(r+1)(h_{r+j-k}(z_1, \dots, z_{k+1}) - h_{r-k}(z_1, \dots, z_{k+1})).$$

The normalized pencil is $I - \lambda K(a)$ after left multiplication by T^{-1} . Its coefficients and their derivatives are evaluated through finite matrix recurrences, not by differentiating numerical eigenvalues.

The saved equally spaced examples for $n = 8, 11, 15$ reach $s = 1$ with positive sampled g , using 60, 80, and 90 decimal working digits respectively. Their final coefficient residuals are approximately $1.6 \cdot 10^{-38}$, $4.3 \cdot 10^{-49}$, and $7.7 \cdot 10^{-49}$. These are high-precision *numerical examples only*; they are not among the twelve interval certificates. A flag named `pass` in those numerical JSON files means that the numerical stopping tests passed, not that a theorem was proved.

8.2 Independent checks included

The exact regression suite performs 1,500 rational interval-arithmetic trials, 50 rational determinant comparisons, formal seed-Jacobian checks through $n = 6$, confluent determinant checks through $n = 5$, exact collocation characteristic checks through $n = 7$, and symbolic Laguerre identities through $n = 15$. It reruns all twelve certificates and confirms rejection of three intentionally corrupted inputs. The previous two- and three-stage symbolic identities are checked separately. These tests help detect implementation errors; finite tests are not substitutes for the analytical all-stage proofs.

8.3 What would still be needed for a full solution

The missing implication is global:

$$\forall n \geq 4 \forall 0 < c_1 < \dots < c_n \leq 1, \quad \exists d \in (0, \infty)^n : F(d, c) = 0.$$

For the particular continuation route above, one would need a theorem that prevents loss of the required solution before $s = 1$: for example, a valid continuation or degree argument controlling finite escape, approach to the boundary of positivity, and singularities of the coefficient Jacobian. Those conditions have not been established here. A nonsingular Jacobian at $s = 0$, or successful Newton tracking for finitely many target node sets, does not establish them.

Several shortcuts therefore remain invalid. The trace and determinant constraints are insufficient beyond the small-stage argument; generic complex multiplicative inverse-eigenvalue theorems [5] do not guarantee positive real diagonals; and the nonnegative-coefficient construction is ruled out near an eleven-stage confluence by Theorem 6.2. Earlier low-order algebraic and higher-order numerical investigations [4] are valuable context, not an all-node proof. The bounded literature search conducted for this attempt did not locate a full theorem or a counterexample settling the stated target.

9 Reproduction and archive map

The main proofs are in this document and its editable LaTeX source. `README.md` gives commands for each check. The key commands, run from the extracted archive root, are:

```
python certification/verify.py \
    certification/point_certificates.json \
    certification/neighborhood_certificates.json \
    --report results/verification_report.json
python cluster/certify_seed.py \
    --report cluster/seed11_certificate.json
python tests/test_exact.py
python prior_work/scripts/check_symbolic.py
```

The first two use the Python standard library only. The symbolic regression requires SymPy. Numerical continuation and candidate generation require mpmath, and the preserved previous numerical examples also use NumPy. Dependencies and the tested environment are recorded in the archive. `prior_work/` preserves the preceding partial writeup and its sources; its status statements describe that earlier version, not the additional results proved in the present document.

References

- [1] *OpenProblemsInNLA*, IE-28, “Positive diagonal preconditioning with a nilpotent stiff limit.” Public problem statement, accessed 12 September 2026; the entry records a bounded status check dated 11 September 2026. <https://github.com/ajt60gaibb/OpenProblemsInNLA/blob/main/linear-systems-and-elimination/IE-28/README.md>.
- [2] P. J. van der Houwen and J. J. B. de Swart, “Triangularly implicit iteration methods for ODE-IVP solvers,” *SIAM Journal on Scientific Computing* 18(1) (1997), 41–55. DOI: 10.1137/S1064827595287456. Open copy: <https://ir.cwi.nl/pub/2191/2191D.pdf>.
- [3] G. Čaklović, T. Lunet, S. Götschel, and D. Ruprecht, “Improving efficiency of parallel across the method spectral deferred corrections,” *SIAM Journal on Scientific Computing* 47(1) (2025), A430–A453. DOI: 10.1137/24M1649800. Open text: <https://epubs.siam.org/doi/10.1137/24M1649800>.
- [4] W. M. Lioen, “On the diagonal approximation of full matrices,” CWI report NM-R9518 (1995); journal version in *Journal of Computational and Applied Mathematics* 75(1) (1996), 35–42. Report record: <https://ir.cwi.nl/pub/4925>; PDF: <https://ir.cwi.nl/pub/4925/04925D.pdf>.
- [5] J. Rosenthal and X. Wang, “The multiplicative inverse eigenvalue problem over an algebraically closed field,” *SIAM Journal on Matrix Analysis and Applications* 23(2) (2001), 517–523. Preprint: <https://arxiv.org/abs/math/0009163>. Its algebraically closed-field setting is essential and is not a positive-real existence theorem for IE-28.