

The association of smoking status with SARS-CoV-2 infection, hospitalisation and mortality from COVID-19: A living rapid evidence review with Bayesian meta-analyses (version 7)

RUNNING HEAD: Living rapid review of smoking and COVID-19

David Simons¹, Lion Shahab², Jamie Brown² & Olga Perski²

¹ Centre for Emerging, Endemic and Exotic Diseases, Royal Veterinary College

² Department of Behavioural Science and Health, University College London, 1-19 Torrington Place, London WC1E 7HB

Word count abstract: 340

Word count manuscript: 3871

Key words: COVID-19; hospitalisation; infection; mortality; SARS-CoV-2; smoking; tobacco; e-cigarettes; nicotine replacement therapy; living review

Declaration of conflicts of interest

DS and OP have no conflicts of interest to declare. LS has received a research grant and honoraria for a talk and travel expenses from manufacturers of smoking cessation medications (Pfizer and Johnson & Johnson). JB has received unrestricted research funding to study smoking cessation from companies who manufacture smoking cessation medications. All authors declare no financial links with tobacco companies or e-cigarette manufacturers or their representatives.

Funding statement

DS is supported by a PhD studentship from the UK Biotechnology and Biological Sciences Research Council [BB/M009513/1]. OP receives salary support from Cancer Research UK (C1417/A22962). JB, LS, & OP are members of SPECTRUM, a UK Prevention Research Partnership Consortium (MR/S037519/1). UKPRP is an initiative funded by the UK Research and Innovation Councils, the

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/add.152776

Department of Health and Social Care (England) and the UK devolved administrations, and leading health research charities.

Abstract

Aims: To estimate the association of smoking status with rates of i) infection, ii) hospitalisation, iii) disease severity, and iv) mortality from SARS-CoV-2/COVID-19 disease.

Design: Living rapid review of observational and experimental studies with random-effects hierarchical Bayesian meta-analyses. Published articles and pre-prints were identified via MEDLINE and medRxiv.

Setting: Community or hospital. No restrictions on location.

Participants: Adults who received a SARS-CoV-2 test or a COVID-19 diagnosis.

Measurements: Outcomes were SARS-CoV-2 infection, hospitalisation, disease severity and mortality stratified by smoking status. Study quality was assessed (i.e. 'good', 'fair' and 'poor').

Findings: Version 7 (searches up to 25 August 2020) included 233 studies with 32 'good' and 'fair' quality studies included in meta-analyses. Fifty-seven studies (24.5%) reported current, former and never smoking status. Recorded smoking prevalence among people with COVID-19 was generally lower than national prevalence. Current compared with never smokers were at reduced risk of SARS-CoV-2 infection ($RR=0.74$, 95% Credible Interval (CrI) = 0.58-0.93, $\tau = 0.41$). Data for former smokers were inconclusive ($RR=1.05$, 95% CrI = 0.95-1.17, $\tau = 0.17$) but favoured there being no important association (21% probability of $RR \geq 1.1$). Former compared with never smokers were at somewhat increased risk of hospitalisation ($RR=1.20$, CrI = 1.03-1.44, $\tau = 0.17$), greater disease severity ($RR=1.52$, CrI = 1.13-2.07, $\tau = 0.29$), and mortality ($RR=1.39$, 95% CrI = 1.09-1.87, $\tau = 0.27$). Data for current smokers were inconclusive ($RR=1.06$, CrI = 0.82-1.35, $\tau = 0.27$; $RR=1.25$, CrI = 0.85-1.93, $\tau = 0.34$; $RR=1.22$, 95% CrI = 0.78-1.94, $\tau = 0.49$ respectively) but favoured there being no important associations with hospitalisation and mortality (35% and 70% probability of $RR \geq 1.1$, respectively) and a small but important association with disease severity (79% probability of $RR \geq 1.1$).

Conclusions: Compared with never smokers, current smokers appear to be at reduced risk of SARS-CoV-2 infection while former smokers appear to be at increased risk of hospitalisation, increased

disease severity and mortality from COVID-19. However, it is uncertain whether these associations are causal.

Introduction

COVID-19 is a respiratory disease caused by the SARS-CoV-2 virus. Large age and gender differences in case severity and mortality have been observed in the ongoing COVID-19 pandemic¹; however, these differences are currently unexplained. SARS-CoV-2 enters epithelial cells through the angiotensin-converting enzyme 2 (ACE-2) receptor². Some evidence suggests that gene expression and subsequent receptor levels are elevated in the airway and oral epithelium of current smokers^{3,4}, thus putting smokers at higher risk of contracting SARS-CoV-2. Other studies, however, suggest that nicotine downregulates the ACE-2 receptor⁵. These uncertainties notwithstanding, both former and current smoking is known to increase the risk of respiratory viral^{6,7} and bacterial^{8,9} infections and is associated with worse outcomes once infected. Cigarette smoke reduces the respiratory immune defence through peri-bronchiolar inflammation and fibrosis, impaired mucociliary clearance and disruption of the respiratory epithelium¹⁰. There is also reason to believe that behavioural factors (e.g. regular hand-to-mouth movements) involved in smoking may increase SARS-CoV-2 infection and transmission in current smokers. However, early data from the COVID-19 pandemic have not provided clear evidence for a negative impact of current or former smoking on SARS-CoV-2 infection or COVID-19 disease outcomes, such as hospitalisation or mortality¹¹. It has also been hypothesised that nicotine might protect against a hyper-inflammatory response to SARS-CoV-2 infection, which may lead to adverse outcomes in patients with COVID-19 disease¹².

There are several reviews that fall within the scope of smoking and COVID-19^{11,13–18}. We aimed to produce a rapid synthesis of available evidence pertaining to the rates of infection, hospitalisation, disease severity and mortality from SARS-CoV-2/COVID-19 stratified by smoking status. Given the increasing availability of data on this topic, this is a living review with regular updates. As evidence accumulates, the review will be expanded to include studies reporting COVID-19 outcomes by alternative nicotine use (e.g., nicotine replacement therapy or e-cigarettes).

Methods

Study design

This is a living evidence review which is updated as new evidence becomes available¹⁹. We adopted recommended best practice for rapid evidence reviews, which involved limiting the search to main databases and having one reviewer extract the data and another verify²⁰. This study was not pre-registered but evolved from a report written for a UK medical society²¹. The most recent (and all future) version(s) of this living review is available here (<https://www.geios.com/read/latest-UJR2AW>). A completed Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) checklist is included in Supplementary file 1.

Eligibility criteria

Studies were included if they:

- 1) Were primary research studies using experimental (e.g. randomised controlled trial), quasi-experimental (e.g. pre- and post-test) or observational (e.g. case-control, retrospective cohort, prospective cohort) study designs;
- 2) Included adults aged 16+ years;
- 3) Recorded as outcome i) results of a SARS-CoV-2 diagnostic test (including antibody assays), ii) clinical diagnosis of COVID-19, iii) hospitalisation with COVID-19, iv) severity of COVID-19 disease in those hospitalised or v) mortality from COVID-19;
- 4) Reported any of the outcomes of interest by self-reported or biochemically verified smoking status (e.g. current smoker, former smoker, never smoker) or current vaping and nicotine replacement therapy (NRT) use;
- 5) Were available in English;
- 6) Were published in a peer-reviewed journal, as a pre-print or a public health report by reputable agents (e.g. governments, scientific societies).

Search strategy

The following terms were searched for in Ovid MEDLINE (2019-search date) as free text or Medical Subject Headings:

1. Tobacco Smoking/ or Smoking Cessation/ or Water Pipe Smoking/ or Smoking/ or Smoking Pipes/ or Cigar Smoking/ or Smoking Prevention/ or Cigarette Smoking/ or smoking.mp. or Pipe Smoking/ or Smoking, Non-Tobacco Products/ or Smoking Water Pipes/
2. Nicotine/ or nicotine.mp. or Electronic Nicotine Delivery Systems/ or Nicotine Chewing Gum/
3. vaping.mp. or Vaping/
4. 1 or 2 or 3
5. Coronavirus/ or Severe Acute Respiratory Syndrome/ or Coronavirus Infections/ or covid.mp.
6. 4 and 5

The following terms were searched for in titles, abstracts and full texts in [medRxiv](#) (no time limitations):

1. covid (this term captures both covid and SARS-CoV-2) AND smoking
2. covid AND nicotine
3. covid AND vaping

Additional articles/reports of interest were identified through mailing lists, Twitter, the International Severe Acute Respiratory and Emerging Infection Consortium ([ISARIC](#)) and the US Centers for Disease Control and Prevention ([CDC](#)). Where updated versions of pre-prints or public health reports were available, old versions were superseded.

Selection of studies

One reviewer screened titles, abstracts and full texts against the inclusion criteria.

Data extraction

Data were extracted by one reviewer and verified (i.e. independently checked against pre-prints and published reports) by another on i) author (year); ii) date published; iii) country; iv) study design; v) study setting; vi) sample size; vii) sex; viii) age; ix) smoking status (e.g. current, former, never, not stated, missing); x) use of alternative nicotine products; xi) SARS-CoV-2 testing; xii) SARS-CoV-2

infection; xiii) diagnosis of COVID-19; xiv) hospitalisation with COVID-19; xv) disease severity in those hospitalised with COVID-19; and xvi) mortality.

Quality appraisal

The quality of included studies was assessed to determine suitability for inclusion in meta-analyses. Studies were judged as 'good' quality if they: i) had <20% missing data on smoking status and used a reliable self-report measure that distinguished between current, former and never smoking status; AND ii) used biochemical verification of smoking status and reported results from adjusted analyses; OR reported data from a representative/random sample. Studies were rated as 'fair' if they fulfilled only criterion i) and were otherwise rated as 'poor'. The quality appraisal was conducted by one reviewer and verified by a second.

Evidence synthesis

A narrative synthesis was conducted. Data from 'good' and 'fair' quality studies were pooled in R v.3.6.3²². In a living review where new data are regularly added to the analyses, it may be more appropriate to use a Bayesian (as opposed to frequentist) approach where prior knowledge is used in combination with new data to estimate a posterior risk distribution. A Bayesian approach mitigates against the issue of performing multiple statistical tests, which can inflate family-wise error. A series of random-effects hierarchical Bayesian meta-analyses were performed with the *brms*²³ package to estimate the relative risk for each comparison with accompanying 95% credible intervals (CrIs). We first defined prior distributions for the true pooled effect size (μ) and the between-study heterogeneity (τ), with μ specified as a normal distribution with a mean equal to the derived point estimate from each comparison of interest in the immediately preceding version of this living review²⁴, and τ specified as a half-Cauchy distribution with a mean of 0 and standard deviation of 1. The half-Cauchy distribution was selected to reflect prior knowledge that high levels of between-study heterogeneity are more likely than lower levels. Markov Chain Monte Carlo methods (20,000 burn-ins followed by 80,000 iterations) were then used to generate a risk distribution for each study, in addition to a pooled effect for the posterior risk distribution. We report forest plots with the pooled effect for the posterior risk distribution displayed as the median relative risk with an accompanying 95% CrIs. We used the empirical cumulative distribution function (ECDF) to estimate the probability of there being a 10% reduction or 10% increase in relative risk (RR) (i.e. $RR \geq 1.1$ or $RR \leq 0.9$). Due to a lack of indication as to what constitutes a clinically or epidemiologically meaningful effect (e.g. with regards to onward disease transmission or

requirements for intensive care beds), we deemed a 10% change in risk as small but important. Where data were inconclusive (as indicated by CIs crossing $RR = 1.0$), to disambiguate whether data favoured no effect or there being a small but important association, we estimated whether there was $\geq 75\%$ probability of $RR \geq 1.1$ or $RR \leq 0.9$.

Two sensitivity analyses were performed. First, a minimally informative prior for μ was specified as a normal distribution with a mean of 0 and standard deviation of 1 and τ as described above. Second, an informative prior as described above for μ was used with τ specified as a half-Cauchy distribution with a mean of 0.3 and standard deviation of 1 to reflect greater between-study heterogeneity.

To aid in the visualisation of smoking prevalence in the included studies, 95% bootstrap percentile confidence intervals were calculated for each study. We performed 1,000 bootstrap replications, with the 2.5th and 97.5th percentiles of the empirical distribution forming the 95% bootstrap percentile confidence intervals²⁵ (CIs). It should be noted that prevalence estimates in the included studies were not adjusted for age, sex, socioeconomic position, or region within countries.

Results

In the current review version (v7) with searches up to 25 August 2020, a total of 347 new records were identified, with 233 studies included in a narrative synthesis and 32 studies included in meta-analyses (see Figure 1).

Study characteristics

Characteristics of included studies are presented in Table 1. Studies were conducted across 33 countries. Sixty-two studies were conducted in the US, 53 in China, 26 in the UK, 13 in Spain, 12 in Mexico, 11 in France, seven in Italy, six across multiple international sites, four in Brazil and Iran, three in Israel and Turkey, two in Bangladesh, Chile, Denmark, Finland, India, Japan and Qatar and one from 15 further countries (see Supplementary figure S1). The majority of studies used observational designs (see Supplementary table S1). One-hundred-and-fifty-five studies were conducted in hospital settings, 62 studies included a community component in addition to hospitalised patients and 14 studies were conducted exclusively in the community, one in a quarantine centre and one did not state the study setting. Studies had a median of 404 (interquartile range = 115-1,631) participants. The majority of studies (93.5%) used reverse transcriptase polymerase chain reaction (RT-PCR) for confirmation of SARS-CoV-2 infection, 2.6% used an antibody

test to confirm prior infection, and 3.9% further studies relied on a combination of RT-PCR and clinical diagnosis (see Supplementary table S1).

Smoking status

Categorisation of smoking status was heterogeneous (see Table 1). One-hundred-and-forty-five studies collected data on smoking status through routine electronic health records (EHRs), 59 studies used a bespoke case report form for COVID-19 and 29 studies did not state the source for information on smoking status. None of the studies verified smoking status biochemically. Notably, only 57 (24.4%) studies reported current, former and never smoking status (see Supplementary table S2a), with a further 17 studies reporting ever and never smoking status (see Supplementary table S2b). The remaining 159 studies reported current, current/former or current and former smoking status but did not explicitly state whether remaining participants were never smokers or if data were missing on smoking status (see Supplementary table S2c). Seventy-eight studies explicitly reported the proportion with missing data on smoking status, which ranged from 0.08% to 96.4%.

Use of alternative nicotine products

Five studies recorded the use of alternative nicotine products in current and/or former smokers but did not report COVID-19 outcomes stratified by nicotine use^{26–30}.

Quality appraisal

One study was performed in a random, representative population sample and was rated as ‘good’ quality. Forty-six studies were rated as ‘fair’ quality. The remaining 186 studies were rated as ‘poor’ quality (see Table 1).

Smoking prevalence by country

Unadjusted smoking prevalence compared with overall estimates for national adult smoking prevalence split by country and study setting is presented in Figure 2a and 2b. Lower than expected current smoking prevalence was generally observed. Former smoking prevalence was more similar to expected prevalence when reported. National smoking prevalence estimates used for comparison are presented in Supplementary table 3.

SARS-CoV-2 testing by smoking status

Three studies provided data on access to SARS-CoV-2 diagnostic testing for those meeting local testing criteria by smoking status. In a cohort study of US military veterans aged 54-75³¹, current smokers were more likely to receive a test: 42.3% (1,603/3,789) of the sample were current smokers compared with 23.8% of all veterans aged 50+ years using any tobacco product between 2010-2015³². In the UK Biobank cohort³³, former (RR = 1.29, 95% CI = 1.14-1.45, $p < .001$) and current (RR = 1.44, 95% CI = 1.20-1.71, $p < .001$) compared with never smokers were more likely to receive a test in a multivariable analysis. In an Australian rapid assessment screening clinic for COVID-19³⁴, 9.4% (397/4,226) of the self-referred sample (subsequently assessed by a healthcare professional to decide on testing) were current smokers. Current compared with former or never smokers were less likely to require a test (RR = 0.93, 95% CI = 0.86-1.0, $p = 0.045$).

SARS-CoV-2 infection by smoking status

Forty-five studies provided data on SARS-CoV-2 infection for people meeting local testing criteria by smoking status (see Table 2). Meta-analyses were performed for one 'good' and 16 'fair' quality studies (see Figure 4 and 5). Current smokers were at reduced risk of testing positive for SARS-CoV-2 compared with never smokers (RR = 0.74, 95% CrI = 0.58-0.93, $\tau = 0.41$, 95% CI = 0.24-0.64). The probability of current smokers being at reduced risk of infection compared with never smokers (RR ≤ 0.9) was 95%. Former compared with never smokers were at increased risk of testing positive, but data were inconclusive (RR = 1.05, 95% CrI = 0.95-1.17, $\tau = 0.17$, 95% CI = 0.10-0.26) and favoured there being no important association. The probability of former smokers being at increased risk of infection (RR ≥ 1.1) compared with never smokers was 21%. Results were materially unchanged in the two sensitivity analyses (see Supplementary figure S2).

Hospitalisation for COVID-19 by smoking status

Twenty-nine studies examined hospitalisation for COVID-19 disease stratified by smoking status (see Table 3). Meta-analyses were performed for eight 'fair' quality studies (see Figure 6 and 7). Current (RR = 1.06, CrI = 0.82-1.35, $\tau = 0.27$, 95% CI = 0.08-0.55) and former (RR = 1.20, CrI = 1.03-1.44, $\tau = 0.17$, 95% CI = 0.06-0.37) compared with never smokers were at increased risk of hospitalisation with COVID-19, but data for current smokers were inconclusive and favoured there being no important association. The probability of current and former smokers being at increased risk of

hospitalisation compared with never smokers was 35% and 89%, respectively. Results were materially unchanged in two sensitivity analyses (see Supplementary figure S3).

Disease severity by smoking status

Sixty studies reported disease severity in hospitalised patients stratified by smoking status (see Table 4). Severe (as opposed to non-severe) disease was broadly defined as requiring intensive treatment unit (ITU) admission, requiring oxygen as a hospital inpatient or in-hospital death. Meta-analyses were performed for eight 'fair' quality studies (see Figure 8 and 9). Current (RR = 1.25, CrI = 0.85-1.93, τ = 0.34, 95% CI = 0.01-0.86) and former (RR = 1.52, CrI = 1.13-2.07, τ = 0.29, 95% CI = 0.47-0.66) compared with never smokers were at increased risk of greater disease severity; data for current smokers were inconclusive but favoured there being a small but important association. The probability of current and former smokers having increased risk of greater disease severity compared with never smokers was 79% and 98%, respectively. Results were materially unchanged in two sensitivity analyses (see Supplementary figure S4).

Mortality by smoking status

Fifty studies reported mortality from COVID-19 by smoking status (see Table 5), with nine 'fair' quality studies included in meta-analyses (see Figure 10 and 11). Current (RR = 1.22, 95% CrI = 0.78-1.94, τ = 0.49, 95% CI = 0.16-0.99) and former (RR = 1.39, 95% CrI = 1.09-1.87, τ = 0.27, 95% CI = 0.05-0.58) compared with never smokers were at increased risk of in-hospital mortality from COVID-19. Data for current smokers were inconclusive but favoured there being no important association. The probability of current and former smokers being at greater risk of in-hospital mortality compared with never smokers was 70% and 97%, respectively. Results were materially unchanged in two sensitivity analyses (see Supplementary figure S5).

Discussion

This living rapid review found uncertainty in the majority of 233 studies arising from the recording of smoking status. Notwithstanding these uncertainties, compared with overall adult national prevalence estimates, recorded current smoking rates in most countries were lower than expected. In a subset of better of quality studies (n = 17), current smokers had a reduced risk of testing positive

for SARS-CoV-2 but appeared more likely to present for testing and/or receive a test. Data for current smokers on the risk of hospitalisation, disease severity and mortality were inconclusive, but favoured there being no important associations with hospitalisation and mortality and a small but important increase in the risk of severe disease. Former smokers were at increased risk of hospitalisation, disease severity and mortality compared with never smokers.

Issues complicating interpretation

Interpretation of results from studies conducted during the first phase of the SARS-CoV-2 pandemic is complicated by several factors (see Figure 12):

- 1) Exposure to SARS-CoV-2 is heterogeneous with different subgroups at heightened risk of infection at different stages of the pandemic. This will likely introduce bias in studies assessing the rate of infection by smoking status conducted early on.
- 2) Current and former smokers may be more likely to meet local criteria for community testing due to increased prevalence of symptoms consistent with SARS-CoV-2 infection, such as cough, increased sputum production or altered sense of smell or taste³⁵. Evidence from a small number of studies indicates that current smokers may be more likely to present for testing, hence increasing the denominator in comparisons with never smokers and potentially inflating the rate of negative tests in current smokers. Infection positivity rates estimated among random samples will be more informative than currently available data. We identified one population study conducted in Hungary reporting on seroprevalence and smoking status³⁶; however, the response rate was fairly low at 58.8% and the current smoking rate was 10 percentage points below national prevalence estimates, thus questioning the representativeness of the final sample. Smoking status is being collected in at least two large representative infection and antibody surveys in the UK^{37,38}.
- 3) Testing for acute infection requires swabbing of the mucosal epithelium, which may be disrupted in current smokers, potentially altering the sensitivity of assays³⁹.
- 4) Diagnostic criteria for SARS-CoV-2 infection and COVID-19 have changed during the course of the pandemic⁴⁰. It was not possible to extract details on the specific RT-PCR technique or platforms used across the included studies due to reporting gaps. Different platforms have varying sensitivity and specificity to detect SARS-CoV-2 infection.
- 5) Most included studies relied on EHRs as the source of information on smoking status. Research shows large discrepancies between EHRs and actual behaviour⁴¹. Known failings of EHRs include implausible longitudinal changes, such as former smokers being recorded as never smokers at subsequent hospital visits⁴¹. Misreporting on the part of the patient (perhaps due to perceived

stigmatisation) has also been observed, with biochemical measures showing higher rates of smoking compared with self-report in hospitalised patients in the US⁴². It is hence possible that under-reporting of current and former smoking status in hospitals occurred across the included studies.

6) Individuals with severe COVID-19 symptoms may have stopped smoking immediately before admission to hospital and may therefore not have been recorded as current smokers (i.e. reverse causality).

7) Smokers with COVID-19 may be less likely to receive a SARS-CoV-2 test or present to hospital due to lack of access to healthcare and may be more likely to die in the community from sudden complications (i.e. self-selection bias) and thus not be recorded.

8) If there is a protective effect of nicotine on COVID-19 disease outcomes, abrupt nicotine withdrawal upon hospitalisation may lead to worse outcomes¹².

9) During periods of heightened demand of limited healthcare resources, current and former smokers with extensive comorbidities may have reduced priority for intensive care admission, thus leading to higher in-hospital mortality.

10) Given lack of knowledge of the disease progression and long-term outcomes of COVID-19, it is unclear whether studies conducted thus far in the pandemic have monitored patients for a sufficient time period to report complete survival outcomes or whether they are subject to early censoring.

11) Reasons for hospitalisation vary by country and time in the pandemic. For example, early cases may have been hospitalised for isolation and quarantine reasons and not due to medical necessity. It is plausible this may have skewed early data towards less severe cases. In addition, the observed association between former smoking and greater disease severity may be explained by collider bias⁴³, where conditioning on a collider (e.g. testing or hospitalisation) by design or analysis may introduce a spurious association between current or former smoking (a potential cause of testing or hospitalisation) and SARS-CoV-2 infection/adverse outcomes from COVID-19 (potentially exacerbated by smoking)⁴⁴.

Limitations

This living rapid evidence review was limited by having a single reviewer extracting data with a second independently verifying the data extracted to minimise errors, restricting the search to one electronic database and one pre-print server and by not including at least three large population surveys due to their reliance on self-reported suspected or confirmed SARS-CoV-2 infection (which means they do not meet our eligibility criteria)^{35,45,46}. We also did not include a large, UK-based,

representative seroprevalence study⁴⁷ in our meta-analyses as the odds of testing positive in former smokers was not reported. However, the odds of infection for current smokers (OR = 0.64, 95% CI = 0.58-0.71) was in concordance with the pooled estimate in our meta-analysis. Population surveys – particularly with linked data on confirmed infection or antibodies – will be included in future review versions to help mitigate some of the limitations of healthcare based observational studies. The comparisons of current and former smoking prevalence in the included studies with national prevalence estimates did not adjust observed prevalence for the demographic profile of those tested/admitted to hospital. Other reviews focused on this comparison have applied adjustments for sex and age, and continue to find lower than expected prevalence – notwithstanding the issues complicating interpretation described above¹⁷.

Implications for research, policy and practice

Further scientific research is needed to resolve the mixed findings summarised in our review. First, clinical trials of the posited therapeutic effect of nicotine could have important implications both for smokers and for improved understanding of how the SARS-CoV-2 virus causes disease in humans. Such trials should focus on medicinal nicotine (as smoked tobacco is a dirty delivery mechanism that could mask beneficial effects) and potentially differentiate between different modes of delivery (i.e. inhaled vs. ingested) since this can affect pharmacokinetics⁴⁸ and potential therapeutic effects. A second research priority would be a large, representative (randomly sampled) population survey with a validated assessment of smoking status which distinguishes between recent and long-term ex-smokers – ideally biochemically verified – and assesses seroprevalence and links to health records.

In the meantime, public-facing messages about the possible protective effect of smoking or nicotine are premature. In our view, until there is further research, the quality of the evidence does not justify the huge risk associated with a message likely to reach millions of people that a lethal activity, such as smoking, may protect against COVID-19. It continues to be appropriate to recommend smoking cessation and emphasise the role of alternative nicotine products to support smokers to stop as part of public health efforts during COVID-19. At the very least, smoking cessation reduces acute risks from cardiovascular disease and could reduce demands on the healthcare system⁴⁹. GPs and other healthcare providers can play a crucial role – brief, high-quality and free online training is available at [National Centre for Smoking Cessation and Training](#).

Conclusion

Across 234 studies, recorded smoking prevalence was generally lower than national prevalence estimates. Current smokers were at reduced risk of testing positive for SARS-CoV-2 and former smokers were at increased risk of hospitalisation, disease severity and mortality compared with never smokers.

Acknowledgements

An original short review for the Royal College of Physicians was converted to an extended living review after a request by Martin Dockrell, Tobacco Control Lead, Public Health England. All scientific decisions were made by the authors independently of funders and external organisations. The authors would like to thank Rosemary Koper for her assistance in running the electronic searches and data extraction.

Future review versions

<https://www.qeios.com/read/latest-UJR2AW>

Previous review versions

Version 1: <https://doi.org/10.32388/UJR2AW>

Version 2: <https://doi.org/10.32388/UJR2AW.3>

Version 3: <https://doi.org/10.32388/UJR2AW.4>

Version 4: <https://doi.org/10.32388/UJR2AW.5>

Version 5: <https://doi.org/10.32388/UJR2AW.6>

Version 6: <https://doi.org/10.32388/UJR2AW.7>

Data availability

All data contributing to the current and future review versions are available here:

<https://doi.org/10.6084/m9.figshare.12756020>

All code required to reproduce the current and future analyses are available here:

<https://doi.org/10.5281/zenodo.4002046>

References

- 1 Guan W, Ni Z, Hu YY, *et al.* Clinical Characteristics of Coronavirus Disease 2019 in China. *N Engl J Med* 2020; : NEJMoa2002 032.
- 2 Hoffmann M, Kleine-Weber H, Schroeder S, *et al.* SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. *Cell* 2020; published online March 5. DOI:10.1016/j.cell.2020.02.052.
- 3 Brake SJ, Barnsley K, Lu W, McAlinden KD, Eapen MS, Sohal SS. Smoking Upregulates Angiotensin-Converting Enzyme-2 Receptor: A Potential Adhesion Site for Novel Coronavirus SARS-CoV-2 (Covid-19). *J Clin Med* 2020 Vol 9 Page 841 2020; **9**: 841.
- 4 Cai G. Bulk and Single-Cell Transcriptomics Identify Tobacco-Use Disparity in Lung Gene Expression of ACE2, the Receptor of 2019-nCov. 2020; published online March 2. DOI:10.20944/PREPRINTS202002.0051.V3.
- 5 Oakes JM, Fuchs RM, Gardner JD, Lazartigues E, Yue X. Nicotine and the renin-angiotensin system. *Am. J. Physiol. - Regul. Integr. Comp. Physiol.* 2018; **315**: R895–906.
- 6 Denholm JT, Gordon CL, Johnson PD, *et al.* Hospitalised adult patients with pandemic (H1N1) 2009 influenza in Melbourne, Australia. *Med J Aust* 2010; **192**: 84–6.
- 7 Abadom TR, Smith AD, Tempia S, Madhi SA, Cohen C, Cohen AL. Risk factors associated with hospitalisation for influenza-associated severe acute respiratory illness in South Africa: A case-population study. *Vaccine* 2016; **34**: 5649–55.
- 8 Almirall J, González CA, Balanzó X, Bolívar I. Proportion of community-acquired pneumonia cases attributable to tobacco smoking. *Chest* 1999; **116**: 375–9.
- 9 Feldman C, Anderson R. Cigarette smoking and mechanisms of susceptibility to infections of the respiratory tract and other organ systems. *J. Infect.* 2013; **67**: 169–84.
- 10 Dye JA, Adler KB. Occasional review Effects of cigarette smoke on epithelial cells of the respiratory tract. *Thorax* 1994; **49**: 825–34.
- 11 Vardavas CI, Nikitara K. COVID-19 and smoking: A systematic review of the evidence. *Tob Induc Dis* 2020; **18**: 20.
- 12 Farsalinos K, Niaura R, Le Houezec J, *et al.* Editorial: Nicotine and SARS-CoV-2: COVID-19 may be a disease of the nicotinic cholinergic system. *Toxicol Rep* 2020; published online April. DOI:10.1016/j.toxrep.2020.04.012.
- 13 Emami A, Javanmardi F, Pirbonyeh N, Akbari A. Prevalence of Underlying Diseases in Hospitalized Patients with COVID-19: a Systematic Review and Meta-Analysis. *Arch Acad Emerg Med* 2020; **8**: e35.
- 14 Alqahtani JS, Oyelade T, Aldhahir AM, *et al.* Prevalence, Severity and Mortality associated with COPD and Smoking in patients with COVID-19: A Rapid Systematic Review and Meta-Analysis. *medRxiv* 2020; : 2020.03.25.20043745.

- 15 Patanavanich R, Glantz SA. Smoking is Associated with COVID-19 Progression: A Meta-Analysis. *medRxiv* 2020. DOI:10.14171/j.2095-5944.sg.2014.02.004.
- 16 Berlin I, Thomas D, Le Faou A-L, Cornuz J. COVID-19 and Smoking. *Nicotine Tob Res* DOI:10.1093/NTR/NTAA059.
- 17 Farsalinos K, Barbouni A, Niaura R. Systematic review of the prevalence of current smoking among hospitalized COVID-19 patients in China: could nicotine be a therapeutic option? *Intern Emerg Med* 2020; published online May 9. DOI:10.1007/s11739-020-02355-7.
- 18 Grundy* EJ, Suddek* T, Filippidis FT, Majeed A, Coronini-Cronberg S. Smoking, SARS-CoV-2 and COVID-19: A review of reviews considering implications for public health policy and practice. *Tob Induc Dis* 2020; **18**. DOI:10.18332/tid/124788.
- 19 Elliott JH, Turner T, Clavisi O, *et al.* Living Systematic Reviews: An Emerging Opportunity to Narrow the Evidence-Practice Gap. *PLoS Med* 2014; **11**. DOI:10.1371/journal.pmed.1001603.
- 20 Tricco AC, Antony J, Zarin W, *et al.* A scoping review of rapid review methods. *BMC Med* 2015; **13**: 224.
- 21 Simons D, Brown J, Shahab L, Perski O. Smoking and COVID-19: Rapid evidence review for the Royal College of Physicians, London (UK). *Qeios* 2020; published online April 1. DOI:10.32388/VGJCUN.
- 22 R Core Team. The R Project for Statistical Computing. 2013; : 1–12.
- 23 Bürkner P-C. Advanced Bayesian Multilevel Modeling with the R Package brms. *ArXiv170511123 Stat* 2017; published online Oct 15. <http://arxiv.org/abs/1705.11123> (accessed July 26, 2020).
- 24 Simons D, Shahab L, Brown J, Perski O. The association of smoking status with SARS-CoV-2 infection, hospitalisation and mortality from COVID-19: A living rapid evidence review (version 5). *Qeios* 2020; published online July 1. DOI:10.32388/UJR2AW.6.
- 25 Efron B. Better bootstrap confidence intervals. *J Am Stat Assoc* 1987; **82**: 171–85.
- 26 Miyara M, Tubach F, Martinez V, *et al.* Low rate of daily smokers in patients with symptomatic COVID-19. *medrxiv* 2020; : 2020.06.10.20127514.
- 27 Rimland CA, Morgan CE, Bell GJ, *et al.* Clinical characteristics and early outcomes in patients with COVID-19 treated with tocilizumab at a United States academic center. *medRxiv* 2020; : 2020.05.13.20100404.
- 28 Yannick Girardeau, Yoan Gallous, Guillaume de Bonnecaze, *et al.* Confirmed central olfactory system lesions on brain MRI in COVID-19 patients with anosmia: a case-series | *medRxiv*. <https://doi.org/10.1101/2020.07.08.20148692> (accessed Aug 25, 2020).

- 29Ebinger J, Botwin GJ, Albert CM, *et al.* SARS-CoV-2 Seroprevalence Across a Diverse Cohort of Healthcare Workers. *medRxiv* 2020; : 2020.07.31.20163055.
- 30Islam MZ, Riaz BK, Islam AS, *et al.* Risk factors associated with morbidity and mortality outcomes of COVID-19 patients on the 14th and 28th day of the disease course: a retrospective cohort study in Bangladesh. *medRxiv* 2020; : 2020.08.17.20176586.
- 31Rentsch CT, Kidwai-Khan F, Tate JP, *et al.* Covid-19 Testing, Hospital Admission, and Intensive Care Among 2,026,227 United States Veterans Aged 54-75 Years. *medRxiv* 2020; : 2020.04.09.20059964.
- 32Odani S. Tobacco Product Use Among Military Veterans — United States, 2010–2015. *MMWR Morb Mortal Wkly Rep* 2018; **67**. DOI:10.15585/mmwr.mm6701a2.
- 33Niedzwiedz CL, O'Donnell CA, Jani BD, *et al.* Ethnic and socioeconomic differences in SARS-CoV-2 infection: prospective cohort study using UK Biobank. *BMC Med* 2020; **18**: 160.
- 34Trubiano JA, Vogrin S, Smibert OC, *et al.* COVID-MATCH65 - A prospectively derived clinical decision rule for severe acute respiratory syndrome coronavirus 2. *medRxiv* 2020; : 2020.06.30.20143818.
- 35Hopkinson NS, Rossi NN, Moustafa JE-SSE, *et al.* Current tobacco smoking and risk from COVID-19 results from a population symptom app in over 2.4 million people. *medrxiv* 2020; **44**: 2020.05.18.20105288.
- 36Merkely B, Szabó AJ, Kosztin A, *et al.* Novel coronavirus epidemic in the Hungarian population, a cross-sectional nationwide survey to support the exit policy in Hungary. *GeroScience* 2020; published online July 17. DOI:10.1007/s11357-020-00226-9.
- 37Major home testing programme for coronavirus will track levels of infection in the community - GOV.UK. <https://www.gov.uk/government/news/major-home-testing-programme-for-coronavirus-will-track-levels-of-infection-in-the-community> (accessed May 22, 2020).
- 38COVID-19 Infection Survey (CIS) - Office for National Statistics. <https://www.ons.gov.uk/surveys/informationforhouseholdsandindividuals/householdandindividualsurveys/covid19infectionsurveycis> (accessed June 30, 2020).
- 39de Lusignan S, Dorward J, Correa A, *et al.* Risk factors for SARS-CoV-2 among patients in the Oxford Royal College of General Practitioners Research and Surveillance Centre primary care network: a cross-sectional study. *Lancet Infect Dis* 2020; **0**. DOI:10.1016/S1473-3099(20)30371-6.
- 40World Health Organisation. Laboratory testing for 2019 novel coronavirus (2019-nCoV) in suspected human cases. <https://www.who.int/publications-detail-redirect/10665-331501> (accessed July 29, 2020).

- 41 Polubriaginof F, Salmasian H, Albert DA, Vawdrey DK. Challenges with Collecting Smoking Status in Electronic Health Records. *AMIA Annu Symp Proc AMIA Symp* 2017; **2017**: 1392–400.
- 42 Benowitz NL, Schultz KE, Haller CA, Wu AHB, Dains KM, Jacob P. Prevalence of smoking assessed biochemically in an urban public hospital: a rationale for routine cotinine screening. *Am J Epidemiol* 2009; **170**: 885–91.
- 43 Griffith G, Morris TT, Tudball M, *et al.* Collider bias undermines our understanding of COVID-19 disease risk and severity. *medRxiv* 2020; : 2020.05.04.20090506.
- 44 Murray E. Causation in smoking and COVID-19. Twitter. 2020. <https://twitter.com/EpiEllie/status/1258607277357006849?s=20>.
- 45 Bowyer RCE, Varsavsky T, Carole H. Geo-social gradients in predicted COVID-19 prevalence and severity in Great Britain : results from Affiliations : Corresponding authors : Understanding the geographical distribution of COVID-19 through the general population is key to the provision of ade. 2020.
- 46 Jackson SE, Brown J, Shahab L, Steptoe A, Fancourt D. COVID-19, smoking, and inequalities: a cross-sectional survey of adults in the UK. *Submitted* 2020.
- 47 Ward H, Atchison CJ, Whitaker M, *et al.* Antibody prevalence for SARS-CoV-2 in England following first peak of the pandemic: REACT2 study in 100,000 adults. *medRxiv* 2020; : 2020.08.12.20173690.
- 48 Shahab L, Brose LS, West R. Novel delivery systems for nicotine replacement therapy as an aid to smoking cessation and for harm reduction: Rationale, and evidence for advantages over existing systems. *CNS Drugs* 2013; **27**: 1007–19.
- 49 Stead LF, Buitrago D, Preciado N, Sanchez G, Hartmann-Boyce J, Lancaster T. Physician advice for smoking cessation. *Cochrane Database Syst. Rev.* 2013; **2017**. DOI:10.1002/14651858.CD000165.pub4.
- 50 Guan W, Liang W, Zhao Y, *et al.* Comorbidity and its impact on 1590 patients with COVID-19 in China: a nationwide analysis. *Eur Respir J* 2020; **55**. DOI:10.1183/13993003.00547-2020.
- 51 Lian J, Jin X, Hao S, *et al.* Analysis of Epidemiological and Clinical Features in Older Patients With Coronavirus Disease 2019 (COVID-19) Outside Wuhan. *Clin Infect Dis* DOI:10.1093/cid/ciaa242.
- 52 Jin X, Lian J-S, Hu J-H, *et al.* Epidemiological, clinical and virological characteristics of 74 cases of coronavirus-infected disease 2019 (COVID-19) with gastrointestinal symptoms. *Gut* 2020; **69**: 1002–9.
- 53 Chen T, Wu D, Chen H, *et al.* Clinical characteristics of 113 deceased patients with coronavirus disease 2019: retrospective study. *BMJ* 2020; **368**. DOI:10.1136/bmj.m1091.

- 54 Zhou F, Yu T, Du R, *et al.* Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *The Lancet* 2020; **395**: 1054–62.
- 55 Mo P, Xing Y, Xiao Y, *et al.* Clinical characteristics of refractory COVID-19 pneumonia in Wuhan, China. *Clin Infect Dis* DOI:10.1093/cid/ciaa270.
- 56 Zhang J, Dong X, Cao Y, *et al.* Clinical characteristics of 140 patients infected with SARS-CoV-2 in Wuhan, China. *Allergy* 2020; **75**: 1730–41.
- 57 Wan S, Xiang Y, Fang W, *et al.* Clinical features and treatment of COVID-19 patients in northeast Chongqing. *J Med Virol* 2020; **92**: 797–806.
- 58 Liu W, Tao Z-W, Wang L, *et al.* Analysis of factors associated with disease outcomes in hospitalized patients with 2019 novel coronavirus disease. *Chin Med J (Engl)* 2020; **133**: 1032–1038.
- 59 Huang C, Wang Y, Li X, *et al.* Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *The Lancet* 2020; **395**: 497–506.
- 60 Zhang X, Cai H, Hu J, *et al.* Epidemiological, clinical characteristics of cases of SARS-CoV-2 infection with abnormal imaging findings. *Int J Infect Dis* 2020; **94**: 81–7.
- 61 Guo T, Fan Y, Chen M, *et al.* Cardiovascular Implications of Fatal Outcomes of Patients With Coronavirus Disease 2019 (COVID-19). *JAMA Cardiol* 2020; **5**: 811–8.
- 62 Liu R, Ming X, Xu O, *et al.* Association of Cardiovascular Manifestations with In-hospital Outcomes in Patients with COVID-19: A Hospital Staff Data. *medRxiv* 2020; : 2020.02.29.20029348.
- 63 Huang Y, Yang R, Xu Y, Gong P. Clinical characteristics of 36 non-survivors with COVID-19 in Wuhan, China. *medRxiv* 2020; : 2020.02.27.20029009.
- 64 Xu H, Hou K, Xu H, *et al.* Acute Myocardial Injury of Patients with Coronavirus Disease 2019. *medRxiv* 2020; : 2020.03.05.20031591.
- 65 Li J, Li S, Cai Y, *et al.* Epidemiological and Clinical Characteristics of 17 Hospitalized Patients with 2019 Novel Coronavirus Infections Outside Wuhan, China. *medRxiv* 2020; : 2020.02.11.20022053.
- 66 Hu L, Chen S, Fu Y, *et al.* Risk Factors Associated with Clinical Outcomes in 323 COVID-19 Patients in Wuhan, China. *medRxiv* 2020; : 2020.03.25.20037721.
- 67 Wang R, Pan M, Zhang X, *et al.* Epidemiological and clinical features of 125 Hospitalized Patients with COVID-19 in Fuyang, Anhui, China. *Int J Infect Dis* 2020; **95**: 421–8.
- 68 CDCMMWR. Preliminary Estimates of the Prevalence of Selected Underlying Health Conditions Among Patients with Coronavirus Disease 2019 — United States, February 12–March 28, 2020. *MMWR Morb Mortal Wkly Rep* 2020; **69**. DOI:10.15585/mmwr.mm6913e2.

- 69 Dong X, Cao Y, Lu X, *et al.* Eleven faces of coronavirus disease 2019. *Allergy* 2020; **75**: 1699–709.
- 70 Kim ES, Chin BS, Kang CK, *et al.* Clinical Course and Outcomes of Patients with Severe Acute Respiratory Syndrome Coronavirus 2 Infection: a Preliminary Report of the First 28 Patients from the Korean Cohort Study on COVID-19. *J Korean Med Sci* 2020; **35**. DOI:10.3346/jkms.2020.35.e142.
- 71 Shi Y, Yu X, Zhao H, Wang H, Zhao R, Sheng J. Host susceptibility to severe COVID-19 and establishment of a host risk score: findings of 487 cases outside Wuhan. *Crit Care* 2020; **24**: 108.
- 72 Yang X, Yu Y, Xu J, *et al.* Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study. *Lancet Respir Med* 2020; **8**: 475–81.
- 73 Argenziano MG, Bruce SL, Slater CL, *et al.* Characterization and clinical course of 1000 patients with coronavirus disease 2019 in New York: retrospective case series. *BMJ* 2020; **369**. DOI:10.1136/bmj.m1996.
- 74 Solís P, Carreño H. COVID-19 Fatality and Comorbidity Risk Factors among Confirmed Patients in Mexico. *Epidemiology*, 2020 DOI:10.1101/2020.04.21.20074591.
- 75 Richardson S, Hirsch JS, Narasimhan M, *et al.* Presenting Characteristics, Comorbidities, and Outcomes Among 5700 Patients Hospitalized With COVID-19 in the New York City Area. *JAMA* 2020; **323**: 2052–9.
- 76 Fontanet A, Tondeur L, Madec Y, *et al.* Cluster of COVID-19 in northern France: A retrospective closed cohort study. *medRxiv* 2020; : 2020.04.18.20071134.
- 77 Zheng KI, Gao F, Wang X-B, *et al.* Letter to the Editor: Obesity as a risk factor for greater severity of COVID-19 in patients with metabolic associated fatty liver disease. *Metabolism* 2020; **108**: 154244.
- 78 Liao Y, Feng Y, Wang B, *et al.* Clinical Characteristics and Risk factors for developed COVID-19 patients transferring to designated hospital from Jiangnan Fangcang shelter Hospital: a retrospective, observational study. *medRxiv* 2020; : 2020.04.21.20074724.
- 79 Gil-Agudo A, Rodriguez-Cola M, Jimenez-Velasco I, *et al.* Clinical features of coronavirus disease 2019 (COVID-19) in a cohort of patients with disability due to spinal cord injury. *medRxiv* 2020; : 2020.04.20.20072918.
- 80 Shi P, Ren G, Yang J, *et al.* Clinical characteristics of imported and second-generation COVID-19 cases outside Wuhan, China: A multicenter retrospective study. *medRxiv* 2020; : 2020.04.19.20071472.
- 81 Hadjadj J, Yatim N, Barnabei L, *et al.* Impaired type I interferon activity and exacerbated inflammatory responses in severe Covid-19 patients. *medRxiv* 2020; : 2020.04.19.20068015.

- 82 Gold JAW, Wong KK, Szablewski CM, *et al.* Characteristics and Clinical Outcomes of Adult Patients Hospitalized with COVID-19 — Georgia, March 2020. *MMWR Morb Mortal Wkly Rep* 2020; **69**: 545–50.
- 83 Yu T, Cai S, Zheng Z, *et al.* Association Between Clinical Manifestations and Prognosis in Patients with COVID-19. *Clin Ther* 2020; **42**: 964–72.
- 84 Zheng Y, Xiong C, Liu Y, *et al.* Epidemiological and clinical characteristics analysis of COVID-19 in the surrounding areas of Wuhan, Hubei Province in 2020. *Pharmacol Res* 2020; **157**: 104821.
- 85 Rica R de la, Borges M, Aranda M, *et al.* Low albumin levels are associated with poorer outcomes in a case series of COVID-19 patients in Spain: a retrospective cohort study. *medRxiv* 2020; : 2020.05.07.20094987.
- 86 Yin R, Yang Z, Wei Y, *et al.* Clinical characteristics of 106 patients with neurological diseases and co-morbid coronavirus disease 2019: a retrospective study. *medRxiv* 2020; : 2020.04.29.20085415.
- 87 Shi H, Zuo Y, Yalavarthi S, *et al.* Neutrophil calprotectin identifies severe pulmonary disease in COVID-19. *medRxiv* 2020; : 2020.05.06.20093070.
- 88 Cho ER, Slutsky AS, Jha P. Smoking and the risk of COVID-19 infection in the UK Biobank Prospective Study. *medRxiv* 2020; : 2020.05.05.20092445.
- 89 Allenbach Y, Saadoun D, Maalouf G, *et al.* Multivariable prediction model of intensive care unit transfer and death: a French prospective cohort study of COVID-19 patients. *medRxiv* 2020; : 2020.05.04.20090118.
- 90 Robilotti EV, Babady NE, Mead PA, *et al.* Determinants of Severity in Cancer Patients with COVID-19 Illness. *medRxiv* 2020; : 2020.05.04.20086322.
- 91 Williamson EJ, Walker AJ, Bhaskaran K, *et al.* OpenSAFELY: factors associated with COVID-19 death in 17 million patients. *Nature* 2020; : 1–11.
- 92 Borobia AM, Carcas AJ, Arnalich F, *et al.* A Cohort of Patients with COVID-19 in a Major Teaching Hospital in Europe. *J Clin Med* 2020; **9**: 1733.
- 93 Giacomelli A, Ridolfo AL, Milazzo L, *et al.* 30-day mortality in patients hospitalized with COVID-19 during the first wave of the Italian epidemic: A prospective cohort study. *Pharmacol Res* 2020; **158**: 104931.
- 94 Shah SJ, Barish PN, Prasad PA, *et al.* Clinical features, diagnostics, and outcomes of patients presenting with acute respiratory illness: a comparison of patients with and without COVID-19. *medRxiv* 2020; : 2020.05.02.20082461.
- 95 Kolin DA, Kulm S, Elemento O. Clinical and Genetic Characteristics of Covid-19 Patients from UK Biobank. *medRxiv* 2020; : 2020.05.05.20075507.

- 96 Lubetzky M, Aull M, Craig-Shapiro R, *et al.* Kidney Allograft Recipients Diagnosed with Coronavirus Disease-2019: A Single Center Report. *medRxiv* 2020; : 2020.04.30.20086462.
- 97 Goyal P, Choi JJ, Pinheiro LC, *et al.* Clinical Characteristics of Covid-19 in New York City. *N Engl J Med* 2020; **382**: 2372–4.
- 98 Feng Y, Ling Y, Bai T, *et al.* COVID-19 with Different Severities: A Multicenter Study of Clinical Features. *Am J Respir Crit Care Med* 2020; **201**: 1380–8.
- 99 Yao Q, Wang P, Wang X, Qie G, Chu Y. A retrospective study of risk factors for severe acute respiratory syndrome coronavirus 2 infections in hospitalized adult patients. DOI:10.20452/pamw.15312.
- 100 Sami R, Soltaninejad F, Amra B, *et al.* A one-year hospital-based prospective COVID-19 open-cohort in the Eastern Mediterranean region: The Khorshid COVID Cohort (KCC) study. *medRxiv* 2020; : 2020.05.11.20096727.
- 101 Almazeedi S, Youha SA, Jamal MH, *et al.* Clinical Characteristics, Risk Factors and Outcomes Among the First Consecutive 1,096 Patients Diagnosed with COVID-19: The Kuwait Experience. *medRxiv* 2020; : 2020.05.09.20096495.
- 102 Carrillo-Vega MF, Salinas-Escudero G, Garcia-Peña C, Gutierrez-Robledo LM, Parra-Rodriguez L. Early estimation of the risk factors for hospitalisation and mortality by COVID-19 in Mexico. *medRxiv* 2020; : 2020.05.11.20098145.
- 103 Yanover C, Mizrahi B, Kalkstein N, *et al.* What factors increase the risk of complications in SARS-CoV-2 positive patients? A cohort study in a nationwide Israeli health organization. *medRxiv* 2020; : 2020.05.07.20091652.
- 104 Hamer M, Kivimäki M, Gale CR, Batty GD. Lifestyle risk factors, inflammatory mechanisms, and COVID-19 hospitalization: A community-based cohort study of 387,109 adults in UK. *Brain Behav Immun* 2020; **87**: 184–7.
- 105 Regina J, Papadimitriou-Olivgeris M, Burger R, *et al.* Epidemiology, risk factors and clinical course of SARS-CoV-2 infected patients in a Swiss university hospital: an observational retrospective study. *medRxiv* 2020; : 2020.05.11.20097741.
- 106 de Lusignan S, Dorward J, Correa A, *et al.* Risk factors for SARS-CoV-2 among patients in the Oxford Royal College of General Practitioners Research and Surveillance Centre primary care network: a cross-sectional study. *Lancet Infect Dis* 2020; : S1473309920303716.
- 107 Targher G, Mantovani A, Wang X-B, *et al.* Patients with diabetes are at higher risk for severe illness from COVID-19. *Diabetes Metab* 2020; published online May 13. DOI:10.1016/j.diabet.2020.05.001.
- 108 Valenti L, Bergna A, Pelusi S, *et al.* SARS-CoV-2 seroprevalence trends in healthy blood donors during the COVID-19 Milan outbreak. *medRxiv* 2020; : 2020.05.11.20098442.

- 109 Feuth T, Saaresranta T, Karlsson A, *et al.* Is sleep apnoea a risk factor for Covid-19? Findings from a retrospective cohort study. *medRxiv* 2020; : 2020.05.14.20098319.
- 110 Ge H, Zhu M, Du J, *et al.* Cardiac Structural and Functional Characteristics in Patients with Coronavirus Disease 2019: A Serial Echocardiographic Study. *medRxiv* 2020; : 2020.05.12.20095885.
- 111 Parrotta E, Kister I, Charvet L, *et al.* COVID-19 outcomes in MS: Observational study of early experience from NYU Multiple Sclerosis Comprehensive Care Center. *Neurol - Neuroimmunol Neuroinflammation* 2020; **7**: e835.
- 112 Shekhar R, Sheikh AB, Upadhyay S, Atencio J, Kapuria D. Early experience with COVID-19 patients at academic hospital in Southwestern United States. *Infect Dis* 2020; **52**: 596–9.
- 113 Mejia-Vilet JM, Cordova-Sanchez BM, Fernandez-Camargo D, Mendez-Perez RA, Morales-Buenrostro LE, Hernandez-Gilsoul T. A Risk Score to Predict Admission to Intensive Care Unit in Patients With COVID-19: The ABC-GOALS Score. *medRxiv* 2020; : 2020.05.12.20099416.
- 114 Chen C, Jiang J, Xu X, Hu Y, Hu Y, Zhao Y. Dynamic liver function indexes monitoring and clinical characteristics in three types of COVID-19 patients. *medRxiv* 2020; : 2020.05.13.20099614.
- 115 Li J, Chen Y, Chen S, *et al.* Derivation and validation of a prognostic model for predicting in-hospital mortality in patients admitted with COVID-19 in Wuhan, China: the PLANS (Platelet Lymphocyte Age Neutrophil Sex) model. *medRxiv* 2020; : 2020.05.13.20100370.
- 116 Palaodimos L, Kokkinidis DG, Li W, *et al.* Severe obesity, increasing age and male sex are independently associated with worse in-hospital outcomes, and higher in-hospital mortality, in a cohort of patients with COVID-19 in the Bronx, New York. *Metabolism* 2020; **108**: 154262.
- 117 Ip A, Berry DA, Hansen E, *et al.* Hydroxychloroquine and Tocilizumab Therapy in COVID-19 Patients - An Observational Study. *medRxiv* 2020; : 2020.05.21.20109207.
- 118 Heili-Frades S. COVID-19 Outcomes in 4712 consecutively confirmed SARS-CoV2 cases in the city of Madrid. *medRxiv* <https://doi.org/10.1101/2020.05.22.20109850> (accessed July 27, 2020).
- 119 Vaquero LM, Barrado MES, Escobar D, *et al.* C-Reactive protein and SOFA score as early predictors of critical care requirement in patients with COVID-19 pneumonia in Spain. *medRxiv* 2020; : 2020.05.22.20110429.
- 120 Kim L, Garg S, O'Halloran A, *et al.* Interim Analysis of Risk Factors for Severe Outcomes among a Cohort of Hospitalized Adults Identified through the U.S. Coronavirus Disease 2019 (COVID-19)-Associated Hospitalization Surveillance Network (COVID-NET). *medRxiv* 2020; : 2020.05.18.20103390.

- 121 Wu MA, Fossali T, Pandolfi L, *et al.* COVID-19: the key role of pulmonary capillary leakage. An observational cohort study. *medRxiv* 2020; : 2020.05.17.20104877.
- 122 Shi Q, Zhao K, Yu J, *et al.* Clinical characteristics of 101 COVID-19 nonsurvivors in Wuhan, China: a retrospective study. *medRxiv* 2020; : 2020.03.04.20031039.
- 123 Al-Hindawi A, Sokhi J, Cuddihy J, *et al.* COVID-19 in London, a Case Series Demonstrating Late Improvement in Survivors. *medRxiv* 2020; : 2020.05.16.20103853.
- 124 Basse C, Diakite S, Servois V, *et al.* Characteristics and outcome of SARS-CoV-2 infection in cancer patients. *medRxiv* 2020; : 2020.05.14.20101576.
- 125 Freitas D, Leon L, Mucientes A, *et al.* Risk factors for hospital admission related to COVID-19 in inflammatory rheumatic diseases. *medRxiv* 2020; : 2020.05.14.20101584.
- 126 Alshami AA, Alattas RA, Anan HF, *et al.* Silent Disease and Loss of Taste and Smell are Common Manifestations of SARS-COV-2 Infection in a Quarantine Facility: First report from Saudi Arabia. *medRxiv* 2020; : 2020.05.13.20100222.
- 127 Berumen J, Schmulson M, Alegre J, *et al.* Risk of infection and hospitalization by Covid-19 in Mexico: a case-control study. *medRxiv* 2020; : 2020.05.24.20104414.
- 128 Gianfrancesco M, Hyrich KL, Al-Adely S, *et al.* Characteristics associated with hospitalisation for COVID-19 in people with rheumatic disease: data from the COVID-19 Global Rheumatology Alliance physician-reported registry. *Ann Rheum Dis* 2020; **79**: 859–66.
- 129 Li J, Long X, Zhu C, *et al.* Olfactory Dysfunction in Recovered Coronavirus Disease 2019 (COVID-19) Patients. *Mov Disord*; **n/a**. DOI:10.1002/mds.28172.
- 130 Batty GD, Deary I, Luciano M, Altschul D, Kivimaki M, Gale C. Psychosocial factors and hospitalisations for COVID-19: Prospective cohort study of the general population. *medRxiv* 2020; : 2020.05.29.20100735.
- 131 Israel A, Feldhamer I, Lahad A, Levin-Zamir D, Lavie G. Smoking and the risk of COVID-19 in a large observational population study. *medRxiv* 2020; : 2020.06.01.20118877.
- 132 Valle DMD, Kim-schulze S, Hsin-hui H, *et al.* An inflammatory cytokine signature helps predict COVID-19 severity and death. *medRxiv* 2020; : 2020.05.28.20115758.
- 133 Chaudhry F, Bulka H, Rathnam AS, *et al.* COVID-19 in Multiple Sclerosis Patients and Risk Factors for Severe Infection. *medRxiv* 2020; : 2020.05.27.20114827.
- 134 Louis S, Dhawan A, Newey C, *et al.* Continuous Electroencephalography (cEEG) Characteristics and Acute Symptomatic Seizures in COVID-19 Patients. *medRxiv* 2020; : 2020.05.26.20114033.

- 135 Soto-Mota A, Garza BAM, Rodriguez EM, *et al.* THE LOW-HARM SCORE FOR PREDICTING MORTALITY IN PATIENTS DIAGNOSED WITH COVID-19: A MULTICENTRIC VALIDATION STUDY. *medRxiv* 2020; : 2020.05.26.20111120.
- 136 Garibaldi BT, Fiksel J, Muschelli J, *et al.* Patient trajectories and risk factors for severe outcomes among persons hospitalized for COVID-19 in the Maryland/DC region. *medRxiv* 2020; : 2020.05.24.20111864.
- 137 Docherty AB, Harrison EM, Green CA, *et al.* Features of 20 133 UK patients in hospital with covid-19 using the ISARIC WHO Clinical Characterisation Protocol: prospective observational cohort study. *BMJ* 2020; **369**. DOI:10.1136/bmj.m1985.
- 138 Boulware DR, Pullen MF, Bangdiwala AS, *et al.* A Randomized Trial of Hydroxychloroquine as Postexposure Prophylaxis for Covid-19. *N Engl J Med* 2020; published online June 3. DOI:10.1056/NEJMoa2016638.
- 139 Kuderer NM, Choueiri TK, Shah DP, *et al.* Clinical impact of COVID-19 on patients with cancer (CCC19): a cohort study. *The Lancet* 2020; **395**: 1907–18.
- 140 Romão VC, Oliveira-Ramos F, Cruz-Machado AR, *et al.* A COVID-19 outbreak in a rheumatology department upon the early days of the pandemic. *medRxiv* 2020; : 2020.06.05.20107011.
- 141 Giannouchos T, Sussman R, Mier JM, Poulas K, Farsalinos K. Characteristics and risk factors for COVID-19 diagnosis and adverse outcomes in Mexico: an analysis of 89,756 laboratory-confirmed COVID-19 cases. *medRxiv* 2020; : 2020.06.04.20122481.
- 142 Ramlall V, Thangaraj P, Meydan C, *et al.* Identification of Immune complement function as a determinant of adverse SARS-CoV-2 infection outcome. *medRxiv* 2020; : 2020.05.05.20092452.
- 143 Wang B, Oekelen OV, Mouhieddine T, *et al.* A tertiary center experience of multiple myeloma patients with COVID-19: lessons learned and the path forward. *medRxiv* 2020; : 2020.06.04.20122846.
- 144 Perrone F, Piccirillo MC, Ascierto PA, *et al.* Tocilizumab for patients with COVID-19 pneumonia. The TOCIVID-19 prospective phase 2 trial. *medRxiv* 2020; : 2020.06.01.20119149.
- 145 Sharma AK, Ahmed A, Baig VN, *et al.* Characteristics and Outcomes of Hospitalized Young Adults with Mild to Moderate Covid-19 at a University Hospital in India. *medRxiv* 2020; : 2020.06.02.20106310.
- 146 Eugen-Olsen J, Altintas I, Tingleff J, *et al.* Low levels of the prognostic biomarker suPAR are predictive of mild outcome in patients with symptoms of COVID-19 - a prospective cohort study. *medRxiv* 2020; : 2020.05.27.20114678.
- 147 Martinez-Portilla RJ, Sotiriadis A, Torres-Torres J, *et al.* Risk factors for mortality in pregnant women with SARS-CoV-2 infection. *medRxiv* 2020; : 2020.05.31.20107276.

- 148 Raisi-Estabragh Z, McCracken C, Bethell MS, *et al.* Greater risk of severe COVID-19 in Black, Asian and Minority Ethnic populations is not explained by cardiometabolic, socioeconomic or behavioural factors, or by 25(OH)-vitamin D status: study of 1326 cases from the UK Biobank. *J Public Health* DOI:10.1093/pubmed/fdaa095.
- 149 Luo H, Liu S, Wang Y, *et al.* Age differences in clinical features and outcomes in patients with COVID-19, Jiangsu, China: a retrospective, multi-center cohort study. *medRxiv* 2020; : 2020.06.01.20086025.
- 150 Houlihan CF, Vora N, Byrne T, *et al.* Pandemic peak SARS-CoV-2 infection and seroconversion rates in London frontline health-care workers. *The Lancet* 2020; : S0140673620314847.
- 151 Cen Y, Chen X, Shen Y, *et al.* Risk factors for disease progression in patients with mild to moderate coronavirus disease 2019—a multi-centre observational study. *Clin Microbiol Infect* 2020; : S1198743X20303414.
- 152 Klang E, Kassim G, Soffer S, Freeman R, Levin MA, Reich DL. Morbid Obesity as an Independent Risk Factor for COVID-19 Mortality in Hospitalized Patients Younger than 50. *Obesity*; n/a. DOI:10.1002/oby.22913.
- 153 Maraschini A, Corsi E, Salvatore MA, Donati S. Coronavirus and birth in Italy: results of a national population-based cohort study. *medRxiv* 2020; : 2020.06.11.20128652.
- 154 Wang A-L, Zhong X, Hurd Y. Comorbidity and Sociodemographic determinants in COVID-19 Mortality in an US Urban Healthcare System. *medRxiv* 2020; : 2020.06.11.20128926.
- 155 McQueenie R, Foster H, Jani BD, *et al.* Multimorbidity, Polypharmacy, and COVID-19 infection within the UK Biobank cohort. *medRxiv* 2020; : 2020.06.10.20127563.
- 156 Apea VJ, Wan YI, Dhairyan R, *et al.* Ethnicity and outcomes in patients hospitalised with COVID-19 infection in East London: an observational cohort study. *medRxiv* 2020; : 2020.06.10.20127621.
- 157 Woolford SJ, D'angelo S, Curtis EM, *et al.* COVID-19 and associations with frailty and multimorbidity: a prospective analysis of UK Biobank participants. *medRxiv* 2020; : 2020.06.09.20126292.
- 158 Hultcrantz M, Richter J, Rosenbaum C, *et al.* COVID-19 infections and outcomes in patients with multiple myeloma in New York City: a cohort study from five academic centers. *medRxiv* 2020; : 2020.06.09.20126516.
- 159 Rajter JC, Sherman M, Fatteh N, Vogel F, Sacks J, Rajter J-J. ICON (Ivermectin in Covid Nineteen) study: Use of Ivermectin is Associated with Lower Mortality in Hospitalized Patients with COVID19. *medRxiv* 2020; : 2020.06.06.20124461.

- 160 Lan F-Y, Suharlim C, Kales SN, Yang J. Association between SARS-CoV-2 infection, exposure risk and mental health among a cohort of essential retail workers in the United States. *medRxiv* 2020; : 2020.06.08.20125120.
- 161 Zeng H, Zhang T, He X, *et al.* Impact of Chronic Comorbidities on Progression and Prognosis in Patients with COVID-19: A Retrospective Cohort Study in 1031 Hospitalized Cases in Wuhan, China. *medRxiv* 2020; : 2020.06.14.20125997.
- 162 Suleyman G, Fadel RA, Malette KM, *et al.* Clinical Characteristics and Morbidity Associated With Coronavirus Disease 2019 in a Series of Patients in Metropolitan Detroit. *JAMA Netw Open* 2020; **3**: e2012270–e2012270.
- 163 Chen L, Yu J, He W, *et al.* Risk factors for death in 1859 subjects with COVID-19. *Leukemia* 2020; : 1–11.
- 164 Garassino MC, Whisenant JG, Huang L-C, *et al.* COVID-19 in patients with thoracic malignancies (TERAVOLT): first results of an international, registry-based, cohort study. *Lancet Oncol* 2020; **21**: 914–22.
- 165 Hernández-Garduño E. Obesity is the comorbidity more strongly associated for Covid-19 in Mexico. A case-control study. *Obes Res Clin Pract* 2020; published online June 12. DOI:10.1016/j.orpc.2020.06.001.
- 166 Govind R, Freitas DF de, Pritchard MR, Hayes RD, MacCabe JH. Clozapine treatment and risk of COVID-19. *medRxiv* 2020; : 2020.06.17.20133595.
- 167 Sisó-Almirall A, Kostov B, Mas-Heredia M, *et al.* PROGNOSTIC FACTORS IN SPANISH COVID-19 PATIENTS: A CASE SERIES FROM BARCELONA. *medRxiv* 2020; : 2020.06.18.20134510.
- 168 Gu T, Mack JA, Salvatore M, *et al.* COVID-19 outcomes, risk factors and associations by race: a comprehensive analysis using electronic health records data in Michigan Medicine. *medRxiv* 2020; : 2020.06.16.20133140.
- 169 Kibler M, Carmona A, Marchandot B, *et al.* Risk and severity of COVID-19 and ABO blood group in transcatheter aortic valve patients. *medRxiv* 2020; : 2020.06.13.20130211.
- 170 Ikitimur H, Uysal BB, Cengiz M, *et al.* "Determining Host Factors Contributing to Disease Severity in a Family Cluster of 29 Hospitalized SARS-CoV-2 Patients: Could Genetic Factors Be Relevant in the Clinical Course of COVID-19?". *J Med Virol*; **n/a**. DOI:10.1002/jmv.26106.
- 171 Sierpiński R, Pinkas J, Jankowski M, *et al.* Sex differences in the frequency of gastrointestinal symptoms and olfactory or taste disorders among 1,942 non-hospitalized patients with COVID-19. *Pol Arch Intern Med* 2020; published online June 3. DOI:10.20452/pamw.15414.

- 172 Zhou Y, He X, Zhang J, *et al.* Prolonged SARS-CoV-2 Viral Shedding in Patients with COVID-19 was Associated with Delayed Initiation of Arbidol Treatment: a retrospective cohort study. *medRxiv* 2020; : 2020.06.09.20076646.
- 173 Crovetto F, Crispi F, Llurba E, Figueras F, Gomez-Roig MD, Gratacos E. SEROPREVALENCE AND CLINICAL SPECTRUM OF SARS-CoV-2 INFECTION IN THE FIRST VERSUS THIRD TRIMESTER OF PREGNANCY. *medRxiv* 2020; : 2020.06.17.20134098.
- 174 Veras FP, Pontelli M, Silva C, *et al.* SARS-CoV-2 triggered neutrophil extracellular traps (NETs) mediate COVID-19 pathology. *medRxiv* 2020; : 2020.06.08.20125823.
- 175 Sterlin D, Mathian A, Miyara M, *et al.* IgA dominates the early neutralizing antibody response to SARS-CoV-2. *medRxiv* 2020; : 2020.06.10.20126532.
- 176 Rossi B, Nguyen LS, Zimmermann P, *et al.* Effect of tocilizumab in hospitalized patients with severe pneumonia COVID-19: a cohort study. *medRxiv* 2020; : 2020.06.06.20122341.
- 177 Duan L, Zhang S, Guo M, *et al.* Epidemiological and clinical characteristics in patients with SARS-CoV-2 antibody negative probable COVID-19 in Wuhan. *medRxiv* 2020; : 2020.06.18.20134619.
- 178 Martin-Jimenez P, Munoz-Garcia MI, Seoane D, *et al.* Cognitive impairment is a common comorbidity in COVID-19 deceased patients. A hospital-based retrospective cohort study. *medRxiv* 2020; : 2020.06.08.20125872.
- 179 Elezkurtaj S, Greuel S, Ihlow J, *et al.* Causes of Death and Comorbidities in Patients with COVID-19. *medRxiv* 2020; : 2020.06.15.20131540.
- 180 Lenka J, Chhabria MS, Sharma N, *et al.* Clinical characteristics and outcomes of critically ill patients with COVID-19 in a tertiary community hospital in upstate New York. *medRxiv* 2020; : 2020.06.18.20135046.
- 181 Olivares F, Munoz D, Fica A, *et al.* Covid-19 in Chile. The experience of a Regional reference Center. Preliminary report. *medRxiv* 2020; : 2020.06.14.20130898.
- 182 Salton F, Confalonieri P, Santus P, *et al.* Prolonged low-dose methylprednisolone in patients with severe COVID-19 pneumonia. *medRxiv* 2020; : 2020.06.17.20134031.
- 183 Wei W, Ortwine JK, Mang NS, Joseph C, Hall BC, Prokesch BC. Limited Role for Antibiotics in COVID-19: Scarce Evidence of Bacterial Coinfection. *medRxiv* 2020; : 2020.06.16.20133181.
- 184 Zuo Y, Estes SK, Gandhi AA, *et al.* Prothrombotic antiphospholipid antibodies in COVID-19. *medRxiv* 2020; : 2020.06.15.20131607.
- 185 Killerby ME. Characteristics Associated with Hospitalization Among Patients with COVID-19 — Metropolitan Atlanta, Georgia, March–April 2020. *MMWR Morb Mortal Wkly Rep* 2020; **69**. DOI:10.15585/mmwr.mm6925e1.

- 186 Petrilli CM, Jones SA, Yang J, *et al.* Factors associated with hospital admission and critical illness among 5279 people with coronavirus disease 2019 in New York City: prospective cohort study. *BMJ* 2020; **369**. DOI:10.1136/bmj.m1966.
- 187 Magagnoli J, Narendran S, Pereira F, *et al.* Outcomes of Hydroxychloroquine Usage in United States Veterans Hospitalized with COVID-19. *Med* 2020; : S2666634020300064.
- 188 Bello-Chavolla OY, Bahena-López JP, Antonio-Villa NE, *et al.* Predicting Mortality Due to SARS-CoV-2: A Mechanistic Score Relating Obesity and Diabetes to COVID-19 Outcomes in Mexico. *J Clin Endocrinol Metab* 2020; **105**. DOI:10.1210/clinem/dgaa346.
- 189 Zuo Y, Zuo M, Yalavarthi S, *et al.* Neutrophil extracellular traps and thrombosis in COVID-19. *medRxiv* 2020; : 2020.04.30.20086736.
- 190 Sigel K, Swartz T, Golden E, *et al.* Covid-19 and People with HIV Infection: Outcomes for Hospitalized Patients in New York City. *Clin Infect Dis* DOI:10.1093/cid/ciaa880.
- 191 Nguyen AB, Upadhyay GA, Chung B, *et al.* Outcomes and Cardiovascular Comorbidities in a Predominantly African-American Population with COVID-19. *medRxiv* 2020; : 2020.06.28.20141929.
- 192 Melo AC de, Thuler LCS, Silva JL da, *et al.* Cancer inpatient with COVID-19: a report from the Brazilian National Cancer Institute. *medRxiv* 2020; : 2020.06.27.20141499.
- 193 Auvinen R, Nohynek H, Syrjänen R, *et al.* Comparison of the clinical characteristics and outcomes of hospitalized adult COVID-19 and influenza patients: a prospective observational study. *medRxiv* 2020; : 2020.06.29.20140632.
- 194 Souza FSH, Hojo-Souza NS, Santos EB, Silva CM, Guidoni DL. Predicting the disease outcome in COVID-19 positive patients through Machine Learning: a retrospective cohort study with Brazilian data. *medRxiv* 2020; : 2020.06.26.20140764.
- 195 Mendy A, Apewokin S, Wells AA, Morrow AL. Factors Associated with Hospitalization and Disease Severity in a Racially and Ethnically Diverse Population of COVID-19 Patients. *medRxiv* 2020; : 2020.06.25.20137323.
- 196 Pongpirul WA, Wiboonchutikul S, Charoenpong L, *et al.* Clinical course and potential predicting factors of pneumonia of adult patients with coronavirus disease 2019 (COVID-19): A retrospective observational analysis of 193 confirmed cases in Thailand. *medRxiv* 2020; : 2020.06.24.20139642.
- 197 Jin C, Gu J, Yuan Y, *et al.* Treatment of Six COVID-19 Patients with Convalescent Plasma. *medRxiv* 2020; : 2020.05.21.20109512.
- 198 Favara DM, Cooke A, Doffinger R, *et al.* First results from the UK COVID-19 Serology in Oncology Staff Study (CSOS). *medRxiv* 2020; : 2020.06.22.20136838.
- 199 Fisman D, Greer AL, Tuite A. Derivation and Validation of Clinical Prediction Rule for COVID-19 Mortality in Ontario, Canada. *medRxiv* 2020; : 2020.06.21.20136929.

- 200 Madariaga MLL, Guthmiller J, Schrantz S, *et al.* Clinical predictors of donor antibody titer and correlation with recipient antibody response in a COVID-19 convalescent plasma clinical trial. *medRxiv* 2020; : 2020.06.21.20132944.
- 201 Senkal N. Chronic ACE Inhibitor use is Associated with Decreased Odds of Severe Disease in Patients with COVID-19. *Anatol J Cardiol* 2020. DOI:10.14744/AnatolJCardiol.2020.57431.
- 202 Mohamud AY, Griffith B, Rehman M, *et al.* Intraluminal Carotid Artery Thrombus in COVID-19: Another Danger of Cytokine Storm? *Am J Neuroradiol* 2020; published online July 2. DOI:10.3174/ajnr.A6674.
- 203 Magleby R, Westblade LF, Trzebucki A, *et al.* Impact of SARS-CoV-2 Viral Load on Risk of Intubation and Mortality Among Hospitalized Patients with Coronavirus Disease 2019. *Clin Infect Dis* DOI:10.1093/cid/ciaa851.
- 204 Kimmig LM, Wu D, Gold M, *et al.* IL6 inhibition in critically ill COVID-19 patients is associated with increased secondary infections. *medRxiv* 2020; : 2020.05.15.20103531.
- 205 Bello-Chavolla OY, Antonio-Villa NE, Vargas-Vázquez A, Fermín-Martínez CA, Márquez-Salinas A, Bahena-López JP. Profiling pre-symptomatic and asymptomatic cases with confirmed SARS-CoV-2 infection in Mexico City. *medRxiv* 2020; : 2020.07.02.20145516.
- 206 Zacharioudakis IM, Prasad PJ, Zervou FN, *et al.* Association of SARS-CoV-2 Genomic Load with COVID-19 Patient Outcomes. *medRxiv* 2020; : 2020.07.02.20145151.
- 207 Antonio-Villa NE, Bello-Chavolla OY, Vargas-Vazquez A, Fermin-Martinez CA, Marquez-Salinas A, Bahena-Lopez JP. Health-care workers with COVID-19 living in Mexico City: clinical characterization and related outcomes. *medRxiv* 2020; : 2020.07.02.20145169.
- 208 Patel M, Chowdhury J, Mills N, *et al.* ROX Index Predicts Intubation in Patients with COVID-19 Pneumonia and Moderate to Severe Hypoxemic Respiratory Failure Receiving High Flow Nasal Therapy. *medRxiv* 2020; : 2020.06.30.20143867.
- 209 Merzon E, Tworowski D, Gorohovski A, *et al.* Low plasma 25(OH) vitamin D3 level is associated with increased risk of COVID-19 infection: an Israeli population-based study. *medRxiv* 2020; : 2020.07.01.20144329.
- 210 Fan X, Yin C, Wang J, *et al.* Pre-diagnostic circulating concentrations of insulin-like growth factor-1 and risk of COVID-19 mortality: results from UK Biobank. *medRxiv* 2020; : 2020.07.09.20149369.
- 211 Shi Z, Resurreccion WK, Wang C-H, *et al.* Association of Cancer with Risk and Mortality of COVID-19: Results from the UK Biobank. *medRxiv* 2020; : 2020.07.10.20151076.

- 212 Maucourant C, Filipovic I, Ponzetta A, *et al.* Natural killer cell activation related to clinical outcome of COVID-19. *medRxiv* 2020; : 2020.07.07.20148478.
- 213 Elmunzer BJ, Spitzer RL, Foster LD, *et al.* Digestive Manifestations in Patients Hospitalized with COVID-19. *medRxiv* 2020; : 2020.07.07.20143024.
- 214 Alizadehsani R, Sani ZA, Behjati M, *et al.* Risk Factors Prediction, Clinical Outcomes, and Mortality of COVID-19 Patients. *medRxiv* 2020; : 2020.07.07.20148569.
- 215 Xie Y, Chen S, Wang X, *et al.* Early Diagnosis and Clinical Significance of Acute Cardiac Injury - Under the Iceberg: A Retrospective Cohort Study of 619 Non-critically Ill Hospitalized COVID-19 Pneumonia Patients. *medRxiv* 2020; : 2020.07.06.20147256.
- 216 Fox TA, Troy-Barnes E, Kirkwood AA, *et al.* Clinical outcomes and risk factors for severe COVID-19 infection in patients with haematological disorders receiving chemo- or immunotherapy. *Br J Haematol*; n/a. DOI:10.1111/bjh.17027.
- 217 Martinez-Resendez MF, Castilleja-Leal F, Torres-Quintanilla A, *et al.* Initial experience in Mexico with convalescent plasma in COVID-19 patients with severe respiratory failure, a retrospective case series. *medRxiv* 2020; : 2020.07.14.20144469.
- 218 Hoertel N, Rico MS, Vernet R, *et al.* Observational Study of Haloperidol in Hospitalized Patients with Covid-19. *medRxiv* 2020; : 2020.07.15.20150490.
- 219 McGrail DE, Edwards D. COVID-19 Case Series at UnityPoint Health St. Lukes Hospital in Cedar Rapids, IA. *medRxiv* 2020; : 2020.07.17.20156521.
- 220 Pandolfi L, Fossali T, Frangipane V, *et al.* Broncho-alveolar inflammation in COVID-19 patients: a correlation with clinical outcome. *medRxiv* 2020; : 2020.07.17.20155978.
- 221 Kazuyoshi Kurashima, Naho Kagiya, Takashi Ishiguro, *et al.* IgG antibody seroconversion and the clinical progression of COVID-19 pneumonia: A retrospective, cohort study | medRxiv. <https://doi.org/10.1101/2020.07.16.20154088> (accessed Aug 25, 2020).
- 222 Zhan Z, Yang X, Du H, *et al.* Early Improvement of Acute Respiratory Distress Syndrome in Patients with COVID-19: Insights from the Data of ICU Patients in Chongqing, China. *medRxiv* 2020; : 2020.07.15.20154047.
- 223 Omrani AS, Almaslamani MA, Daghfal J, *et al.* The First Consecutive 5000 Patients with Coronavirus Disease 2019 from Qatar; a Nation-wide Cohort Study. *medRxiv* 2020; : 2020.07.15.20154690.
- 224 Gupta R, Agrawal R, Bukhari Z, *et al.* Higher Comorbidities and Early Death is Characteristic of Hospitalized African-American Patients with COVID-19. *medRxiv* 2020; : 2020.07.15.20154906.
- 225 Shi H, Zuo Y, Yalavarthi S, *et al.* Neutrophil calprotectin identifies severe pulmonary disease in COVID-19. *medRxiv* 2020; : 2020.05.06.20093070.

- 226 Hussein MH, Toraih EA, Attia AS, *et al.* Asthma in COVID-19: An extra chain fitting around the neck? *medRxiv* 2020; : 2020.07.13.20153130.
- 227 Bian H, Zheng Z-H, Wei D, *et al.* Meplazumab treats COVID-19 pneumonia: an open-labelled, concurrent controlled add-on clinical trial. *medRxiv* 2020; : 2020.03.21.20040691.
- 228 Eiros R, Barreiro-Perez M, Martin-Garcia A, *et al.* Pericarditis and myocarditis long after SARS-CoV-2 infection: a cross-sectional descriptive study in health-care workers. *medRxiv* 2020; : 2020.07.12.20151316.
- 229 Marcos M, Belhassen-Garcia M, Puente AS-, *et al.* Development of a severity of disease score and classification model by machine learning for hospitalized COVID-19 patients. *medRxiv* 2020; : 2020.07.13.20150177.
- 230 Hoertel N, Rico MS, Vernet R, *et al.* Association between SSRI Antidepressant Use and Reduced Risk of Intubation or Death in Hospitalized Patients with Coronavirus Disease 2019: a Multicenter Retrospective Observational Study. *medRxiv* 2020; : 2020.07.09.20143339.
- 231 Soares R de CM, Mattos LR, Raposo LM. Risk Factors for Hospitalization and Mortality due to COVID-19 in Espírito Santo State, Brazil. 2020; : tpm200483.
- 232 Zobairy H, Shamsoddin E, Rasouli MA, *et al.* Association of olfactory dysfunction with hospitalization for COVID-19: a multicenter study in Kurdistan. *medRxiv* 2020; : 2020.07.26.20158550.
- 233 Altamimi H, Alahmad Y, Khazal F, *et al.* The Outcome of COVID-19 Patients with Acute Myocardial Infarction. *medRxiv* 2020; : 2020.07.21.20156349.
- 234 Thompson JV, Meghani N, Powell BM, *et al.* Patient characteristics and predictors of mortality in 470 adults admitted to a district general hospital in England with Covid-19. *medRxiv* 2020; : 2020.07.21.20153650.
- 235 Reiter T, Pajenda S, Wagner L, *et al.* Covid-19 serology in nephrology health care workers. *medRxiv* 2020; : 2020.07.21.20136218.
- 236 Motta JK, Ogunnaike RO, Shah R, *et al.* Clinical Outcomes With the Use of Prophylactic Versus Therapeutic Anticoagulation in COVID-19. *medRxiv* 2020; : 2020.07.20.20147769.
- 237 Santos C, Rhee Y, Hollinger E, *et al.* Comparative Incidence and Outcomes of COVID-19 in Kidney or Kidney-Pancreas Transplant Recipients Versus Kidney or Kidney-Pancreas Waitlisted Patients: A Pilot Study. *medRxiv* 2020; : 2020.07.20.20157990.
- 238 Schneeweiss MC, Leonard S, Weckstein A, Schneeweiss S, Rassen J. Renin-Angiotensin-Aldosterone-System inhibitor use in patients with COVID-19 infection and prevention of serious events: a cohort study in commercially insured patients in the US. *medRxiv* 2020; : 2020.07.22.20159855.

- 239 Concha-Mejia A, Rincon-Sanchez RA. CCOFEE-GI Study: Colombian COVID19 First Experience in Gastroenterology. Characterization of digestive manifestations in patients diagnosed with COVID-19 at a highly complex institution in Bogota D.C., Colombia. *medRxiv* 2020; : 2020.07.24.20161604.
- 240 Izquierdo JL, Almonacid C, Gonzalez Y, *et al.* The impact of COVID-19 on patients with asthma. *medRxiv* 2020; : 2020.07.24.20161596.
- 241 Bernaola N, Mena R, Bernaola A, *et al.* Observational Study of the Efficiency of Treatments in Patients Hospitalized with Covid-19 in Madrid. *medRxiv* 2020; : 2020.07.17.20155960.
- 242 Qi D, Yan X, Tang X, *et al.* Epidemiological and clinical features of 2019-nCoV acute respiratory disease cases in Chongqing municipality, China: a retrospective, descriptive, multiple-center study. *medRxiv* 2020; : 2020.03.01.20029397.
- 243 Peters EJ, Collard D, Assen S van, *et al.* Outcomes of Persons With COVID-19 in Hospitals With and Without Standard Treatment With (Hydroxy)chloroquine. *medRxiv* 2020; : 2020.08.14.20173369.
- 244 Ouyang J, Shan X, Wang X, *et al.* Clinical characteristics of COVID-19 and the model for predicting the occurrence of critically ill patients: a retrospective cohort study. *medRxiv* 2020; : 2020.08.13.20173799.
- 245 Valenzuela O, Ibanez SE, Poli M, *et al.* First report of tocilizumab use in a cohort of Latin American patients hospitalized for severe COVID-19 pneumonia. *medRxiv* 2020; : 2020.08.12.20173104.
- 246 Monteiro ACC, Suri R, Emeruwa IO, *et al.* Obesity and Smoking as Risk Factors for Invasive Mechanical Ventilation in COVID-19: a Retrospective, Observational Cohort Study. *medRxiv* 2020; : 2020.08.12.20173849.
- 247 Philipose Z, Smati N, Wong CSJ, Aspey K, Mendall MA. Obesity, old age and frailty are the true risk factors for COVID-19 mortality and not chronic disease or ethnicity in Croydon. *medRxiv* 2020; : 2020.08.12.20156257.
- 248 Weerahandi H, Hochman KA, Simon E, *et al.* Post-discharge health status and symptoms in patients with severe COVID-19. *medRxiv* 2020; : 2020.08.11.20172742.
- 249 Altibi AM, Bhargava P, Liaqat H, *et al.* Comparative Clinical Outcomes and Mortality in Prisoner and Non-Prisoner Populations Hospitalized with COVID-19: A Cohort from Michigan. *medRxiv* 2020; : 2020.08.08.20170787.
- 250 Izzi-Engbeaya C, Distaso W, Amin A, *et al.* Severe COVID-19 and Diabetes: A Retrospective Cohort Study from Three London Teaching Hospitals. *medRxiv* 2020; : 2020.08.07.20160275.
- 251 Rizzo S, Chawla D, Zalocusky K, *et al.* Descriptive epidemiology of 16,780 hospitalized COVID-19 patients in the United States. *medRxiv* 2020; : 2020.07.17.20156265.

- 252 Dashti HT, Bates D, Fiskio JM, Roche EC, Mora S, Demler O. Clinical Characteristics and Severity of COVID-19 Disease in Patients from Boston Area Hospitals. *medRxiv* 2020; : 2020.07.27.20163071.
- 253 Morshed MS, Mosabbir AA, Chowdhury P, Ashadullah SM, Hossain MS. Clinical manifestations of patients with Coronavirus Disease 2019 (COVID- 19) attending at hospitals in Bangladesh. *medRxiv* 2020; : 2020.07.30.20165100.
- 254 Jun T, Nirenberg S, Kovatch P, Huang K. Sex-specificity of mortality risk factors among hospitalized COVID-19 patients in New York City: prospective cohort study. *medRxiv* 2020; : 2020.07.29.20164640.
- 255 Higuchi T, Nishida T, Iwahashi H, *et al.* Early Clinical Factors Predicting the Development of Critical Disease in Japanese Patients with COVID-19: A Single-Center Retrospective, Observational Study. *medRxiv* 2020; : 2020.07.29.20159442.
- 256 Zhou K, Sun Y, Li L, *et al.* Eleven Routine Clinical Features Predict COVID-19 Severity. *medRxiv* 2020; : 2020.07.28.20163022.
- 257 Salerno S, Zhao Z, Sankar SP, *et al.* Understanding the patterns of repeated testing for COVID-19: Association with patient characteristics and outcomes. *medRxiv* 2020; : 2020.07.26.20162453.
- 258 Kumar A, Prasad G, Srivastav S, Gautam VK, Sharma N. A Retrospective Study on Efficacy and Safety of Guduchi Ghan Vati for Covid-19 Asymptomatic Patients. *medRxiv* 2020; : 2020.07.23.20160424.
- 259 Hao S-R, Zhang S-Y, Lian J-S, *et al.* Liver Enzyme Elevation in Coronavirus Disease 2019: A Multicenter, Retrospective, Cross-Sectional Study. *Am J Gastroenterol* 2020; published online June 1. DOI:10.14309/ajg.0000000000000717.
- 260 Iversen K, Bundgaard H, Hasselbalch RB, *et al.* Risk of COVID-19 in health-care workers in Denmark: an observational cohort study. *Lancet Infect Dis* 2020; **0**. DOI:10.1016/S1473-3099(20)30589-2.
- 261 Hippisley-Cox J, Young D, Coupland C, *et al.* Risk of severe COVID-19 disease with ACE inhibitors and angiotensin receptor blockers: cohort study including 8.3 million people. *Heart* 2020; : heartjnl-2020-317393.
- 262 Fillmore NR, La J, Szalat RE, *et al.* Prevalence and outcome of Covid-19 infection in cancer patients: a national VA study. *medRxiv* 2020; : 2020.08.21.20177923.
- 263 Rashid M, Wu J, Timmis A, *et al.* Clinical Characteristics and Outcomes of COVID-19 Positive Acute Coronary Syndrome Patients; a multisource Electronic Healthcare Records Study from England. *medRxiv* 2020; : 2020.08.20.20175091.
- 264 Pan A, Khan O, Meeks J, *et al.* Disparities in COVID-19 Hospitalizations and Mortality among Black and Hispanic Patients: Cross-Sectional Analysis from the Greater Houston Metropolitan Area. *medRxiv* 2020; : 2020.08.19.20177956.

- 265 Alkurt G, Murt A, Aydin Z, *et al.* Seroprevalence of Coronavirus Disease 2019 (COVID-19) Among Health Care Workers from Three Pandemic Hospitals of Turkey. *medRxiv* 2020; : 2020.08.19.20178095.
- 266 Zhao Z, Chen A, Hou W, *et al.* Prediction model and risk scores of ICU admission and mortality in COVID-19. *PLOS ONE* 2020; **15**: e0236618.
- 267 Holman N, Knighton P, Kar P, *et al.* Risk factors for COVID-19-related mortality in people with type 1 and type 2 diabetes in England: a population-based cohort study. *Lancet Diabetes Endocrinol* 2020; **0**. DOI:10.1016/S2213-8587(20)30271-0.
- 268 Qu J, Chang LK, Tang X, *et al.* Clinical characteristics of COVID-19 and its comparison with influenza pneumonia. *Acta Clin Belg* 2020; **0**: 1–9.
- 269 Chand S, Kapoor S, Orsi D, *et al.* COVID-19-Associated Critical Illness—Report of the First 300 Patients Admitted to Intensive Care Units at a New York City Medical Center: *J Intensive Care Med* 2020; published online Aug 19. DOI:10.1177/0885066620946692.




Figure 1. PRISMA flow diagram of included studies.

| Ref. | Lead author | Date published | Country | Sample size | Study setting | Median (IQR) | Female % | Current smoker % | Former smokers % | Current/former smokers % | Never smokers % | Never/unknown smokers % | Missing % | Study quality |
|------|-------------|----------------|---------|-------------|---------------|--------------|----------|------------------|------------------|--------------------------|-----------------|-------------------------|-----------|---------------|
| 1 | Guan, Ni | 2020-02-28 | China | 1,099 | Hospital | 47 (35-58) | 41.9 | 12.5 | 1.9 | - | 84.3 | - | 1.27 | fair |
| 50 | Guan, Liang | 2020-03-26 | China | 1,590 | Hospital | 49 (33-64) | 42.7 | - | - | 7.0 | 93.0 | - | 0.00 | poor |
| 51 | Lian | 2020-03-25 | China | 788 | Hospital | NA | 38.5 | 6.9 | - | - | - | - | 93.15 | poor |
| 52 | Jin | 2020-03-24 | China | 651 | Hospital | 46 (32-60) | 49.2 | 6.3 | - | - | - | - | 93.70 | poor |
| 53 | Chen | 2020-03-26 | China | 548 | Hospital | 62 (44-70) | 37.6 | 4.4 | 2.6 | - | - | - | 93.07 | poor |
| 54 | Zhou, Yu | 2020-03-11 | China | 191 | Hospital | 56 (46-67) | 38.0 | 5.8 | - | - | - | - | 94.24 | poor |
| 55 | Mo | 2020-03-16 | China | 155 | Hospital | 54 (53-66) | 44.5 | 3.9 | - | - | - | - | 96.13 | poor |
| 56 | Zhang, Dong | 2020-02-19 | China | 140 | Hospital | 57^ (25-87) | 46.3 | 1.4 | 5.0 | - | - | - | 93.57 | poor |
| 57 | Wan | 2020-03-21 | China | 135 | Hospital | 47 (36-55) | 46.7 | 6.7 | - | - | - | - | 93.33 | poor |
| 58 | Liu, Tao | 2020-02-28 | China | 78 | Hospital | 38 (33-57) | 50.0 | - | - | 6.4 | - | - | 93.59 | poor |
| 59 | Huang, Wang | 2020-01-24 | China | 41 | Hospital | 49 (41-58) | 27.0 | 7.3 | - | - | - | - | 92.68 | poor |
| 60 | Zhang, Cai | 2020-03-20 | China | 645 | Hospital | NA | 49.1 | 6.4 | - | - | - | - | 93.64 | poor |
| 61 | Guo | 2020-03-27 | China | 187 | Hospital | 59 (45-73) | 51.3 | 9.6 | - | - | - | - | 90.37 | poor |
| 62 | Liu, Ming | 2020-03-12 | China | 41 | Hospital | 39 (30-48) | 58.5 | 9.8 | - | - | - | - | 90.24 | poor |
| 63 | Huang, Yang | 2020-03-05 | China | 36 | Hospital | 69 (60-78) | 30.6 | - | - | 11.1 | - | - | 88.89 | poor |
| 64 | Xu | 2020-03-08 | China | 53 | Hospital | NA | 47.2 | 11.3 | - | - | - | - | 88.68 | poor |
| 65 | Li | 2020-02-12 | China | 17 | Hospital | 45 (33-57) | 47.1 | 17.6 | - | - | - | - | 82.35 | poor |

| | | | | | | | | | | | | | | |
|----|---------------|------------|-------------|-------|------------------------|-------------|------|------|------|------|------|------|-------|------|
| 31 | Rentsch | 2020-04-14 | USA | 3,528 | Community and Hospital | 66 (60-70) | 4.6 | 27.2 | 30.6 | - | 36.9 | - | 5.30 | fair |
| 66 | Hu | 2020-03-25 | China | 323 | Hospital | 61^ (23-91) | 48.6 | - | - | 11.8 | - | - | 88.24 | poor |
| 67 | Wang, Pan | 2020-03-24 | China | 125 | Hospital | 41 (26-66) | 43.2 | - | - | 12.8 | - | - | 87.20 | poor |
| 68 | Chow (US CDC) | 2020-03-31 | USA | 7,162 | Community and Hospital | NA | - | 1.3 | 2.3 | - | - | - | 96.36 | poor |
| 69 | Dong, Cao | 2020-03-20 | China | 9 | Hospital | 44 (30-46) | 66.7 | 11.1 | - | - | - | - | 88.89 | poor |
| 70 | Kim | 2020-04-01 | South Korea | 28 | Hospital | 43 (30-56) | 46.4 | 17.9 | - | - | - | - | 82.14 | poor |
| 71 | Shi, Yu | 2020-03-18 | China | 487 | Hospital | 46 (27-65) | 46.8 | - | - | 8.2 | - | - | 91.79 | poor |
| 72 | Yang, Yu | 2020-02-24 | China | 52 | Hospital | 60 (47-73) | 37.0 | 3.8 | - | - | - | - | 96.15 | poor |
| 73 | Argenziano | 2020-05-29 | USA | 1,000 | Hospital | 63 (50-75) | 40.4 | 4.9 | 17.9 | - | 77.2 | - | 0.00 | fair |
| 74 | Solis | 2020-04-25 | Mexico | 650 | Hospital | 46 (NA) | 42.1 | 9.4 | - | - | - | - | 90.62 | poor |
| 75 | Richardson | 2020-04-22 | USA | 5,700 | Hospital | 63 (52-75) | 39.7 | - | - | 9.8 | 52.8 | - | 37.42 | poor |
| 76 | Fontanet | 2020-04-23 | France | 661 | Community and Hospital | 37 (16-47) | 62.0 | 10.4 | - | - | - | 89.6 | 0.00 | poor |
| 77 | Zheng, Gao | 2020-04-19 | China | 66 | Hospital | 47^ (NA) | 25.8 | 12.1 | - | - | - | - | 87.88 | poor |
| 78 | Liao, Feng | 2020-04-24 | China | 1,848 | Hospital | 55 (48-61) | 54.7 | - | - | 0.4 | - | - | 99.57 | poor |
| 79 | Gil-Agudo | 2020-04-24 | Spain | 7 | Hospital | 68 (34-75) | 28.6 | - | - | 42.9 | 57.1 | - | 0.00 | poor |
| 80 | Shi, Ren | 2020-04-23 | China | 134 | Hospital | 46 (34-58) | 51.5 | - | - | 10.4 | - | - | 89.55 | poor |
| 81 | Hadjadj | 2020-04-23 | France | 50 | Hospital | 55 (50-63) | 22.0 | 2.0 | 18.0 | - | 80.0 | - | 0.00 | fair |
| 82 | Gold (US CDC) | 2020-04-20 | USA | 305 | Hospital | NA | 50.5 | 5.2 | - | - | - | - | 94.75 | poor |
| 83 | Yu, Cai | 2020-04-27 | China | 95 | Hospital | NA | 44.2 | 8.4 | - | - | - | - | 91.58 | poor |

| | | | | | | | | | | | | | | |
|-----|------------------------------|------------|--------|------------|------------------------|------------------|------|------|------|------|------|------|-------|------|
| 84 | Zheng, Xiong | 2020-04-30 | China | 73 | Hospital | 43^ (NA) | 45.2 | - | - | 11.0 | 89.0 | - | 0.00 | poor |
| 85 | de la Rica | 2020-05-11 | Spain | 48 | Hospital | 66^ (33-88) | 33.0 | - | - | 20.8 | - | - | 79.17 | poor |
| 86 | Yin, Yang | 2020-05-10 | China | 106 | Hospital | 73 (61-85) | 39.6 | - | - | 17.0 | - | - | 83.02 | poor |
| 87 | Shi, Zuo | 2020-05-17 | USA | 172 | Hospital | 63^ (44-82) | 44.0 | - | - | 26.2 | - | - | 73.84 | poor |
| 88 | Cho | 2020-05-11 | UK | 322,341 | Community and Hospital | NA | 49.2 | 14.2 | 21.4 | - | 64.4 | - | 0.00 | fair |
| 89 | Allenbach | 2020-05-08 | France | 152 | Hospital | 77 (60-83) | 31.1 | - | - | 6.6 | - | - | 93.42 | poor |
| 90 | Robilotti | 2020-05-08 | USA | 423 | Hospital | NA | 50.0 | 2.1 | 37.6 | - | 58.6 | - | 1.65 | fair |
| 91 | The Opensafely Collaborative | 2020-07-01 | UK | 17,278,392 | Community and Hospital | NA | 50.1 | 17.0 | 32.9 | - | 45.9 | - | 4.17 | fair |
| 92 | Borobia | 2020-05-06 | Spain | 2,226 | Hospital | 61 (46-78) | 52.0 | 7.1 | - | - | - | - | 92.95 | poor |
| 93 | Giacomelli | 2020-05-06 | Italy | 233 | Hospital | 61 (50-72) | 31.9 | - | - | 30.0 | 70.0 | - | 0.00 | poor |
| 94 | Shah | 2020-05-06 | USA | 316 | Hospital | 63 (43-72) | 48.1 | 16.5 | 17.7 | - | 42.1 | - | 23.73 | poor |
| 95 | Kolin | 2020-05-05 | UK | 502,536 | Community and Hospital | 56.5 (48-64) | 54.4 | 10.5 | 34.4 | - | 54.4 | - | 0.59 | fair |
| 96 | Lubetzky | 2020-05-08 | USA | 54 | Hospital | 57 (29-83) | 62.0 | - | - | 22.2 | - | - | 77.78 | poor |
| 97 | Goyal | 2020-04-17 | USA | 393 | Hospital | 62.2 (49-74) | 39.3 | 5.1 | - | - | - | - | 94.91 | poor |
| 98 | Feng | 2020-04-10 | China | 476 | Hospital | 53 (40-64) | 43.1 | 9.2 | - | - | - | - | 90.76 | poor |
| 99 | Yao | 2020-04-24 | China | 108 | Hospital | 52 (37-58) | 60.2 | 3.7 | - | - | - | - | 96.30 | poor |
| 100 | Sami | 2020-05-19 | Iran | 490 | Hospital | 56.6 (41-71) | 39.0 | 14.1 | - | - | - | 85.9 | 0.00 | poor |
| 101 | Almazeedi | 2020-05-15 | Kuwait | 1,096 | Hospital | 41 (25-57) | 19.0 | 4.0 | - | - | - | 96.0 | 0.00 | poor |
| 102 | Carillo-Vega | 2020-05-14 | Mexico | 10,544 | Community and Hospital | 46.5^ (30-62) | 42.3 | 8.9 | - | - | - | - | 91.12 | poor |

| | | | | | | | | | | | | | | |
|-----|-----------------|------------|-------------|---------|------------------------|-----------------|------|------|------|------|------|------|-------|------|
| 103 | Yanover | 2020-05-13 | Israel | 4,353 | Community and Hospital | 35 (22-54) | 44.5 | 11.8 | 3.0 | - | 85.2 | - | 0.00 | fair |
| 104 | Hamer | 2020-05-13 | UK | 387,109 | Hospital | 56.2 (48-64) | 55.1 | 9.7 | 34.8 | - | 55.5 | - | 0.00 | fair |
| 105 | Regina | 2020-05-14 | Switzerland | 200 | Hospital | 70 (55-81) | 40.0 | 4.5 | - | - | - | - | 95.50 | poor |
| 106 | de Lusignan | 2020-05-15 | UK | 3,802 | Community and Hospital | 58 (34-73) | 57.6 | 10.9 | 46.1 | - | 29.6 | - | 13.44 | fair |
| 107 | Targher | 2020-05-13 | China | 339 | Hospital | 48.4^ (NA) | 52.8 | 8.3 | - | - | - | - | 91.74 | poor |
| 108 | Valenti | 2020-05-18 | Italy | 789 | Community | 40.7^ (NA) | 35.0 | 25.9 | - | - | - | - | 74.14 | poor |
| 109 | Feuth | 2020-05-18 | Finland | 28 | Hospital | 56 (47-72) | 46.0 | 10.7 | 28.6 | - | 60.7 | - | 0.00 | fair |
| 110 | Ge | 2020-05-18 | China | 51 | Hospital | 70 (58-79) | 27.5 | 13.7 | - | - | - | - | 86.27 | poor |
| 111 | Parrotta | 2020-05-18 | USA | 76 | Community and Hospital | 44.9 (13-71) | 61.8 | 2.6 | 26.3 | - | 68.4 | - | 2.63 | fair |
| 112 | Shekhar | 2020-05-18 | USA | 50 | Hospital | 55.5 (20-85) | 54.0 | 48.0 | - | - | - | - | 52.00 | poor |
| 113 | Mejia-Vilet | 2020-05-16 | Mexico | 329 | Hospital | 49 (41-60) | 36.0 | - | - | 7.0 | - | - | 93.01 | poor |
| 114 | Chen, Jiang | 2020-05-16 | China | 135 | Hospital | NA | 42.2 | - | - | 9.6 | - | - | 90.37 | poor |
| 115 | Li, Chen | 2020-05-16 | China | 1,008 | Hospital | 55 (44-65) | 43.6 | 5.7 | - | - | - | - | 94.35 | poor |
| 27 | Rimland | 2020-05-19 | USA | 11 | Hospital | 59 (48-65) | 18.2 | 9.1 | - | - | - | - | 81.82 | poor |
| 116 | Palaiodimos | 2020-05-15 | USA | 200 | Hospital | 64 (50-73.5) | 51.0 | - | - | 32.5 | 67.5 | - | 0.00 | poor |
| 117 | Ip | 2020-05-25 | USA | 2,512 | Hospital | 64 (52-76) | 37.6 | 3.1 | 17.8 | - | 64.5 | - | 14.61 | fair |
| 118 | Heili-Frades | 2020-05-25 | Spain | 4,712 | Hospital | 62 (47-77) | 50.5 | 4.9 | 17.4 | - | - | 66.5 | 11.16 | poor |
| 119 | Vaquero-Roncero | 2020-05-24 | Spain | 146 | Hospital | 66^ (59-72) | 32.2 | - | - | 6.8 | - | - | 93.15 | poor |
| 120 | Kim, Garg | 2020-05-22 | USA | 2,491 | Hospital | 62 (50-75) | 46.8 | 6.0 | 25.8 | - | - | 68.1 | 0.08 | poor |

| | | | | | | | | | | | | | | |
|-----|---------------|------------|--------------|---------|------------------------|-------------------|------|------|------|------|------|------|-------|------|
| 121 | Wu | 2020-05-21 | Italy | 174 | Hospital | 61.2^ (50-71) | 30.5 | - | - | 33.3 | - | - | 66.67 | poor |
| 122 | Shi, Zhao | 2020-05-20 | China | 101 | Hospital | 71 (59-80) | 40.6 | - | - | 5.0 | - | - | 95.05 | poor |
| 123 | Al-Hindawi | 2020-05-20 | UK | 31 | Hospital | 61 (NA) | 12.9 | 3.2 | 71.0 | - | 25.8 | - | 0.00 | fair |
| 124 | Basse | 2020-05-19 | France | 141 | Hospital | 62 (52-72) | 72.0 | 17.7 | - | - | - | - | 82.27 | poor |
| 125 | Freites | 2020-05-19 | Spain | 123 | Hospital | 59.88^ (44-74) | 69.9 | 3.3 | - | - | - | - | 96.75 | poor |
| 126 | Alshami | 2020-05-19 | Saudi Arabia | 128 | Quarantine Centre | 39.6^ (24-55) | 53.9 | 15.6 | 2.3 | - | - | - | 82.03 | poor |
| 127 | Berumen | 2020-05-26 | Mexico | 102,875 | Hospital | NA | 49.1 | - | - | 9.6 | - | 90.4 | 0.00 | poor |
| 128 | Gianfrancesco | 2020-05-29 | Multiple | 600 | Community and Hospital | 56 (45-67) | 71.0 | - | - | 21.5 | 64.8 | - | 13.67 | poor |
| 129 | Li, Long | 2020-05-28 | China | 145 | Not Stated | 49^ (13-80) | 61.0 | - | - | 5.5 | - | - | 94.48 | poor |
| 130 | Batty | 2020-06-17 | UK | 908 | Hospital | 57.27^ (48-66) | 44.3 | 11.2 | - | - | - | - | 88.77 | poor |
| 131 | Israel | 2020-06-01 | Israel | 24,906 | Community and Hospital | 40 (27-59) | 48.7 | 16.8 | 12.7 | - | 70.5 | - | 0.00 | fair |
| 132 | del Valle | 2020-05-30 | USA | 1,484 | Hospital | 62 (52-72) | 40.6 | 5.5 | 23.3 | - | - | - | 71.16 | poor |
| 133 | Chaudhry | 2020-05-29 | USA | 40 | Community and Hospital | 52 (45.5-61) | 60.0 | - | - | 15.0 | - | - | 85.00 | poor |
| 134 | Louis | 2020-05-28 | USA | 22 | Hospital | 66.5^ (55-77) | 36.4 | - | - | 45.5 | - | - | 54.55 | poor |
| 135 | Soto-Mota | 2020-06-05 | Mexico | 400 | Hospital | NA | 30.0 | - | - | 12.0 | - | - | 88.00 | poor |
| 136 | Garibaldi | 2020-05-26 | USA | 832 | Hospital | 63 (49-75) | 47.0 | 5.5 | 22.6 | - | - | - | 71.88 | poor |
| 137 | Docherty | 2020-05-22 | Multiple | 20,133 | Hospital | 72.9 (58-82) | 40.0 | 4.2 | 21.7 | - | 44.5 | - | 29.55 | poor |
| 138 | Boulware | 2020-06-03 | Multiple | 821 | Community | 40 (33-50) | 51.6 | 3.3 | - | - | - | - | 96.71 | poor |
| 139 | Kuderer | 2020-05-28 | Multiple | 928 | Community and Hospital | 66 (57-76) | 50.0 | 4.6 | 35.1 | - | 50.5 | - | 9.70 | fair |

| | | | | | | | | | | | | | | |
|-----|-------------------|------------|----------|---------|------------------------|-------------------|-------|------|------|------|------|------|-------|------|
| 140 | Romao | 2020-06-08 | Portugal | 34 | Community | 41^ (26-66) | 67.7 | - | - | 26.5 | - | - | 73.53 | poor |
| 141 | Giannouchos | 2020-06-07 | Mexico | 236,439 | Community and Hospital | 42.5^ (25-59) | 49.1 | 9.1 | - | - | - | 90.9 | 0.00 | poor |
| 142 | Ramlall | 2020-06-06 | USA | 11,116 | Community and Hospital | 52 (34.7-69.5) | 55.2 | - | - | 26.8 | 73.2 | - | 0.00 | poor |
| 143 | Wang, Oekelen | 2020-06-05 | USA | 58 | Community and Hospital | 67 (NA) | 48.0 | - | - | 36.2 | - | - | 63.79 | poor |
| 144 | Perrone | 2020-06-05 | Italy | 1,189 | Hospital | NA | 21.2 | - | - | 21.9 | - | - | 78.13 | poor |
| 145 | Sharma | 2020-06-05 | India | 501 | Hospital | 35.1^ (18-51) | 36.0 | - | - | 4.2 | - | - | 95.81 | poor |
| 146 | Eugen-Olsen | 2020-06-02 | Denmark | 407 | Hospital | 64 (47-77) | 57.7 | 20.6 | 36.9 | - | 39.6 | - | 2.95 | fair |
| 147 | Martinez-Portilla | 2020-06-02 | Mexico | 224 | Community and Hospital | 29 (26-33) | 100.0 | - | - | 3.1 | - | - | 96.88 | poor |
| 148 | Raisi-Estabragh | 2020-06-02 | UK | 4,510 | Hospital | NA | 48.8 | - | - | 51.8 | - | - | 48.20 | poor |
| 149 | Luo | 2020-06-02 | China | 625 | Hospital | 46 (NA) | 47.7 | 3.0 | - | - | - | - | 96.96 | poor |
| 150 | Houlihan | 2020-06-09 | UK | 200 | Community | 34 (29-44) | 61.0 | 11.0 | 16.5 | - | 66.5 | - | 6.00 | fair |
| 151 | Cen | 2020-06-08 | China | 1,007 | Hospital | 61 (49-68) | 51.0 | - | - | 8.7 | - | - | 91.26 | poor |
| 152 | Klang | 2020-05-23 | USA | 3,406 | Hospital | NA | 61.8 | - | - | 23.3 | - | - | 76.72 | poor |
| 153 | Maraschini | 2020-06-12 | Italy | 146 | Hospital | 32.5^ (27-38) | 100.0 | - | 9.6 | - | 80.8 | - | 9.59 | poor |
| 154 | Wang, Zhong | 2020-06-12 | USA | 7,592 | Community and Hospital | NA | 45.1 | 3.6 | 17.1 | - | 51.9 | - | 27.42 | poor |
| 155 | McQueenie | 2020-06-12 | UK | 428,199 | Community and Hospital | NA | 54.9 | - | - | 44.4 | 55.0 | - | 0.59 | poor |
| 26 | Miyara | 2020-06-12 | France | 479 | Community and Hospital | NA | 44.7 | 6.7 | 31.6 | - | 59.5 | - | 1.87 | fair |
| 156 | Apea | 2020-06-12 | UK | 1,737 | Hospital | 63.4^ (NA) | 30.4 | - | - | 10.0 | - | - | 90.04 | poor |

| | | | | | | | | | | | | | | |
|-----|-------------------|------------|----------|--------|------------------------|---------------|-------|------|------|------|------|------|-------|------|
| 157 | Woolford | 2020-06-11 | UK | 4,510 | Community and Hospital | 70.5 (NA) | 51.2 | 13.0 | 38.1 | - | 48.1 | - | 0.80 | fair |
| 158 | Hultcrantz | 2020-06-11 | USA | 127 | Community and Hospital | 68 (41-91) | 46.0 | - | - | 26.8 | 72.4 | - | 0.79 | poor |
| 159 | Rajter | 2020-06-10 | USA | 280 | Hospital | 59.6^ (41-77) | 45.5 | 5.7 | 10.7 | - | 74.6 | - | 8.93 | fair |
| 160 | Lan | 2020-06-09 | USA | 104 | Community | 49^ (34-63) | 47.1 | - | - | 24.0 | - | - | 75.96 | poor |
| 161 | Zeng | 2020-06-16 | China | 1,031 | Hospital | 60.3^ (46-74) | 47.8 | - | - | 10.2 | - | - | 89.82 | poor |
| 162 | Suleyman | 2020-06-16 | USA | 463 | Hospital | 57.5^ (40-74) | 55.9 | - | - | 34.6 | - | - | 65.44 | poor |
| 163 | Chen, Yu | 2020-06-16 | China | 1,859 | Hospital | 59 (45-68) | 50.0 | 2.4 | 3.6 | - | 94.0 | - | 0.00 | fair |
| 164 | Garassino | 2020-06-12 | Multiple | 200 | Community and Hospital | 68 (61.8-75) | 30.0 | 24.0 | 55.5 | - | 18.5 | - | 2.00 | fair |
| 165 | Hernandez-Garduno | 2020-06-11 | Mexico | 32,583 | Community and Hospital | 45 (34-56) | 48.7 | - | - | 11.0 | - | 88.8 | 0.15 | poor |
| 166 | Govind | 2020-06-20 | UK | 6,309 | Community and Hospital | 46.5^ (31-61) | 38.3 | 66.3 | 26.8 | - | 5.5 | - | 1.49 | fair |
| 167 | Siso-Almirall | 2020-06-20 | Spain | 322 | Community and Hospital | 56.7^ (38-74) | 50.0 | - | - | 25.2 | - | - | 74.84 | poor |
| 168 | Gu | 2020-06-18 | USA | 5,698 | Community and Hospital | 47^ (26-67) | 62.0 | 7.0 | 24.7 | - | 50.8 | - | 17.53 | fair |
| 169 | Kibler | 2020-06-16 | France | 702 | Community and Hospital | 82^ (75-88) | 56.0 | 3.7 | - | - | - | - | 96.30 | poor |
| 170 | Ikitimur | 2020-06-03 | Turkey | 81 | Hospital | 55^ (38-72) | 44.0 | - | - | 28.4 | - | - | 71.60 | poor |
| 171 | Sierpinski | 2020-06-03 | Poland | 1,942 | Community | 50 (NA) | 60.0 | 6.3 | - | - | - | 49.7 | 44.03 | poor |
| 172 | Zhou, He | 2020-06-10 | China | 238 | Hospital | 55.5 (35-67) | 57.0 | 2.9 | - | - | - | - | 97.06 | poor |
| 173 | Crovetto | 2020-06-19 | Spain | 874 | Community and Hospital | 33.7^ (28-38) | 100.0 | 1.1 | - | - | - | 13.2 | 85.70 | poor |
| 174 | Veras | 2020-06-09 | Brazil | 32 | Hospital | 58.9^ (40-77) | 47.0 | - | - | 25.0 | - | - | 75.00 | poor |
| 175 | Sterlin | 2020-06-11 | France | 135 | Hospital | 61 (50-72) | 41.0 | 3.7 | 38.5 | - | 57.8 | - | 0.00 | fair |

| | | | | | | | | | | | | | | |
|-----|-----------------|------------|---------|---------|------------------------|-------------------|------|------|------|------|------|---|-------|------|
| 176 | Rossi | 2020-06-09 | France | 246 | Hospital | 68^ (53-83) | 39.0 | - | - | 25.2 | - | - | 74.80 | poor |
| 177 | Duan | 2020-06-22 | China | 616 | Hospital | 64 (53-70) | 57.5 | 3.7 | - | - | - | - | 96.27 | poor |
| 178 | Martin-Jimenez | 2020-06-09 | Spain | 339 | Hospital | 81.6 (72-87) | 39.5 | - | - | 30.7 | - | - | 69.32 | poor |
| 179 | Elezkurtaj | 2020-06-17 | Germany | 26 | Hospital | 70 (61.8-78.3) | 34.6 | - | - | 19.2 | - | - | 80.77 | poor |
| 180 | Lenka | 2020-06-22 | USA | 32 | Hospital | 62.2^ (51-73) | 37.5 | - | - | 50.0 | - | - | 50.00 | poor |
| 181 | Olivares | 2020-06-16 | Chile | 21 | Hospital | 61^ (26-85) | 76.2 | - | - | 9.5 | - | - | 90.48 | poor |
| 182 | Salton | 2020-06-20 | Italy | 173 | Hospital | 64.4^ (NA) | 34.9 | - | - | 29.5 | - | - | 70.52 | poor |
| 183 | Wei | 2020-06-18 | USA | 147 | Hospital | 52^ (34-70) | 41.0 | 14.3 | - | - | - | - | 85.71 | poor |
| 184 | Zuo, Estes | 2020-06-17 | China | 172 | Hospital | 61^ (25-95) | 44.0 | - | - | 26.2 | - | - | 73.84 | poor |
| 185 | Killerby | 2020-06-17 | USA | 531 | Community and Hospital | 51.6 (38-62) | 57.1 | - | - | 17.1 | 71.4 | - | 11.49 | poor |
| 186 | Petrilli | 2020-05-22 | USA | 5,279 | Community and Hospital | 54 (38-66) | 51.5 | 5.5 | 17.1 | - | 61.9 | - | 15.55 | fair |
| 187 | Magagnoli | 2020-06-05 | USA | 807 | Hospital | 70 (60-75) | 4.3 | - | - | 15.9 | - | - | 84.14 | poor |
| 33 | Niedzwiedz | 2020-05-29 | UK | 392,116 | Community and Hospital | NA | 54.9 | 9.8 | 34.8 | - | 55.4 | - | 0.00 | fair |
| 188 | Bello-Chavolla | 2020-05-31 | Mexico | 177,133 | Community and Hospital | 42.6 (26-59) | 48.9 | - | - | 9.3 | - | - | 90.72 | poor |
| 189 | Zuo, Yalavarthi | 2020-04-24 | USA | 50 | Hospital | 61 (46-76) | 34.0 | - | - | 36.0 | - | - | 64.00 | poor |
| 190 | Sigel | 2020-06-28 | USA | 493 | Hospital | 60 (55-67) | 24.1 | - | - | 28.6 | - | - | 71.40 | poor |
| 191 | Nguyen | 2020-06-29 | USA | 689 | Community and Hospital | 55 (40-68) | 57.0 | - | - | 24.8 | - | - | 75.18 | poor |
| 192 | de Melo | 2020-06-29 | Brazil | 181 | Hospital | 55.3^ (34-76) | 60.8 | 9.9 | 12.2 | - | 38.1 | - | 39.78 | poor |

| | | | | | | | | | | | | | | |
|-----|-------------------------------|------------|-----------|--------|------------------------|---------------------|------|------|------|------|------|------|-------|------|
| 193 | Auvinen | 2020-06-29 | Finland | 61 | Hospital | 53 (41-67) | 36.0 | 18.0 | 27.9 | - | 54.1 | - | 0.00 | fair |
| 194 | Souza | 2020-06-28 | Brazil | 8,443 | Hospital | NA | 53.0 | - | - | 1.7 | - | 96.3 | 2.01 | poor |
| 195 | Mendy | 2020-06-27 | USA | 689 | Community and Hospital | 49.5 (35.2-67.5) | 47.0 | - | - | 24.7 | - | - | 75.33 | poor |
| 196 | Pongpirul | 2020-06-26 | Thailand | 193 | Hospital | 37 (29-53) | 41.5 | - | - | 15.0 | 66.3 | - | 18.65 | poor |
| 197 | Jin, Gu | 2020-06-25 | China | 6 | Hospital | 60.5^ (51-75) | 33.3 | 33.3 | - | - | - | - | 66.67 | poor |
| 198 | Favara | 2020-05-23 | UK | 70 | Community and Hospital | 41 (23-64) | 87.1 | 10.0 | - | - | - | - | 90.00 | poor |
| 199 | Fisman | 2020-06-23 | Canada | 21,922 | Community and Hospital | NA | 57.0 | - | - | 2.3 | - | - | 97.65 | poor |
| 200 | Madariaga | 2020-06-23 | USA | 103 | Community and Hospital | 41.8^ (27-55) | 48.5 | - | - | 25.2 | 74.8 | - | 0.00 | poor |
| 201 | Senkal | 2020-07-07 | Turkey | 611 | Hospital | 57^ (18-98) | 40.6 | 11.3 | - | - | - | - | 88.71 | poor |
| 202 | Mohamud | 2020-07-02 | USA | 6 | Hospital | 65.8^ (55-78) | 16.7 | - | - | 16.7 | - | - | 83.33 | poor |
| 203 | Magleby | 2020-06-30 | USA | 678 | Hospital | 68 (50-81) | 38.9 | - | - | 28.6 | - | - | 71.39 | poor |
| 204 | Kimmig | 2020-07-06 | USA | 111 | Hospital | 63^ (48-78) | 44.1 | 7.2 | 36.0 | - | 56.8 | - | 0.00 | fair |
| 205 | Bello-Chavolla, Antonio-Villa | 2020-07-04 | Mexico | 60,121 | Community and Hospital | 45.5^ (29-61) | 47.0 | - | - | 10.5 | - | - | 89.52 | poor |
| 206 | Zacharioudakis | 2020-07-04 | USA | 314 | Hospital | 64 (54-72) | 34.7 | - | - | 22.8 | - | - | 77.22 | poor |
| 207 | Antonio-Villa | 2020-07-04 | Mexico | 34,263 | Community and Hospital | 40^ (29-50) | 62.9 | 9.7 | - | - | - | - | 90.32 | poor |
| 208 | Patel | 2020-07-03 | USA | 129 | Hospital | 60.8^ (47-74) | 45.0 | 37.2 | - | - | - | 55.8 | 6.98 | poor |
| 209 | Merzon | 2020-07-03 | Israel | 7,807 | Community and Hospital | 46.2^ (NA) | 58.6 | - | - | 16.2 | - | - | 83.82 | poor |
| 34 | Trubiano | 2020-07-02 | Australia | 2,935 | Community and Hospital | 39 (29-53) | 63.5 | - | - | 8.8 | - | - | 91.18 | poor |

| | | | | | | | | | | | | | | |
|-----|--------------------|------------|----------|--------|------------------------|--------------------|------|------|------|------|------|------|-------|------|
| 210 | Fan | 2020-07-11 | UK | 1,425 | Community and Hospital | NA | 46.7 | 12.2 | 40.1 | - | 46.9 | - | 0.84 | fair |
| 211 | Shi, Resurreccion | 2020-07-11 | UK | 1,521 | Community and Hospital | 61.5^ (57-66.8) | 45.9 | - | - | 54.9 | - | - | 45.10 | poor |
| 212 | Maucourant | 2020-07-10 | Sweden | 27 | Hospital | 57 (18-78) | 22.2 | 11.1 | 25.9 | - | 40.7 | - | 22.22 | poor |
| 213 | Elmunzer | 2020-07-09 | Multiple | 1,992 | Hospital | 60^ (43-76) | 43.0 | 6.3 | 28.6 | - | 59.0 | - | 6.12 | fair |
| 214 | Alizadehsani | 2020-07-09 | Iran | 319 | Hospital | 45.48^ (26-63) | 55.5 | - | - | 0.3 | - | - | 99.69 | poor |
| 215 | Xie | 2020-07-07 | China | 619 | Hospital | NA | 52.0 | - | - | 8.2 | - | - | 91.76 | poor |
| 36 | Merkely | 2020-07-17 | Hungary | 10,474 | Community | 48.7^ (30-66) | 53.6 | 28.0 | 20.5 | - | 51.4 | - | 0.16 | good |
| 216 | Fox | 2020-07-17 | UK | 55 | Community and Hospital | 63 (23-88) | 31.0 | 1.8 | 10.9 | - | 56.4 | - | 30.91 | poor |
| 56 | Zhang, Cao | 2020-07-14 | China | 289 | Hospital | 57 (22-88) | 46.6 | 3.5 | 6.2 | - | - | - | 90.31 | poor |
| 217 | Martinez--Resendez | 2020-07-20 | Mexico | 8 | Hospital | 57 (48-69) | 25.0 | - | - | 12.5 | - | - | 87.50 | poor |
| 218 | Hoertel | 2020-07-20 | France | 12,612 | Hospital | 58.7^ (39-77) | 49.6 | - | - | 9.3 | - | - | 90.72 | poor |
| 219 | Mcgrail | 2020-07-19 | USA | 209 | Hospital | 62.5 (NA) | 38.8 | - | - | 18.7 | - | - | 81.34 | poor |
| 220 | Pandolfi | 2020-07-17 | Italy | 33 | Hospital | 62 (52-65) | 21.1 | 3.0 | 24.2 | - | 72.7 | - | 0.00 | fair |
| 28 | Girardeau | 2020-07-17 | France | 10 | Community | 30 (29-33) | 50.0 | 40.0 | 10.0 | - | - | - | 40.00 | poor |
| 221 | Kurashima | 2020-07-17 | Japan | 53 | Hospital | 62.9^ (49-76) | 35.8 | - | - | 50.9 | - | - | 49.06 | poor |
| 222 | Zhan | 2020-07-16 | China | 75 | Hospital | 57 (25-75) | 48.0 | - | - | 12.0 | - | - | 88.00 | poor |
| 223 | Omrani | 2020-07-16 | Qatar | 1,409 | Community and Hospital | 39 (30-50) | 17.2 | - | - | 9.2 | - | - | 90.77 | poor |
| 224 | Gupta | 2020-07-16 | USA | 496 | Hospital | 70 (60-78) | 46.0 | - | - | 7.3 | - | 31.7 | 61.09 | poor |
| 225 | Shi, Zuo | 2020-07-15 | USA | 172 | Hospital | 61.48^ (25-96) | 44.0 | - | - | 26.2 | - | - | 73.84 | poor |

| | | | | | | | | | | | | | | |
|-----|-----------------------|------------|-------------|--------|------------------------|------------------|------|------|------|------|------|------|-------|------|
| 226 | Hussein | 2020-07-15 | USA | 502 | Hospital | 60.9^ (45-76) | 52.0 | 9.0 | 22.1 | - | - | 68.9 | 0.00 | poor |
| 227 | Bian | 2020-07-15 | China | 28 | Hospital | 56^ (42-67) | 42.9 | 7.1 | - | - | - | - | 92.86 | poor |
| 228 | Eiros | 2020-07-14 | Spain | 139 | Community and Hospital | 52 (41-57) | 72.0 | 4.3 | 50.4 | - | - | - | 45.32 | poor |
| 229 | Marcos | 2020-07-14 | Spain | 918 | Hospital | 72.8^ (58-87) | 42.2 | 6.1 | - | 15.3 | - | - | 78.65 | poor |
| 230 | Hoertel, Sanchez-Rico | 2020-07-14 | France | 7,345 | Hospital | NA | 49.3 | 8.5 | - | - | - | - | 91.52 | poor |
| 231 | Soares | 2020-07-16 | Brazil | 10,713 | Community and Hospital | NA | 55.0 | 2.0 | - | - | - | 98.0 | 0.00 | poor |
| 232 | Zobairy | 2020-07-28 | Iran | 203 | Community and Hospital | 49.2^ (32-65) | 44.8 | 5.9 | - | - | - | 94.1 | 0.00 | poor |
| 233 | Altamimi | 2020-07-27 | Qatar | 68 | Hospital | 49^ (40-58) | 2.0 | 16.4 | - | - | - | 83.6 | 0.00 | poor |
| 234 | Thompson | 2020-07-27 | UK | 470 | Hospital | 71 (57-82) | 46.0 | 14.0 | 27.2 | - | 58.7 | - | 0.00 | fair |
| 235 | Reiter | 2020-07-26 | Austria | 235 | Community | 44.2^ (32-55) | 70.0 | 22.6 | 22.6 | - | 54.7 | - | 0.00 | fair |
| 236 | Motta | 2020-07-26 | USA | 374 | Hospital | 64.7^ (46-82) | 41.4 | - | - | 33.2 | 66.8 | - | 0.00 | poor |
| 237 | Santos | 2020-07-25 | USA | 43 | Community and Hospital | 50 (34-73) | 63.0 | - | - | 4.7 | - | - | 95.35 | poor |
| 238 | Schneeweiss | 2020-07-22 | USA | 24,313 | Community and Hospital | 67^ (53-80) | 53.0 | - | - | 2.9 | - | - | 97.12 | poor |
| 239 | Concha-Mejia | 2020-07-24 | Colombia | 72 | Community and Hospital | 46 (28-64) | 47.0 | 8.3 | 11.1 | - | - | - | 80.56 | poor |
| 240 | Izquierdo | 2020-07-24 | Spain | 71,192 | Community and Hospital | 42^ (18-66) | 59.0 | 10.0 | - | - | - | 90.0 | 0.00 | poor |
| 241 | Bernaola | 2020-07-21 | Spain | 1,645 | Hospital | NA | 38.5 | 2.5 | 10.9 | - | 86.6 | - | 0.00 | fair |
| 30 | Islam | 2020-08-18 | Bangladesh | 1,016 | Community and Hospital | 37 (28-49) | 35.9 | 18.2 | - | - | - | - | 77.85 | poor |
| 242 | Qi | 2020-03-03 | China | 267 | Hospital | 48 (35-65) | 45.2 | 19.9 | - | - | - | 80.1 | 0.00 | poor |
| 243 | Peters | 2020-08-15 | Netherlands | 1,893 | Hospital | 66.8^ (52-81) | 39.4 | 4.9 | - | - | - | - | 95.14 | poor |

| | | | | | | | | | | | | | | |
|-----|---------------|------------|------------|--------|------------------------|------------------|------|------|------|------|------|------|-------|-------|
| 244 | Ouyang | 2020-08-14 | China | 217 | Hospital | 46.5^ (30-62) | 53.5 | 16.6 | - | - | - | - | 83.41 | poor |
| 47 | Ward | 2020-08-21 | UK | 99,908 | Community | NA | 56.1 | 10.6 | - | - | - | 88.4 | 0.98 | poor* |
| 245 | Valenzuela | 2020-08-14 | Chile | 29 | Hospital | 56.9^ (43-70) | 6.9 | 17.2 | - | - | - | 82.8 | 0.00 | poor |
| 246 | Monteiro | 2020-08-14 | USA | 112 | Hospital | 61 (45-74) | 34.0 | 6.2 | 17.9 | - | 68.8 | - | 7.14 | fair |
| 247 | Philipose | 2020-08-14 | UK | 466 | Hospital | 67 (6-97) | 41.8 | 6.0 | 73.2 | - | 16.5 | - | 4.29 | fair |
| 248 | Weerahandi | 2020-08-14 | USA | 394 | Community | 63 (55-70) | 37.0 | 5.3 | 25.9 | - | 55.8 | - | 12.94 | fair |
| 29 | Ebinger | 2020-08-04 | USA | 6,062 | Community | 41.5^ (29-53) | 67.8 | 1.7 | - | - | - | - | 96.88 | poor |
| 249 | Altibi | 2020-08-11 | USA | 706 | Hospital | 66.7^ (51-81) | 43.0 | 4.0 | 37.3 | - | 58.8 | - | 0.00 | fair |
| 250 | Izzi-Engbeaya | 2020-08-11 | UK | 889 | Hospital | 65.8^ (48-83) | 40.0 | - | - | 21.3 | 33.2 | - | 45.6 | poor |
| 251 | Rizzo | 2020-08-11 | USA | 76,819 | Hospital | 54 (38-67) | 55.2 | 6.7 | 20.8 | - | 50.4 | - | 22.05 | poor |
| 252 | Dashti | 2020-08-04 | USA | 4,140 | Community and Hospital | 52 (36-65) | 55.0 | - | - | 28.4 | 51.6 | - | 19.95 | poor |
| 253 | Morshed | 2020-08-02 | Bangladesh | 103 | Community | 37 (31-53) | 28.2 | 31.1 | - | - | - | 68.9 | 0.00 | poor |
| 254 | Jun | 2020-08-01 | USA | 3,086 | Hospital | 66 (56-77) | 40.9 | 3.7 | 21.3 | - | 52.8 | - | 22.23 | poor |
| 255 | Higuchi | 2020-07-30 | Japan | 57 | Hospital | 52 (35-70) | 43.9 | 12.3 | 29.8 | - | 57.9 | - | 0.00 | fair |
| 256 | Zhou, Sun | 2020-07-29 | China | 144 | Hospital | 47 (38-56) | 46.5 | 9.0 | - | - | - | 91.0 | 0.00 | poor |
| 257 | Salerno | 2020-08-22 | USA | 15,920 | Hospital | 49 (30-65) | 57.0 | - | - | 36.8 | 55.9 | - | 7.29 | poor |
| 258 | Kumar | 2020-07-29 | India | 91 | Hospital | 47^ (41-52) | 21.0 | 44.0 | - | - | - | - | 56.04 | poor |
| 259 | Hao | 2020-06-01 | China | 788 | Hospital | 46 (35-56) | 48.4 | 6.9 | - | - | - | - | 93.15 | poor |
| 260 | Iversen | 2020-08-03 | Denmark | 28,792 | Community and Hospital | 44.4^ (31-57) | 78.9 | 16.0 | 6.5 | - | 76.8 | - | 0.67 | fair |

| | | | | | | | | | | | | | | |
|-----|---------------|------------|--------|-----------|------------------------|------------------|------|------|------|------|------|---|-------|------|
| 261 | Hippisley-Cox | 2020-07-13 | UK | 8,275,949 | Community and Hospital | 48.5^ (30-66) | 50.3 | 17.2 | 21.4 | - | 57.3 | - | 4.04 | fair |
| 262 | Fillmore | 2020-08-24 | USA | 22,914 | Community and Hospital | NA | - | 37.5 | 40.7 | - | 15.5 | - | 6.38 | fair |
| 263 | Rashid | 2020-08-22 | UK | 517 | Hospital | 72.8^ (59-86) | 31.9 | 9.9 | 29.0 | - | 29.4 | - | 31.72 | poor |
| 264 | Pan | 2020-08-22 | USA | 12,084 | Community and Hospital | 45.5^ (27-63) | 54.3 | - | - | 17.5 | - | - | 82.49 | poor |
| 265 | Alkurt | 2020-08-20 | Turkey | 932 | Community and Hospital | 34.8^ (25-44) | 64.4 | 24.5 | - | - | - | - | 75.54 | poor |
| 266 | Zhao, Chen | 2020-07-30 | USA | 641 | Hospital | 60 (NA) | 40.1 | 21.7 | - | - | - | - | 78.32 | poor |
| 267 | Holman | 2020-08-13 | UK | 10,989 | Community and Hospital | NA | 38.8 | 5.5 | 42.6 | - | 49.0 | - | 2.82 | fair |
| 268 | Qu | 2020-07-29 | China | 246 | Hospital | 53.6^ (38-68) | 53.3 | 42.3 | - | - | - | - | 57.72 | poor |
| 269 | Chand | 2020-08-19 | USA | 300 | Hospital | 58.2^ (45-70) | 39.3 | 22.3 | - | - | - | - | 77.67 | poor |

Note. – Age not provided for total sample; ^ Denotes mean (SD). * This study was rated as ‘poor’ quality as the manuscript only presents data for current (but not former) smokers despite having obtained complete smoking status, thus resulting in >20% missing data on smoking status.

Table 1. Characteristics of included studies.




Figure 2a. Weighted mean prevalence of current smoking in included studies with 95% bootstrap confidence intervals compared with national current smoking prevalence (solid red lines), split by country. Shape corresponds to study setting (community, community and hospital, hospital) and shape size corresponds to relative study sample size.




Figure 2b. Weighted mean prevalence of former smoking in included studies (where this was reported) with 95% bootstrap confidence intervals compared with national former smoking prevalence (solid red lines), split by country. Shape corresponds to study setting (community, community and hospital, hospital) and shape size corresponds to relative study sample size.

| Author | Total population tested | SARS-CoV-2 negative | | | | | | SARS-CoV-2 positive | | | | | |
|-----------------|-------------------------|---------------------|--------------------|-------------------|---------------------------|------------------|----------------|---------------------|--------------------|-------------------|---------------------------|-------------------|----------------|
| | | N (%) | Current smoker (%) | Former smoker (%) | Current/former smoker (%) | Never smoker (%) | Not stated (%) | N (%) | Current smoker (%) | Former smoker (%) | Current/former smoker (%) | Never smoker (%) | Not stated (%) |
| Rentsch | 3528 | 2974 (84.30%) | 1444 (48.55%) | 704 (23.67%) | - | 826 (27.77%) | - | 554 (15.70%) | 159 (28.70%) | 179 (32.31%) | - | 216 (38.99%) | - |
| Fontanet | 661 | 490 (74.13%) | 64 (13.06%) | - | - | 426 (86.94%) | - | 171 (25.87%) | 5 (2.92%) | - | - | 166 (97.08%) | - |
| Cho | 1331 | 793 (59.58%) | 142 (17.91%) | 214 (26.99%) | - | 437 (55.11%) | - | 538 (40.42%) | 111 (20.63%) | 145 (26.95%) | - | 282 (52.42%) | - |
| Shah | 243 | 212 (87.24%) | 52 (24.53%) | 47 (22.17%) | - | 113 (53.30%) | - | 29 (11.93%) | 0 (0.00%) | 9 (31.03%) | - | 20 (68.97%) | - |
| Kolin | 1474 | 805 (54.61%) | 141 (17.52%) | 307 (38.14%) | - | 354 (43.98%) | 3 (0.37%) | 669 (45.39%) | 72 (10.76%) | 285 (42.60%) | - | 303 (45.29%) | 9 (1.35%) |
| de Lusignan | 3291 | 2740 (83.26%) | 366 (13.36%) | 1450 (52.92%) | - | 924 (33.72%) | - | 551 (16.74%) | 47 (8.53%) | 303 (54.99%) | - | 201 (36.48%) | - |
| Valenti | 789 | 689 (87.33%) | 197 (28.59%) | - | - | - | 492 (71.41%) | 40 (5.07%) | 7 (17.50%) | - | - | - | 33 (82.50%) |
| Parrotta | 76 | 39 (51.32%) | 1 (2.56%) | 10 (25.64%) | - | 27 (69.23%) | 1 (2.56%) | 37 (48.68%) | 1 (2.70%) | 10 (27.03%) | - | 25 (67.57%) | 1 (2.70%) |
| Berumen | 102875 | 71353 (69.36%) | - | - | 7173 (10.05%) | 64180 (89.95%) | - | 31522 (30.64%) | - | - | 2748 (8.72%) | 28774 (91.28%) | - |
| Israel | 24906 | 20755 (83.33%) | 3783 (18.23%) | 2671 (12.87%) | - | 14301 (68.90%) | - | 41151 (165.23%) | 406 (0.99%) | 483 (1.17%) | - | 3262 (7.93%) | - |
| del Valle | 1108 | 143 (12.91%) | 27 (18.88%) | 53 (37.06%) | - | - | 63 (44.06%) | 965 (87.09%) | 55 (5.70%) | 293 (30.36%) | - | - | 617 (63.94%) |
| Romao | 34 | 20 (58.82%) | - | - | 5 (25.00%) | - | 15 (75.00%) | 14 (41.18%) | - | - | 4 (28.57%) | - | 10 (71.43%) |
| Ramlall | 11116 | 4723 (42.49%) | - | - | - | - | - | 6393 (57.51%) | - | - | 1643.001 (25.70%) | 4749.999 (74.30%) | - |
| Sharma | 501 | 267 (53.29%) | - | - | 1 (0.37%) | - | 266 (99.63%) | 234 (46.71%) | - | - | 20 (8.55%) | - | 214 (91.45%) |
| Eugen-Olsen | 407 | 290 (71.25%) | 76 (26.21%) | 104 (35.86%) | - | 102 (35.17%) | - | 117 (28.75%) | 8 (6.84%) | 46 (39.32%) | - | 59 (50.43%) | - |
| Raisi-Estabragh | 4510 | 3184 (70.60%) | - | - | 1653 (51.92%) | - | 1531 (48.08%) | 1326 (29.40%) | - | - | 683 (51.51%) | - | 643 (48.49%) |

| | | | | | | | | | | | | | |
|-----------------------|--------|--------------------|------------------|------------------|-----------------|--------------------|-------------------|-------------------|------------------|-----------------|--------------|-------------------|-------------------|
| Houlihan | 177 | 97 (54.80%) | 14 (14.43%) | 14 (14.43%) | - | 69 (71.13%) | - | 80 (45.20%) | 7 (8.75%) | 19 (23.75%) | - | 54 (67.50%) | - |
| McQueenie | 428199 | 424355 (99.10%) | - | - | 189299 (44.61%) | 235056 (55.39%) | - | 1311 (0.31%) | - | - | 669 (51.03%) | 642 (48.97%) | - |
| Woolford | 4474 | 3161 (70.65%) | 441 (13.95%) | 1194 (37.77%) | - | 1526 (48.28%) | - | 1313 (29.35%) | 145 (11.04%) | 525 (39.98%) | - | 643 (48.97%) | - |
| Lan | 104 | 83 (79.81%) | - | - | 24 (28.92%) | - | 59 (71.08%) | 21 (20.19%) | - | - | 1 (4.76%) | - | 20 (95.24%) |
| Hernandez- Garduno | 32583 | 20279 (62.24%) | - | - | 2399 (11.83%) | 17861 (88.08%) | - | 12304 (37.76%) | - | - | 1191 (9.68%) | 11083 (90.08%) | - |
| Govind | 6215 | 6207 (99.87%) | 4104 (66.12%) | 1669 (26.89%) | - | 342 (5.51%) | - | 102 (1.64%) | 78 (76.47%) | 20 (19.61%) | - | 2 (1.96%) | - |
| Gu | 4699 | 3815 (81.19%) | 360 (9.44%) | 1142 (29.93%) | - | 2313 (60.63%) | - | 884 (18.81%) | 40 (4.52%) | 264 (29.86%) | - | 580 (65.61%) | - |
| Kibler | 702 | 680 (96.87%) | 25 (3.68%) | - | - | - | 655 (96.32%) | 22 (3.13%) | 1 (4.55%) | - | - | - | 21 (95.45%) |
| Petrilli | 10620 | 5341 (50.29%) | 3454 (64.67%) | 816 (15.28%) | - | 541 (10.13%) | 530 (9.92%) | 5279 (49.71%) | 3268 (61.91%) | 902 (17.09%) | - | 288 (5.46%) | 821 (15.55%) |
| Bello-Chavolla | 150200 | 98567 (65.62%) | - | - | 9624 (9.76%) | - | 88943 (90.24%) | 51633 (34.38%) | - | - | 4366 (8.46%) | - | 47267 (91.54%) |
| Auvinen | 61 | 33 (54.10%) | 10 (30.30%) | 8 (24.24%) | - | 15 (45.45%) | - | 28 (45.90%) | 1 (3.57%) | 9 (32.14%) | - | 18 (64.29%) | - |
| Favara | 70 | 55 (78.57%) | 5 (9.09%) | - | - | - | 50 (90.91%) | 15 (21.43%) | 2 (13.33%) | - | - | - | 13 (86.67%) |
| Antonio-Villa | 34263 | 23338 (68.11%) | 2293 (9.83%) | - | - | - | 21045 (90.17%) | 10925 (31.89%) | 1023 (9.36%) | - | - | - | 9902 (90.64%) |
| Merzon | 7807 | 7025 (89.98%) | - | - | 1136 (16.17%) | - | 5889 (83.83%) | 782 (10.02%) | - | - | 127 (16.24%) | - | 655 (83.76%) |
| Trubiano | 2676 | 2827 (105.64%) | - | - | 256 (9.06%) | - | 2586 (91.48%) | 108 (4.04%) | - | - | 3 (2.78%) | - | 105 (97.22%) |
| Shi, Resurreccion | 1521 | 1265 (83.17%) | - | - | 681 (53.83%) | - | 584 (46.17%) | 256 (16.83%) | - | - | 154 (60.16%) | - | 102 (39.84%) |
| Riley | 120620 | 120461 (99.87%) | 2594 (2.15%) | - | - | 19914 (16.53%) | 97953 (81.32%) | 159 (0.13%) | 3 (1.89%) | - | - | 17 (10.69%) | 139 (87.42%) |
| Alizadehsani | 319 | 196 (61.44%) | - | - | - | - | 196 (100.00%) | 123 (38.56%) | - | - | 1 (0.81%) | - | 122 (99.19%) |
| Merkely | 10474 | 10336 (98.68%) | 2904 (28.10%) | 2107 (20.39%) | - | 5310 (51.37%) | 15 (0.15%) | 70 (0.67%) | 16 (22.86%) | 15 (21.43%) | - | 38 (54.29%) | 1 (1.43%) |

| | | | | | | | | | | | | | |
|---------------|---------|-------------------|-------------------|------------------|---------------|-------------------|-------------------|------------------|-----------------|------------------|--------------|-------------------|------------------|
| Mcgrail | 209 | 118 (56.46%) | - | - | 31 (26.27%) | - | 87 (73.73%) | 91 (43.54%) | - | - | 8 (8.79%) | - | 83 (91.21%) |
| Izquierdo | 71192 | NA (NA%) | - | - | - | - | - | 1006 (1.41%) | 111 (11.03%) | - | - | - | 895 (88.97%) |
| Ward | 99908 | 94416 (94.50%) | 10202 (10.81%) | - | - | - | 84214 (89.19%) | 5492 (5.50%) | 433 (7.88%) | - | - | - | 5059 (92.12%) |
| Ebinger | 6062 | 5850 (96.50%) | 99 (1.69%) | - | - | - | 5668 (96.89%) | 212 (3.50%) | 3 (1.42%) | - | - | - | 205 (96.70%) |
| Salerno | 15920 | 14753 (92.67%) | - | - | 5517 (37.40%) | 8278 (56.11%) | 958 (6.49%) | 1167 (7.33%) | - | - | 339 (29.05%) | 626 (53.64%) | 202 (17.31%) |
| Iversen | 28792 | 27629 (95.96%) | 4430 (16.03%) | 1799 (6.51%) | - | 21217 (76.79%) | 246 (0.89%) | 1163 (4.04%) | 177 (15.22%) | 78 (6.71%) | - | 898 (77.21%) | 10 (0.86%) |
| Hippisley-Cox | 8275949 | NA (NA%) | - | - | - | - | - | 19486 (0.24%) | 1354 (6.95%) | 5715 (29.33%) | - | 12036 (61.77%) | 381 (1.96%) |
| Fillmore | 22914 | 21120 (92.17%) | 8137 (38.53%) | 8416 (39.85%) | - | 3227 (15.28%) | 1340 (6.34%) | 1794 (7.83%) | 452 (25.20%) | 899 (50.11%) | - | 322 (17.95%) | 121 (6.74%) |
| Alkurt | 119 | NA (NA%) | - | - | - | - | - | 119 (100.00%) | 14 (11.76%) | - | - | - | 105 (88.24%) |

Note. Niedzwiedz et al. reported on SARS-CoV-2 infection by smoking status in multivariable analyses but did not present raw data.

Table 2. SARS-CoV-2 infection by smoking status.




Figure 3. Forest plot for risk of testing positive for SARS-CoV-2 in current vs. never smokers. * This was a 'good' quality study.




Figure 4. Forest plot for risk of testing positive for SARS-CoV-2 in former vs. never smokers.

* This was a 'good' quality study.

| Community | | | | | | | | | Hospitalised | | | | | | |
|---------------|-------------------------|--------------|--------------------|-------------------|---------------------------|------------------|--------------------------|----------------|--------------|--------------------|-------------------|---------------------------|------------------|--------------------------|----------------|
| Author | Population with outcome | N (%) | Current smoker (%) | Former smoker (%) | Current/former smoker (%) | Never smoker (%) | Never/unknown smoker (%) | Not stated (%) | N (%) | Current smoker (%) | Former smoker (%) | Current/former smoker (%) | Never smoker (%) | Never/unknown smoker (%) | Not stated (%) |
| Rentsch | 554 | 269 (48%) | 69 (25.65%) | 90 (33.46%) | - | 110 (40.89%) | - | - | 285 (51%) | 90 (31.58%) | 89 (31.23%) | - | 106 (37.19%) | - | - |
| Chow (US CDC) | 6637 | 5143 (77%) | 61 (1.19%) | 80 (1.56%) | - | - | - | 5002 (97.26%) | 1494 (22%) | 27 (1.81%) | 78 (5.22%) | - | - | - | 1389 (92.97%) |
| Argenziano | 1000 | 151 (15%) | 14 (9.27%) | 18 (11.92%) | - | 119 (78.81%) | - | - | 849 (84%) | 35 (4.12%) | 161 (18.96%) | - | 653 (76.91%) | - | - |
| Lubetzky | 54 | 15 (27%) | - | - | 4 (26.67%) | - | - | 11 (73.33%) | 39 (72%) | - | - | 8 (20.51%) | - | - | 31 (79.49%) |
| Carillo-Vega | 9946 | 3922 (39%) | 408 (10.40%) | - | - | - | - | 3514 (89.60%) | 6024 (60%) | 486 (8.07%) | - | - | - | - | 5538 (91.93%) |
| Yanover | 4353 | 4180 (96%) | 484 (11.58%) | 118 (2.82%) | - | 3578 (85.60%) | - | - | 173 (3%) | 30 (17.34%) | 11 (6.36%) | - | 132 (76.30%) | - | - |
| Hamer | 387109 | 386349 (99%) | 37333 (9.66%) | 134542 (34.82%) | - | 214474 (55.51%) | - | - | 760 (0%) | 93 (12.24%) | 313 (41.18%) | - | 354 (46.58%) | - | - |
| Heili-Frades | 4712 | 1973 (41%) | 121 (6.13%) | 222 (11.25%) | - | - | 1630 (82.62%) | 1630 (82.62%) | 2739 (58%) | 112 (4.09%) | 598 (21.83%) | - | - | 2029 (74.08%) | - |
| Freites | 123 | 69 (56%) | 1 (1.45%) | - | - | - | - | 68 (98.55%) | 54 (43%) | 3 (5.56%) | - | - | - | - | 51 (94.44%) |
| Berumen | 102875 | 18832 (18%) | - | - | 1546 (8.21%) | - | 17286 (91.79%) | - | 12690 (12%) | - | - | 1202 (9.47%) | - | 11488 (90.53%) | - |
| Gianfrancesco | 600 | 323 (53%) | - | - | 61 (18.89%) | - | - | 262 (81.11%) | 277 (46%) | - | - | 68 (24.55%) | - | - | 209 (75.45%) |

| | | | | | | | | | | | | | | | |
|------------------|-------|----------------|-----------------|-----------------|-------------|------------------|-------------------|-----------------|--------------------|-----------------|-----------------|--------------|------------------|-------------------|-----------------|
| Chaudhry | 40 | 19 (47%) | - | - | 0 (0.00%) | - | - | 19 (100.00%) | 21 (52%) | - | - | 6 (28.57%) | - | - | 15 (71.43%) |
| Giannouchos | 89756 | 58485 (65%) | 4679 (8.00%) | - | - | - | 53806 (92.00%) | - | 3127 1 (34%) | 2721 (8.70%) | - | - | - | 28550 (91.30%) | - |
| Wang, Oekelen | 57 | 22 (38%) | - | - | 6 (27.27%) | - | - | 16 (72.73%) | 36 (63%) | - | - | 15 (41.67%) | - | - | 20 (55.56%) |
| Miyara | 470 | 132 (28%) | 14 (10.61%) | 41 (31.06%) | - | 77 (58.33%) | - | - | 338 (71%) | 18 (5.33%) | 111 (32.84%) | - | 209 (61.83%) | - | - |
| Suleyman | 463 | 108 (23%) | - | - | 23 (21.30%) | - | - | 85 (78.70%) | 355 (76%) | - | - | 137 (38.59%) | - | - | 218 (61.41%) |
| Garassino | 196 | 48 (24%) | 10 (20.83%) | 27 (56.25%) | - | 11 (22.92%) | - | - | 152 (77%) | 38 (25.00%) | 84 (55.26%) | - | 26 (17.11%) | - | - |
| Siso-Almirall | 260 | 119 (45%) | - | - | 31 (26.05%) | - | - | 88 (73.95%) | 141 (54%) | - | - | 50 (35.46%) | - | - | 91 (64.54%) |
| Gu | 884 | 511 (57%) | 30 (5.87%) | 126 (24.66%) | - | 355 (69.47%) | - | - | 373 (42%) | 10 (2.68%) | 138 (37.00%) | - | 225 (60.32%) | - | - |
| Killerby | 531 | 311 (58%) | - | - | 37 (11.90%) | 222 (71.38%) | - | 52 (16.72%) | 220 (41%) | - | - | 54 (24.55%) | 157 (71.36%) | - | 9 (4.09%) |
| Petrilli | 5279 | 2538 (48%) | 147 (5.79%) | 337 (13.28%) | - | 1678 (66.12%) | - | 376 (14.81%) | 2741 (51%) | 141 (5.14%) | 565 (20.61%) | - | 1590 (58.01%) | - | 445 (16.23%) |
| Nguyen | 689 | 333 (48%) | - | - | 57 (17.12%) | - | - | 276 (82.88%) | 356 (51%) | - | - | 114 (32.02%) | - | - | 242 (67.98%) |
| Mendy | 689 | 473 (68%) | - | - | 84 (17.76%) | - | - | 389 (82.24%) | 216 (31%) | - | - | 86 (39.81%) | - | - | 130 (60.19%) |
| Soares | 10713 | 9561 (89%) | 132 (1.38%) | - | - | - | 9429 (98.62%) | - | 1152 (10%) | 77 (6.68%) | - | - | - | 1075 (93.32%) | - |
| Zobairy | 203 | 65 (32%) | 1 (1.54%) | - | - | - | 64 (98.46%) | - | 138 (67%) | 11 (7.97%) | - | - | - | 127 (92.03%) | - |

| | | | | | | | | | | | | | | | |
|-----------|-------|----------------|-----------------|-------------------|---------------|-------------------|--------------|-------------------|--------------------|-----------------|------------------|--------------|------------------|--------------|------------------|
| Izquierdo | 1006 | 743 (73%) | 52 (7.00%) | - | - | - | 691 (93.00%) | - | 263 (26%) | 16 (6.08%) | - | - | - | 247 (93.92%) | - |
| Rizzo | 76819 | 60039 (78%) | 3931 (6.55%) | 11379 (18.95%) | - | 30042 (50.04%) | - | 14687 (24.46%) | 1678 0 (21%) | 1254 (7.47%) | 4585 (27.32%) | - | 8693 (51.81%) | - | 2248 (13.40%) |
| Dashti | 4140 | 2759 (66%) | - | - | 600 (21.75%) | 1541 (55.85%) | - | 618 (22.40%) | 1381 (33%) | - | - | 577 (41.78%) | - | 596 (43.16%) | 208 (15.06%) |
| Pan | 12084 | 8548 (70%) | - | - | 1263 (14.78%) | - | - | 7285 (85.22%) | 3536 (29%) | - | - | 874 (24.72%) | - | - | 2662 (75.28%) |

Table 3. Hospitalisation with COVID-19 by smoking status.




Figure 5. Forest plot for risk of hospitalisation in current vs. never smokers.




Figure 6. Forest plot for risk of hospitalisation in former vs. never smokers.

| Author | Population with severity | Non severe disease | | | | | | | Severe disease | | | | | | |
|--------------|--------------------------|--------------------|--------------------|-------------------|---------------------------|------------------|--------------------------|----------------|----------------|--------------------|-------------------|---------------------------|------------------|--------------------------|----------------|
| | | N (%) | Current smoker (%) | Former smoker (%) | Current/former smoker (%) | Never smoker (%) | Never/unknown smoker (%) | Not stated (%) | N (%) | Current smoker (%) | Former smoker (%) | Current/former smoker (%) | Never smoker (%) | Never/unknown smoker (%) | Not stated (%) |
| Guan, Ni | 1085 | 913 (84%) | 108 (11.83%) | 12 (1.31%) | - | 793 (86.86%) | - | - | 172 (15%) | 29 (16.86%) | 9 (5.23%) | - | 134 (77.91%) | - | - |
| Zhang, Dong | 9 | 3 (33%) | 0 (0.00%) | 3 (100.00%) | - | 0 (0.00%) | - | - | 6 (66%) | 2 (33.33%) | 4 (66.67%) | - | 0 (0.00%) | - | - |
| Wan | 9 | 8 (88%) | 8 (100.00%) | 0 (0.00%) | - | 0 (0.00%) | - | - | 1 (11%) | 1 (100.00%) | 0 (0.00%) | - | 0 (0.00%) | - | - |
| Huang, Wang | 3 | 3 (100%) | 3 (100.00%) | 0 (0.00%) | - | 0 (0.00%) | - | - | 0 (0%) | 0 (-%) | 0 (-%) | - | 0 (-%) | - | - |
| Rentsch | 285 | 168 (58%) | 47 (27.98%) | 53 (31.55%) | - | 68 (40.48%) | - | - | 117 (41%) | 43 (36.75%) | 36 (30.77%) | - | 38 (32.48%) | - | - |
| Hu | 323 | 151 (46%) | - | - | 12 (7.95%) | - | 139 (92.05%) | - | 172 (53%) | - | - | 26 (15.12%) | - | 146 (84.88%) | - |
| Wang, Pan | 125 | 100 (80%) | - | - | 9 (9.00%) | - | 91 (91.00%) | - | 25 (20%) | - | - | 7 (28.00%) | - | 18 (72.00%) | - |
| Kim | 27 | 21 (77%) | 3 (14.29%) | - | - | - | 18 (85.71%) | - | 6 (22%) | 2 (33.33%) | 0 (0.00%) | - | - | 4 (66.67%) | - |
| Shi, Yu | 474 | 425 (89%) | - | - | 34 (8.00%) | - | 391 (92.00%) | - | 49 (10%) | - | - | 6 (12.24%) | - | 43 (87.76%) | - |
| Liao, Feng | 148 | 92 (62%) | - | - | 5 (5.43%) | - | - | 87 (94.57%) | 56 (37%) | 3 (5.36%) | - | - | - | - | 53 (94.64%) |
| Shi, Ren | 134 | 88 (65%) | - | - | 8 (9.09%) | - | - | 80 (90.91%) | 46 (34%) | - | - | 6 (13.04%) | - | - | 40 (86.96%) |
| Hadjadj | 50 | 15 (30%) | 1 (6.67%) | 2 (13.33%) | - | 12 (80.00%) | - | - | 35 (70%) | 0 (0.00%) | 7 (20.00%) | - | 28 (80.00%) | - | - |
| Zheng, Xiong | 73 | 43 (58%) | - | - | 6 (13.95%) | 37 (86.05%) | - | - | 30 (41%) | - | - | 2 (6.67%) | 28 (93.33%) | - | - |
| de la Rica | 48 | 26 (54%) | - | - | 6 (23.08%) | - | - | 20 (76.92%) | 20 (41%) | - | - | 4 (20.00%) | - | - | 16 (80.00%) |
| Yin, Yang | 106 | 47 (44%) | - | - | 6 (12.77%) | - | - | 41 (87.23%) | 59 (55%) | - | - | 12 (20.34%) | - | - | 47 (79.66%) |
| Allenbach | 147 | 100 (68%) | - | - | 9 (9.00%) | - | - | 91 (91.00%) | 47 (31%) | - | - | 0 (0.00%) | - | - | 47 (100.00%) |

| | | | | | | | | | | | | | | | |
|-----------------|-------|----------------|-----------------|-----------------|-------------|-----------------|----------------|-----------------|----------------|-----------------|-----------------|-------------|----------------|----------------|-----------------|
| Goyal | 393 | 263 (66%) | 14 (5.32%) | - | - | - | - | 249 (94.68%) | 130 (33%) | 6 (4.62%) | - | - | - | - | 124 (95.38%) |
| Feng | 454 | 333 (73%) | 27 (8.11%) | - | - | - | - | 306 (91.89%) | 121 (26%) | 17 (14.05%) | - | - | - | - | 104 (85.95%) |
| Yao | 108 | 83 (76%) | 1 (1.20%) | - | - | - | - | 82 (98.80%) | 25 (23%) | 3 (12.00%) | - | - | - | - | 22 (88.00%) |
| Sami | 490 | 400 (81%) | 53 (13.25%) | - | - | - | - | 347 (86.75%) | 90 (18%) | 16 (17.78%) | - | - | - | - | 74 (82.22%) |
| Regina | 200 | 163 (81%) | 9 (5.52%) | - | - | - | - | 154 (94.48%) | 37 (18%) | 0 (0.00%) | - | - | - | - | 37 (100.00%) |
| Feuth | 28 | 21 (75%) | 1 (4.76%) | 7 (33.33%) | - | 13 (61.90%) | - | - | 7 (25%) | 2 (28.57%) | 1 (14.29%) | - | 4 (57.14%) | - | - |
| Mejia-Vilet | 329 | 214 (65%) | - | - | 13 (6.07%) | - | - | 201 (93.93%) | 115 (34%) | - | - | 10 (8.70%) | - | - | 105 (91.30%) |
| Chen, Jiang | 135 | 54 (40%) | - | - | 4 (7.41%) | - | - | 50 (92.59%) | 81 (60%) | - | - | 9 (11.11%) | - | - | 72 (88.89%) |
| Vaquero-Roncero | 146 | 75 (51%) | - | - | 4 (5.33%) | - | - | 71 (94.67%) | 71 (48%) | - | - | 6 (8.45%) | - | - | 65 (91.55%) |
| Kim, Garg | 2490 | 1692 (67%) | 112 (6.62%) | 395 (23.35%) | - | - | 1185 (70.04%) | - | 798 (32%) | 38 (4.76%) | 247 (30.95%) | - | - | 512 (64.16%) | - |
| Wu | 174 | 92 (52%) | - | - | 47 (51.09%) | - | 45 (48.91%) | - | 82 (47%) | 11 (13.41%) | - | - | - | 71 (86.59%) | - |
| Chaudhry | 40 | 34 (85%) | - | - | 5 (14.71%) | - | - | 29 (85.29%) | 6 (15%) | - | - | 1 (16.67%) | - | - | 5 (83.33%) |
| Garibaldi | 832 | 532 (63%) | 25 (4.70%) | 107 (20.11%) | - | - | - | 400 (75.19%) | 300 (36%) | 21 (7.00%) | 81 (27.00%) | - | - | - | 198 (66.00%) |
| Kuderer | 928 | 686 (73%) | 35 (5.10%) | 210 (30.61%) | - | 370 (53.94%) | - | 29 (4.23%) | 242 (26%) | 8 (3.31%) | 116 (47.93%) | - | 99 (40.91%) | 15 (6.20%) | 4 (1.65%) |
| Romao | 14 | 14 (100%) | - | - | 4 (28.57%) | - | - | 10 (71.43%) | 0 (0%) | - | - | - | - | - | - |
| Giannouchos | 89756 | 78050 (86%) | 6322 (8.10%) | - | - | - | 71728 (91.90%) | - | 11706 (13%) | 1089 (9.30%) | - | - | - | 10617 (90.70%) | - |
| Cen | 1007 | 720 (71%) | - | - | 70 (9.72%) | - | - | 650 (90.28%) | 287 (28%) | - | - | 18 (6.27%) | - | - | 269 (93.73%) |
| Maraschini | 132 | 89 (67%) | - | 11 (12.36%) | - | 78 (87.64%) | - | - | 43 (32%) | - | 3 (6.98%) | - | 40 (93.02%) | - | - |
| Siso-Almirall | 260 | 212 (81%) | - | - | 60 (28.30%) | - | - | 152 (71.70%) | 48 (18%) | - | - | 21 (43.75%) | - | - | 27 (56.25%) |

| | | | | | | | | | | | | | | | |
|------------------------------|------|---------------|----------------|-----------------|--------------|------------------|--------------|------------------|---------------|-----------------|-----------------|-------------|-----------------|-------------|------------------|
| Gu | 884 | 511 (57%) | 30 (5.87%) | 126 (24.66%) | - | 355 (69.47%) | - | - | 134 (15%) | 3 (2.24%) | 61 (45.52%) | - | 70 (52.24%) | - | - |
| Petrilli | 2729 | 1739 (63%) | 97 (5.58%) | 325 (18.69%) | - | 1067 (61.36%) | - | 250 (14.38%) | 990 (36%) | 44 (4.44%) | 236 (23.84%) | - | 517 (52.22%) | - | 193 (19.49%) |
| Mendy | 689 | 598 (86%) | - | - | 133 (22.24%) | - | - | 465 (77.76%) | 91 (13%) | - | - | 37 (40.66%) | - | - | 54 (59.34%) |
| Pongpirul | 193 | 161 (83%) | - | - | 25 (15.53%) | 106 (65.84%) | - | 30 (18.63%) | 32 (16%) | - | - | 4 (12.50%) | 21 (65.62%) | - | 7 (21.88%) |
| Jin, Gu | 6 | 2 (33%) | - | - | 0 (0.00%) | - | - | 4 (200.00%) | 4 (66%) | - | - | 2 (50.00%) | - | - | 2 (50.00%) |
| Senkal | 611 | 446 (73%) | 48 (10.76%) | - | - | - | - | 398 (89.24%) | 165 (27%) | 21 (12.73%) | - | - | - | - | 144 (87.27%) |
| Patel | 129 | 89 (68%) | 26 (29.21%) | - | - | - | 58 (65.17%) | 5 (5.62%) | 40 (31%) | 22 (55.00%) | - | - | - | 14 (35.00%) | 4 (10.00%) |
| Maucourant | 27 | 10 (37%) | 1 (10.00%) | 2 (20.00%) | - | 2 (20.00%) | - | 5 (50.00%) | 17 (62%) | 2 (11.76%) | 5 (29.41%) | - | 9 (52.94%) | - | 1 (5.88%) |
| Xie | 619 | 469 (75%) | - | - | 32 (6.82%) | - | - | 437 (93.18%) | 150 (24%) | - | - | 19 (12.67%) | - | - | 131 (87.33%) |
| Fox | 55 | 30 (54%) | 1 (3.33%) | 4 (13.33%) | - | 17 (56.67%) | - | 8 (26.67%) | 25 (45%) | 0 (0.00%) | 2 (8.00%) | - | 14 (56.00%) | - | 9 (36.00%) |
| Zhang, Cao | 240 | 162 (67%) | 2 (1.23%) | 6 (3.70%) | - | - | - | 154 (95.06%) | 78 (32%) | 4 (5.13%) | 4 (5.13%) | - | - | - | 70 (89.74%) |
| Kurashima | 53 | 10 (18%) | - | - | 3 (30.00%) | - | - | 7 (70.00%) | 43 (81%) | - | - | 24 (55.81%) | - | - | 19 (44.19%) |
| Zhan | 75 | NA (NA%) | - | - | - | - | - | - | 75 (100%) | - | - | 9 (12.00%) | - | - | 66 (88.00%) |
| Omrani | 858 | 806 (93%) | - | - | 121 (15.01%) | - | - | 685 (84.99%) | 52 (6%) | - | - | 9 (17.31%) | - | - | 43 (82.69%) |
| Marcos | 918 | 555 (60%) | 38 (6.85%) | - | 69 (12.43%) | - | - | 448 (80.72%) | 363 (39%) | 18 (4.96%) | - | 71 (19.56%) | - | - | 292 (80.44%) |
| Hoertel, Sanchez- Rico | 7345 | 6014 (81%) | 433 (7.20%) | - | - | - | - | 5581 (92.80%) | 1331 (18%) | 190 (14.27%) | - | - | - | - | 1141 (85.73%) |
| Qi | 267 | 217 (81%) | 22 (10.14%) | - | - | - | 195 (89.86%) | - | 50 (18%) | 31 (62.00%) | - | - | - | 19 (38.00%) | - |
| Monteiro | 112 | 84 (75%) | 3 (3.57%) | 14 (16.67%) | - | 63 (75.00%) | - | 4 (4.76%) | 28 (25%) | 4 (14.29%) | 6 (21.43%) | - | 14 (50.00%) | - | 4 (14.29%) |

| | | | | | | | | | | | | | | | |
|---------------|------|--------------|----------------|---|--------------|-----------------|-------------|-----------------|--------------|----------------|-----------------|--------------|-----------------|-------------|-----------------|
| Dashti | 1381 | 619 (44%) | - | - | 239 (38.61%) | 292 (47.17%) | - | 88 (14.22%) | 762 (55%) | - | - | 338 (44.36%) | 304 (39.90%) | - | 120 (15.75%) |
| Morshed | 103 | 87 (84%) | 28 (32.18%) | - | - | - | 59 (67.82%) | - | 16 (15%) | 4 (25.00%) | - | - | - | 12 (75.00%) | - |
| Zhou, Sun | 144 | 108 (75%) | 11 (10.19%) | - | - | - | - | 97 (89.81%) | 36 (25%) | 2 (5.56%) | - | - | - | - | 34 (94.44%) |
| Hippisley-Cox | - | NA | - | - | - | - | - | - | 1286 | 56 (4.35%) | 427 (33.20%) | - | 791 (61.51%) | - | 12 (0.93%) |
| Zhao, Chen | 641 | 398 (62%) | 87 (21.86%) | - | - | - | - | 311 (78.14%) | 195 (30%) | 52 (26.67%) | - | - | - | - | 143 (73.33%) |
| Qu | 246 | 226 (91%) | 90 (39.82%) | - | - | - | - | 136 (60.18%) | 20 (8%) | 14 (70.00%) | - | - | - | - | 6 (30.00%) |

Table 4. Disease severity by smoking status.




Figure 7. Forest plot for the risk of severe disease in current vs. never smokers.




Figure 8. Forest plot for the risk of severe disease in former vs. never smokers.

Accepted

| Recovered | | | | | | | | | Died | | | | | | |
|--------------|---------------------------|------------|--------------------|-------------------|---------------------------|------------------|--------------------------|----------------|------------|--------------------|-------------------|---------------------------|------------------|--------------------------|----------------|
| Author | Population with mortality | N (%) | Current smoker (%) | Former smoker (%) | Current/former smoker (%) | Never smoker (%) | Never/unknown smoker (%) | Not stated (%) | N (%) | Current smoker (%) | Former smoker (%) | Current/former smoker (%) | Never smoker (%) | Never/unknown smoker (%) | Not stated (%) |
| Chen | 274 | 161 (58%) | 5 (3.11%) | 5 (3.11%) | - | - | - | 151 (93.79%) | 113 (41%) | 7 (6.19%) | 2 (1.77%) | - | - | - | 104 (92.04%) |
| Zhou, Yu | 191 | 137 (71%) | 6 (4.38%) | - | - | - | - | 131 (95.62%) | 54 (28%) | 5 (9.26%) | - | - | - | - | 49 (90.74%) |
| Yang, Yu | 52 | 20 (38%) | 2 (10.00%) | - | - | - | 18 (90.00%) | - | 32 (61%) | - | - | - | - | 32 (100.00%) | - |
| Borobia | 2226 | 1766 (79%) | 113 (6.40%) | - | - | - | - | 1653 (93.60%) | 460 (20%) | 44 (9.57%) | - | - | - | - | 416 (90.43%) |
| Giacomelli | 233 | 185 (79%) | - | - | 53 (28.65%) | 132 (71.35%) | - | - | 48 (20%) | - | - | 17 (35.42%) | 31 (64.58%) | - | 0 (0.00%) |
| Yao | 108 | 96 (88%) | 1 (1.04%) | - | - | - | - | 95 (98.96%) | 12 (11%) | 3 (25.00%) | - | - | - | - | 9 (75.00%) |
| Carillo-Vega | 9946 | 8983 (90%) | 795 (8.85%) | - | - | - | - | 8188 (91.15%) | 963 (9%) | 99 (10.28%) | - | - | - | - | 864 (89.72%) |
| Heng | 51 | 39 (76%) | 6 (15.38%) | - | - | - | - | 33 (84.62%) | 12 (23%) | 1 (8.33%) | - | - | - | - | 11 (91.67%) |
| Chen, Jiang | 135 | NA (NA%) | - | - | - | - | - | - | 31 (22%) | - | - | 4 (12.90%) | - | - | 27 (87.10%) |
| Heili-Frades | 4712 | 4086 (86%) | 210 (5.14%) | 659 (16.13%) | - | - | 3217 (78.73%) | - | 626 (13%) | 23 (3.67%) | 161 (25.72%) | - | - | 442 (70.61%) | - |
| Kim, Garg | 2490 | 2070 (83%) | 128 (6.18%) | 481 (23.24%) | - | - | 1461 (70.58%) | - | 420 (16%) | 22 (5.24%) | 161 (38.33%) | - | - | 236 (56.19%) | - |
| Al-Hindawi | 31 | 15 (48%) | 0 (0.00%) | 10 (66.67%) | - | 5 (33.33%) | - | - | 16 (51%) | 1 (6.25%) | 12 (75.00%) | - | 3 (18.75%) | - | - |
| Louis | 22 | 16 (72%) | - | - | 7 (43.75%) | - | - | 9 (56.25%) | 6 (27%) | - | - | 3 (50.00%) | - | - | 3 (50.00%) |
| Soto-Mota | 400 | 200 (50%) | - | - | 23 (11.50%) | - | - | 177 (88.50%) | 200 (50%) | - | - | 25 (12.50%) | - | - | 175 (87.50%) |
| Garibaldi | 747 | 634 (84%) | 36 (5.68%) | 129 (20.35%) | - | - | - | 469 (73.97%) | 113 (15%) | 6 (5.31%) | 36 (31.86%) | - | - | - | 71 (62.83%) |
| Docherty | 13364 | 8199 (61%) | 370 (4.51%) | 1832 (22.34%) | - | 4179 (50.97%) | - | 1818 (22.17%) | 5165 (38%) | 214 (4.14%) | 1350 (26.14%) | - | 2105 (40.76%) | - | 1496 (28.96%) |

| | | | | | | | | | | | | | | | |
|-----------------------|-------|----------------|----------------|------------------|---------------|------------------|---------------|------------------|---------------|---------------|-----------------|--------------|-----------------|--------------|-----------------|
| Kuderer | 928 | 807 (86%) | 38 (4.71%) | 262 (32.47%) | - | 425 (52.66%) | - | 31 (3.84%) | 121 (13%) | 5 (4.13%) | 64 (52.89%) | - | 44 (36.36%) | - | 2 (1.65%) |
| Ramlall | 11116 | 10498 (94%) | - | - | 2771 (26.40%) | 7727 (73.60%) | - | - | 618 (5%) | - | - | 208 (33.66%) | 410 (66.34%) | - | - |
| Wang, Oekelen | 57 | 43 (75%) | - | - | 14 (32.56%) | - | - | 29 (67.44%) | 14 (24%) | - | - | 7 (50.00%) | - | - | 7 (50.00%) |
| Martinez- Portilla | 224 | 217 (96%) | - | - | 7 (3.23%) | - | - | 210 (96.77%) | 7 (3%) | - | - | 0 (0.00%) | - | - | 7 (100.00%) |
| Cen | 1007 | 964 (95%) | - | - | 87 (9.02%) | - | - | 877 (90.98%) | 43 (4%) | - | - | 1 (2.33%) | - | - | 42 (97.67%) |
| Klang | 3406 | 2270 (66%) | - | - | 492 (21.67%) | - | - | 1778 (78.33%) | 1136 (33%) | - | - | 301 (26.50%) | - | - | 835 (73.50%) |
| Wang, Zhong | 5510 | 4874 (88%) | 247 (5.07%) | 1083 (22.22%) | - | 3544 (72.71%) | - | - | 636 (11%) | 28 (4.40%) | 214 (33.65%) | - | 394 (61.95%) | - | - |
| Miyara | 338 | 211 (62%) | 13 (6.16%) | 58 (27.49%) | - | 141 (66.82%) | - | - | 46 (13%) | 1 (2.17%) | 23 (50.00%) | - | 21 (45.65%) | - | - |
| Rajter | 255 | 209 (81%) | - | - | 28 (13.40%) | 181 (86.60%) | - | - | 53 (20%) | - | - | 18 (33.96%) | 28 (52.83%) | - | - |
| Zeng | 1031 | 866 (84%) | - | - | 69 (7.97%) | - | - | 797 (92.03%) | 165 (16%) | - | - | 36 (21.82%) | - | - | 129 (78.18%) |
| Chen, Yu | 1859 | 1651 (88%) | 32 (1.94%) | 54 (3.27%) | - | 1565 (94.79%) | - | - | 208 (11%) | 13 (6.25%) | 12 (5.77%) | - | 183 (87.98%) | - | - |
| Garassino | 190 | 124 (65%) | - | - | 92 (74.19%) | 32 (25.81%) | - | - | 66 (34%) | - | 61 (92.42%) | - | 5 (7.58%) | - | - |
| Gu | 884 | 864 (97%) | 40 (4.63%) | 250 (28.94%) | - | 219 (25.35%) | - | - | 20 (2%) | 0 (0.00%) | 14 (70.00%) | - | 6 (30.00%) | - | - |
| Sigel | 88 | 70 (79%) | - | - | 37 (52.86%) | - | - | 33 (47.14%) | 18 (20%) | - | - | 11 (61.11%) | - | - | 7 (38.89%) |
| Nguyen | 356 | 308 (86%) | - | - | 91 (29.55%) | - | - | 217 (70.45%) | 45 (12%) | - | - | 23 (51.11%) | - | - | 22 (48.89%) |
| de Souza | 8443 | 7826 (92%) | - | - | 95 (1.21%) | - | 7571 (96.74%) | 160 (2.04%) | 617 (7%) | - | - | 47 (7.62%) | - | 560 (90.76%) | 10 (1.62%) |
| Mendy | 532 | 663 (124%) | - | - | 160 (24.13%) | - | - | 502 (75.72%) | 26 (4%) | - | - | 10 (38.46%) | - | - | 16 (61.54%) |
| Shi, Resurreccion | 256 | 210 (82%) | - | - | 128 (60.95%) | - | - | 82 (39.05%) | 46 (17%) | - | - | 26 (56.52%) | - | - | 20 (43.48%) |
| Xie | 619 | 591 (95%) | - | - | 43 (7.28%) | - | - | 548 (92.72%) | 28 (4%) | - | - | 8 (28.57%) | - | - | 20 (71.43%) |

| | | | | | | | | | | | | | | | |
|------------|-------|---------------|-----------------|-----------------|---------------|------------------|--------------|------------------|-----------------|----------------|------------------|--------------|------------------|--------------|-----------------|
| Fox | 54 | 35 (64%) | 1 (2.86%) | 4 (11.43%) | - | 18 (51.43%) | - | 12 (34.29%) | 19 (35%) | 0 (0.00%) | 2 (10.53%) | - | 12 (63.16%) | - | 5 (26.32%) |
| Zhang, Cao | 289 | 240 (83%) | 10 (4.17%) | 6 (2.50%) | - | - | - | 224 (93.33%) | 49 (16%) | 4 (8.16%) | 8 (16.33%) | - | - | - | 37 (75.51%) |
| Gupta | 496 | 255 (51%) | - | - | 15 (5.88%) | - | 80 (31.37%) | 160 (62.75%) | 241 (48%) | - | - | 21 (8.71%) | 77 (31.95%) | - | 143 (59.34%) |
| Soares | 1075 | 696 (64%) | 38 (5.46%) | - | - | - | 658 (94.54%) | - | 456 (42%) | 39 (8.55%) | - | - | - | 417 (91.45%) | - |
| Thompson | 470 | 301 (64%) | 39 (12.96%) | 79 (26.25%) | - | 183 (60.80%) | - | - | 169 (35%) | 27 (15.98%) | 49 (28.99%) | - | 93 (55.03%) | - | - |
| Bernaola | 1645 | 1382 (84%) | 35 (2.53%) | 146 (10.56%) | - | 1201 (86.90%) | - | - | 263 (15%) | 6 (2.28%) | 33 (12.55%) | - | 218 (82.89%) | - | - |
| Islam | 654 | 631 (96%) | 103 (16.32%) | - | - | - | - | 507 (80.35%) | 23 (3%) | 3 (13.04%) | - | - | - | - | - |
| Philipose | 466 | 267 (57%) | 19 (7.12%) | 204 (76.40%) | - | 44 (16.48%) | - | - | 199 (42%) | 9 (4.52%) | 137 (68.84%) | - | 33 (16.58%) | - | 20 (10.05%) |
| Dashti | 4140 | 3953 (95%) | - | - | 1068 (27.02%) | 2078 (52.57%) | - | 804 (20.34%) | 187 (4%) | - | - | 109 (58.29%) | 56 (29.95%) | - | 22 (11.76%) |
| Fillmore | 1794 | 1566 (87%) | 408 (26.05%) | 758 (48.40%) | - | 279 (17.82%) | - | 98 (6.26%) | 228 (12%) | 44 (19.30%) | 141 (61.84%) | - | 43 (18.86%) | - | 23 (10.09%) |
| Pan | 3536 | 3302 (93%) | - | - | 862 (26.11%) | - | - | 2440 (73.89%) | 234 (6%) | - | - | 82 (35.04%) | - | - | 152 (64.96%) |
| Zhao, Chen | 474 | 398 (83%) | 87 (21.86%) | - | - | - | - | 311 (78.14%) | 82 (17%) | 36 (43.90%) | - | - | - | - | 46 (56.10%) |
| Holman | 10989 | NA (NA%) | - | - | - | - | - | - | 10989 (100%) | 609 (5.54%) | 4684 (42.62%) | - | 5386 (49.01%) | - | 310 (2.82%) |
| Chand | 300 | 143 (47%) | 23 (16.08%) | - | - | - | - | 120 (83.92%) | 157 (52%) | 44 (28.03%) | - | - | - | - | 113 (71.97%) |

Note. Solis et al. and the OpenSAFELY Collaborative reported on mortality by smoking status in a multivariable analysis but did not present raw data for both the exposure and outcome variables.

Table 5. Mortality by smoking status.




Figure 9. Forest plot for the risk of mortality in current vs. never smokers.




Figure 10. Forest plot for the risk of mortality in former vs. never smokers.




Figure 11. A schematic of some of the interpretation issues for the association of smoking and SARS-CoV-2/COVID-19. * Indicates potential confounding with smoking status.