


# Daily Inspiration



Don't stop  
ignoring the data

I'm InspiroBot.

I am an artificial intelligence dedicated to generating unlimited amounts of unique inspirational quotes for endless enrichment of pointless human existence.



# Today

---

- Recap yesterday
- Logistic regression: using regression tools for classification
- Neural network basics



# Yesterday

---

- Cost function: (differentiable) function that shows how wrong an estimate is for given parameters.
- Gradient descent: one common way to minimise the cost function automatically, i.e. to get optimal parameters
- Linear regression: very simple model that assumes that value to predict is linear combination of input features.
- Overfitting and underfitting, bias and variance: want our model to work well for unseen data. Need just enough model freedom given the complexity of our problem. How:
  - Cross-validation to measure ability to generalise + get best hyperparameters
  - Use learning curves to diagnose bias vs. variance

# Gradient descent in linear algebra

---

- Goal gradient descent: take a small step in every parameter such that you get closer to the minimum of the cost. Return new theta's.

$$\theta_{0_{new}} = \theta_0 - \frac{\alpha}{m} \sum_{i=1}^m ((h_{\theta}(x^{(i)}) - y^{(i)}) \cdot 1)$$

$$\theta_{1_{new}} = \theta_1 - \frac{\alpha}{m} \sum_{i=1}^m ((h_{\theta}(x^{(i)}) - y^{(i)}) \cdot x_1^{(i)})$$

$$\theta_{2_{new}} = \theta_2 - \frac{\alpha}{m} \sum_{i=1}^m ((h_{\theta}(x^{(i)}) - y^{(i)}) \cdot x_2^{(i)})$$

# Gradient descent in linear algebra

We have data, known values, and initial theta's:

$$X = \begin{bmatrix} 1 & feat_1val_1 & feat_2val_1 \\ 1 & feat_1val_2 & feat_2val_2 \end{bmatrix}; y = \begin{bmatrix} 10.23 \\ -4 \end{bmatrix}; params = \begin{bmatrix} \theta_0 \\ \theta_1 \\ \theta_2 \end{bmatrix}$$

Get predicted values:

$$\begin{bmatrix} 1 & feat_1val_1 & feat_2val_1 \\ 1 & feat_1val_2 & feat_2val_2 \end{bmatrix} @ \begin{bmatrix} \theta_0 \\ \theta_1 \\ \theta_2 \end{bmatrix} = \begin{bmatrix} 9.23 \\ -2.5 \end{bmatrix}$$

2 by 3 times 3 by 1 gives 2 by 1 (rows by columns)

Get errors:

$$errs = \begin{bmatrix} 9.23 \\ -2.5 \end{bmatrix} - y = \begin{bmatrix} 9.23 \\ -2.5 \end{bmatrix} - \begin{bmatrix} 10.23 \\ -4 \end{bmatrix} = \begin{bmatrix} -1 \\ 1.5 \end{bmatrix}$$



# Gradient descent in linear algebra

---

$$errs = \begin{bmatrix} -1 \\ 1.5 \end{bmatrix}$$

$$\theta_{0_{new}} = \theta_0 - \frac{\alpha}{m} \sum_{i=1}^m ((h_{\theta}(x^{(i)}) - y^{(i)}) \cdot 1)$$

$$\theta_{1_{new}} = \theta_1 - \frac{\alpha}{m} \sum_{i=1}^m ((h_{\theta}(x^{(i)}) - y^{(i)}) \cdot x_1^{(i)})$$

$$\theta_{2_{new}} = \theta_2 - \frac{\alpha}{m} \sum_{i=1}^m ((h_{\theta}(x^{(i)}) - y^{(i)}) \cdot x_2^{(i)})$$

# Gradient descent in linear algebra

Calculate, for each feature, sum of each error times that feature:

$$\begin{bmatrix} -1 \\ 1.5 \end{bmatrix}^T = [-1 \quad 1.5]$$

$$errs = \begin{bmatrix} -1 \\ 1.5 \end{bmatrix}$$

$$[-1 \quad 1.5] @ \begin{bmatrix} 1 & feat_1 val_1 & feat_2 val_1 \\ 1 & feat_1 val_2 & feat_2 val_2 \end{bmatrix} =$$

$$[-1 \cdot 1 + 1.5 \cdot 1 \quad -1 \cdot feat_1 val_1 + 1.5 \cdot feat_1 val_2 \quad -1 \cdot feat_2 val_1 + 1.5 \cdot feat_2 val_2]$$

# Gradient descent in linear algebra

Calculate, for each feature, sum of each error times that feature:

$$\begin{bmatrix} -1 \\ 1.5 \end{bmatrix}^T = \begin{bmatrix} -1 & 1.5 \end{bmatrix}$$

$$errs = \begin{bmatrix} -1 \\ 1.5 \end{bmatrix}$$

$$\begin{bmatrix} -1 & 1.5 \end{bmatrix} @ \begin{bmatrix} 1 & feat_1 val_1 & feat_2 val_1 \\ 1 & feat_1 val_2 & feat_2 val_2 \end{bmatrix} =$$

$$[-1 \cdot 1 + 1.5 \cdot 1]$$

$$[-1 \cdot feat_1 val_1 + 1.5 \cdot feat_1 val_2]$$

$$[-1 \cdot feat_2 val_1 + 1.5 \cdot feat_2 val_2]$$

$$\theta_{0new} = \theta_0 - \frac{\alpha}{m} \sum_{i=1}^m ((h_{\theta}(x^{(i)}) - y^{(i)}) \cdot 1)$$

$$\theta_{1new} = \theta_1 - \frac{\alpha}{m} \sum_{i=1}^m ((h_{\theta}(x^{(i)}) - y^{(i)}) \cdot x_1^{(i)})$$

$$\theta_{2new} = \theta_2 - \frac{\alpha}{m} \sum_{i=1}^m ((h_{\theta}(x^{(i)}) - y^{(i)}) \cdot x_2^{(i)})$$



# Gradient descent in linear algebra

Now all that we need to do is multiply with  $\alpha/m$  and subtract from our old theta's:

$$\begin{aligned} & \alpha/m \cdot \begin{bmatrix} -1 \cdot 1 + 1.5 \cdot 1 & -1 \cdot feat_1 val_1 + 1.5 \cdot feat_1 val_2 & -1 \cdot feat_2 val_1 + 1.5 \cdot feat_2 val_2 \end{bmatrix} \\ &= \begin{bmatrix} \frac{\alpha}{m}(-1 \cdot 1 + 1.5 \cdot 1) & \frac{\alpha}{m}(-1 \cdot feat_1 val_1 + 1.5 \cdot feat_1 val_2) & \frac{\alpha}{m}(-1 \cdot feat_2 val_1 + 1.5 \cdot feat_2 val_2) \end{bmatrix} \end{aligned}$$

Transpose it:

$$\begin{bmatrix} \frac{\alpha}{m}(-1 \cdot 1 + 1.5 \cdot 1) & \frac{\alpha}{m}(-1 \cdot feat_1 val_1 + 1.5 \cdot feat_1 val_2) & \frac{\alpha}{m}(-1 \cdot feat_2 val_1 + 1.5 \cdot feat_2 val_2) \end{bmatrix}^T = \begin{bmatrix} \frac{\alpha}{m}(-1 \cdot 1 + 1.5 \cdot 1) \\ \frac{\alpha}{m}(-1 \cdot feat_1 val_1 + 1.5 \cdot feat_1 val_2) \\ \frac{\alpha}{m}(-1 \cdot feat_2 val_1 + 1.5 \cdot feat_2 val_2) \end{bmatrix}$$

So finally:

$$\begin{bmatrix} \theta_{0old} \\ \theta_{1old} \\ \theta_{2old} \end{bmatrix} - \begin{bmatrix} \frac{\alpha}{m}(-1 \cdot 1 + 1.5 \cdot 1) \\ \frac{\alpha}{m}(-1 \cdot feat_1 val_1 + 1.5 \cdot feat_1 val_2) \\ \frac{\alpha}{m}(-1 \cdot feat_2 val_1 + 1.5 \cdot feat_2 val_2) \end{bmatrix} = \begin{bmatrix} \theta_{0new} \\ \theta_{1new} \\ \theta_{2new} \end{bmatrix}$$

$$\theta_{1new} = \theta_{1old} - \frac{\alpha}{m} \sum_{i=1}^m ((h_{\theta}(x^{(i)}) - y^{(i)}) \cdot x_1^{(i)})$$


# Logistic regression

---

- Use regression-like framework for classification

# Logistic regression


- Naïve idea:  
Train a linear regression. If  
Class  $\geq 0.5$ , predict class 1.  
Otherwise, class 0.






# Logistic regression

- Naïve idea:  
Train a linear regression. If  
Class  $\geq 0.5$ , predict class 1.  
Otherwise, class 0.




# Logistic regression

- Naïve idea:  
Train a linear regression. If  $\text{Class} \geq 0.5$ , predict class 1. Otherwise, class 0.
- Problems:
  - You can predict class  $> 1$  and  $< 0$ , while that is not possible in reality.



# Logistic regression


- Naïve idea:  
Train a linear regression. If  $\text{Class} \geq 0.5$ , predict class 1. Otherwise, class 0.
- Problems:
  - You can predict class  $> 1$  and  $< 0$ , while that is not possible in reality.
  - This example seemed to work, but quickly breaks down  $\rightarrow$





# Logistic regression

- Naïve idea:  
Train a linear regression. If  $\text{Class} \geq 0.5$ , predict class 1. Otherwise, class 0.
- Problems:
  - You can predict class  $> 1$  and  $< 0$ , while that is not possible in reality.
  - This example seemed to work, but quickly breaks down  $\rightarrow$  get what is basically confirmation of hypothesis, but perform worse!



# Logistic regression

---

- What we want:
  - Use the information that we only have two classes, 0 or 1.
  - Hypothesis function should output only numbers between 0 or 1.

# Sigmoid or logistic function

---

- Before, our hypothesis function was of the form:

$$h_{\theta}(x) = \theta^T \cdot x$$



# Sigmoid or logistic function

---

- Before, our hypothesis function was of the form:

$$h_{\theta}(x) = \theta^T \cdot x \longrightarrow [0.5 \quad 3 \quad -1.5] \cdot \begin{bmatrix} 1 \\ 3 \\ 8 \end{bmatrix}$$

# Sigmoid or logistic function

- Before, our hypothesis function was of the form:

$$h_{\theta}(x) = \theta^T \cdot x \quad \longrightarrow \quad \underbrace{[0.5 \quad 3 \quad -1.5]}_{\text{Learned parameters (theta 0 – theta 2)}} \cdot \underbrace{\begin{bmatrix} 1 \\ 3 \\ 8 \end{bmatrix}}_{\text{Features for one sample (x0 = 1, intercept term, 2 data-derived features x1 and x2)}}$$



# Sigmoid or logistic function

---

- Before, our hypothesis function was of the form:

$$h_{\theta}(x) = \theta^T \cdot x \quad \longrightarrow \quad [0.5 \quad 3 \quad -1.5] \cdot \begin{bmatrix} 1 \\ 3 \\ 8 \end{bmatrix} = 0.5 \cdot 1 + 3 \cdot 3 - 1.5 \cdot 8 = -2.5$$



# Sigmoid or logistic function

---

- Before, our hypothesis function was of the form:

$$h_{\theta}(x) = \theta^T \cdot x$$

- Change that to the following:

$$h_{\theta}(x) = g(\theta^T \cdot x)$$

# Sigmoid or logistic function

---

- Before, our hypothesis function was of the form:

$$h_{\theta}(x) = \theta^T \cdot x$$

- Change that to the following:

$$h_{\theta}(x) = g(\theta^T \cdot x) \longrightarrow g(z) = \frac{1}{1 + e^{-z}}$$



# Sigmoid or logistic function

---

- Before, our hypothesis function was of the form:

$$h_{\theta}(x) = \theta^T \cdot x$$

- Change that to the following:

$$h_{\theta}(x) = g(\theta^T \cdot x) \longrightarrow g(z) = \frac{1}{1 + e^{-z}}$$

- What does that look like?



# Sigmoid or logistic function

---

- Before, our hypothesis function was of the form:

$$h_{\theta}(x) = \theta^T \cdot x$$


- Change that to the following:

$$h_{\theta}(x) = g(\theta^T \cdot x) \longrightarrow g(z) = \frac{1}{1 + e^{-z}}$$

- What does that look like?  $z \rightarrow \infty, e^{-z} \rightarrow 0$


$$z \rightarrow -\infty, e^{-z} \rightarrow \infty$$

# What does the sigmoid function look like?






# What does the sigmoid function look like?






# Sigmoid or logistic function

- How do we work with this?  $h_{\theta}(x) = \frac{1}{1 + e^{-(\theta^T \cdot x)}}$



# Sigmoid or logistic function

- How do we work with this?  $h_{\theta}(x) = \frac{1}{1 + e^{-(\theta^T \cdot x)}}$ 
  - Interpret outcome of  $h_{\theta}(x)$  as probability that class = 1 given the features.






# Sigmoid or logistic function

- How do we work with this?  $h_{\theta}(x) = \frac{1}{1 + e^{-(\theta^T \cdot x)}}$

- Interpret outcome of  $h_{\theta}(x)$  as probability that class = 1 given the features. Example:

$$x = \begin{bmatrix} x_0 \\ x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 1 \\ \text{Tumor size} \\ \text{Neovascularisation level} \end{bmatrix}$$


$h_{\theta}(x) = 0.8 \longrightarrow$  80% chance of tumor being malignant





# Sigmoid or logistic function

- How do we work with this?  $h_{\theta}(x) = \frac{1}{1 + e^{-(\theta^T \cdot x)}}$



- Interpret outcome of  $h_{\theta}(x)$  as probability that class = 1 given the features. Example:

$$x = \begin{bmatrix} x_0 \\ x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 1 \\ \text{Tumor size} \\ \text{Neovascularisation level} \end{bmatrix}$$

$h_{\theta}(x) = 0.8 \longrightarrow$  80% chance of tumor being malignant (class 1)  
100% - 80%  $\rightarrow$  20 % chance of being benign (class 0)


# Sigmoid or logistic function

▪ How do we work with this?  $h_{\theta}(x) = \frac{1}{1 + e^{-(\theta^T \cdot x)}}$

- Interpret outcome of  $h_{\theta}(x)$  as probability that class = 1 given the features.
- Formally:

$$h_{\theta}(x) = \frac{1}{1 + e^{-(\theta^T \cdot x)}} = p(y=1|x;\theta)$$


$$p(y=0|x;\theta) = 1 - h_{\theta}(x)$$





# Decision boundary


- Threshold:






# Decision boundary

- Threshold:




# Decision boundary

- How does it look?  $g(z) = \frac{1}{1 + e^{-z}}$   
 $h_{\theta}(x) = g(\theta_0 \cdot x_0, \theta_1 \cdot x_1, \theta_2 \cdot x_2)$



# Decision boundary


- How does it look?  $g(z) = \frac{1}{1 + e^{-z}}$   
 $h_{\theta}(x) = g(\theta_0 \cdot x_0, \theta_1 \cdot x_1, \theta_2 \cdot x_2)$   
 $\theta = \begin{bmatrix} -3 \\ 1 \\ 1 \end{bmatrix}$







# Decision boundary

- How does it look?  $g(z) = \frac{1}{1 + e^{-z}}$   
 $h_{\theta}(x) = g(\theta_0 \cdot x_0, \theta_1 \cdot x_1, \theta_2 \cdot x_2)$   
 $\theta = \begin{bmatrix} -3 \\ 1 \\ 1 \end{bmatrix}$   
 $y = 1$  if  $-3 \cdot 1 + 1 \cdot x_1 + 1 \cdot x_2 \geq 0$




# Decision boundary

- How does it look?  $g(z) = \frac{1}{1 + e^{-z}}$



# Decision boundary


- How does it look?  $g(z) = \frac{1}{1 + e^{-z}}$   
 $h_{\theta}(x) = g(\theta_0 \cdot x_0, \theta_1 \cdot x_1, \theta_2 \cdot x_2)$   
 $\theta = \begin{bmatrix} -3 \\ 1 \\ 1 \end{bmatrix}$   
 $y = 1$  if  $-3 \cdot 1 + 1 \cdot x_1 + 1 \cdot x_2 \geq 0$   
 $-3 \cdot 1 + \cancel{1 \cdot x_1} + \cancel{1 \cdot x_2} \geq 0$   
 $x_1 + x_2 = 3$






# Decision boundary

- How does it look?  $g(z) = \frac{1}{1 + e^{-z}}$   
 $h_{\theta}(x) = g(\theta_0 \cdot x_0, \theta_1 \cdot x_1, \theta_2 \cdot x_2)$   
 $\theta = \begin{bmatrix} -3 \\ 1 \\ 1 \end{bmatrix}$   
 $y = 1$  if  $-3 \cdot 1 + 1 \cdot x_1 + 1 \cdot x_2 \geq 0$   
 $-3 \cdot 1 + \cancel{1 \cdot x_1} + \cancel{1 \cdot x_2} \geq 0$   
 $x_1 + x_2 = 3$




# Decision boundary

- How does it look?  $g(z) = \frac{1}{1 + e^{-z}}$   
 $h_{\theta}(x) = g(\theta_0 \cdot x_0, \theta_1 \cdot x_1, \theta_2 \cdot x_2)$   
 $\theta = \begin{bmatrix} -3 \\ 1 \\ 1 \end{bmatrix}$   
 $y = 1$  if  $-3 \cdot 1 + 1 \cdot x_1 + 1 \cdot x_2 \geq 0$



# Decision boundary


- How does it look?  $g(z) = \frac{1}{1 + e^{-z}}$   
 $h_{\theta}(x) = g(\theta_0 \cdot x_0, \theta_1 \cdot x_1, \theta_2 \cdot x_2)$   
 $\theta = \begin{bmatrix} -3 \\ 1 \\ 1 \end{bmatrix}$   
 $y = 1$  if  $-3 \cdot 1 + 1 \cdot x_1 + 1 \cdot x_2 \geq 0$





# Decision boundary


- How does it look?  $g(z) = \frac{1}{1 + e^{-z}}$   
 $h_{\theta}(x) = g(\theta_0 \cdot x_0, \theta_1 \cdot x_1, \theta_2 \cdot x_2)$   
 $\theta = \begin{bmatrix} -3 \\ 1 \\ 1 \end{bmatrix}$



$$y=1 \text{ if } -3 \cdot 1 + 1 \cdot x_1 + 1 \cdot x_2 \geq 0$$

# Non-linear decision boundary

- How does it look?  $g(z) = \frac{1}{1 + e^{-z}}$   
 $h_{\theta}(x) = g(\theta_0 x_0, \theta_1 x_1, \theta_2 x_2, \theta_3 x_1^2, \theta_4 x_2^2)$
- Add two polynomial features






# Non-linear decision boundary

- How does it look?  $g(z) = \frac{1}{1 + e^{-z}}$   
 $h_{\theta}(x) = g(\theta_0 x_0, \theta_1 x_1, \theta_2 x_2, \theta_3 x_1^2, \theta_4 x_2^2)$
- Add two polynomial features

$$\theta = \begin{bmatrix} -1 \\ 0 \\ 0 \\ 1 \\ 1 \end{bmatrix}$$






# Non-linear decision boundary

- How does it look?  $g(z) = \frac{1}{1 + e^{-z}}$   
 $h_{\theta}(x) = g(\theta_0 x_0, \theta_1 x_1, \theta_2 x_2, \theta_3 x_1^2, \theta_4 x_2^2)$
- Add two polynomial features


$$\theta = \begin{bmatrix} -1 \\ 0 \\ 0 \\ 1 \\ 1 \end{bmatrix} \longrightarrow -1 + x_1^2 + x_2^2 \geq 0$$



# Non-linear decision boundary

- How does it look?  $g(z) = \frac{1}{1 + e^{-z}}$   
 $h_{\theta}(x) = g(\theta_0 x_0, \theta_1 x_1, \theta_2 x_2, \theta_3 x_1^2, \theta_4 x_2^2)$
- Add two polynomial features

$$\theta = \begin{bmatrix} -1 \\ 0 \\ 0 \\ 1 \\ 1 \end{bmatrix} \longrightarrow -1 + x_1^2 + x_2^2 \geq 0$$
$$\downarrow$$
$$x_1^2 + x_2^2 \geq 1$$






# Non-linear decision boundary

- How does it look?  $g(z) = \frac{1}{1 + e^{-z}}$   
 $h_{\theta}(x) = g(\theta_0 x_0, \theta_1 x_1, \theta_2 x_2, \theta_3 x_1^2, \theta_4 x_2^2)$
- Add two polynomial features




$$\theta = \begin{bmatrix} -1 \\ 0 \\ 0 \\ 1 \\ 1 \end{bmatrix} \longrightarrow -1 + x_1^2 + x_2^2 \geq 0$$
$$\downarrow$$
$$x_1^2 + x_2^2 \geq 1$$





# Non-linear decision boundary

- How does it look?
- If you add more and higher-order polynomial features, you can get complex boundaries:



# So how do we get theta's?

---

- Need a cost function

# So how do we get theta's?

---

- Need a cost function

- Before:  $J(\theta) = \frac{1}{2m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2$



# So how do we get theta's?

---

- Need a cost function

- Before:  $J(\theta) = \frac{1}{2m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2$




$$J(\theta) = \frac{1}{m} \sum_{i=1}^m \frac{1}{2} (h_{\theta}(x^{(i)}) - y^{(i)})^2$$

# So how do we get theta's?

---

- Need a cost function

- Before:  $J(\theta) = \frac{1}{2m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2$

$$J(\theta) = \frac{1}{m} \sum_{i=1}^m \left( \frac{1}{2} (h_{\theta}(x^{(i)}) - y^{(i)})^2 \right)$$


$$\text{Cost}(x) = \frac{1}{2} (h_{\theta}(x) - y)^2$$



# So how do we get theta's?

- Need a cost function

- Before:  $J(\theta) = \frac{1}{2m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2$

$$\begin{array}{l} \updownarrow \\ J(\theta) = \frac{1}{m} \sum_{i=1}^m \left\{ \frac{1}{2} (h_{\theta}(x^{(i)}) - y^{(i)})^2 \right\} \\ \text{Cost}(x) = \frac{1}{2} (h_{\theta}(x) - y)^2 \end{array} \left. \vphantom{\begin{array}{l} J(\theta) = \frac{1}{m} \sum_{i=1}^m \left\{ \frac{1}{2} (h_{\theta}(x^{(i)}) - y^{(i)})^2 \right\} \\ \text{Cost}(x) = \frac{1}{2} (h_{\theta}(x) - y)^2 \end{array}} \right\} J(\theta) = \frac{1}{m} \sum_{i=1}^m \text{Cost}(x^{(i)})$$





# So how do we get theta's?

---

- Need a cost function  $J(\theta) = \frac{1}{m} \sum_{i=1}^m \text{Cost}(x^i)$        $\text{Cost}(x) = \frac{1}{2} (h_{\theta}(x) - y)^2$
- Why not MSE? → not convex

# So how do we get theta's?



- Need a cost function  $J(\theta) = \frac{1}{m} \sum_{i=1}^m \text{Cost}(x^i)$        $\text{Cost}(x) = \frac{1}{2} (h_{\theta}(x) - y)^2$
- Why not MSE? → not convex





# So how do we get theta's?



- Need a cost function  $J(\theta) = \frac{1}{m} \sum_{i=1}^m \text{Cost}(x^i)$        $\text{Cost}(x) = \frac{1}{2} (h_{\theta}(x) - y)^2$
- Why not MSE? → not convex







# So how do we get theta's?

- Need a cost function  $J(\theta) = \frac{1}{m} \sum_{i=1}^m \text{Cost}(x^i)$        $\text{Cost}(x) = \frac{1}{2} (h_{\theta}(x) - y)^2$
- Why not MSE? → not convex



# So how do we get theta's?



- Need a cost function  $J(\theta) = \frac{1}{m} \sum_{i=1}^m \text{Cost}(x^i)$        $\text{Cost}(x) = \frac{1}{2} (h_{\theta}(x) - y)^2$
- Why not MSE? → not convex





# So how do we get theta's?



- Need a cost function  $J(\theta) = \frac{1}{m} \sum_{i=1}^m \text{Cost}(x^i)$        $\text{Cost}(x) = \frac{1}{2} (h_{\theta}(x) - y)^2$
- Why not MSE? → not convex





# So how do we get theta's?

- Need a cost function  $J(\theta) = \frac{1}{m} \sum_{i=1}^m \text{Cost}(x^i)$        $\text{Cost}(x) = \frac{1}{2} (h_{\theta}(x) - y)^2$
- Why not MSE?  $\rightarrow$  not convex



# So how do we get theta's?

---

- Need a cost function  $J(\theta) = \frac{1}{m} \sum_{i=1}^m \text{Cost}(x^i)$
  - What then?
- ~~$\text{Cost}(x) = \frac{1}{2} (h_{\theta}(x) - y)^2$~~



# So how do we get theta's?

- Need a cost function  $J(\theta) = \frac{1}{m} \sum_{i=1}^m \frac{1}{2} \text{Cost}(x^i)$   ~~$\text{Cost}(x) = \frac{1}{2} (h_{\theta}(x) - y)^2$~~
- What then?


$$\text{Cost}(x) = \begin{cases} -\log(h_{\theta}(x)) & \text{if } y=1 \\ -\log(1 - h_{\theta}(x)) & \text{if } y=0 \end{cases}$$



# What does this function look like?

$$\text{Cost}(x) = \begin{cases} -\log(h_{\theta}(x)) & \text{if } y=1 \\ -\log(1-h_{\theta}(x)) & \text{if } y=0 \end{cases}$$


$$J(\theta) = \frac{1}{m} \sum_{i=1}^m \text{Cost}(x^i)$$



# What does this function look like?

$$\text{Cost}(x) = \begin{cases} -\log(h_{\theta}(x)) & \text{if } y=1 \\ -\log(1-h_{\theta}(x)) & \text{if } y=0 \end{cases}$$


$$J(\theta) = \frac{1}{m} \sum_{i=1}^m \text{Cost}(x^i)$$



# What does this function look like?

$$\text{Cost}(x) = \begin{cases} -\log(h_{\theta}(x)) & \text{if } y=1 \\ -\log(1-h_{\theta}(x)) & \text{if } y=0 \end{cases}$$

$$J(\theta) = \frac{1}{m} \sum_{i=1}^m \text{Cost}(x^i)$$






# What does this function look like?

$$\text{Cost}(x) = \begin{cases} -\log(h_{\theta}(x)) & \text{if } y=1 \\ -\log(1-h_{\theta}(x)) & \text{if } y=0 \end{cases}$$


$$J(\theta) = \frac{1}{m} \sum_{i=1}^m \text{Cost}(x^i)$$



# What does this function look like?

$$\text{Cost}(x) = \begin{cases} -\log(h_\theta(x)) & \text{if } y=1 \\ -\log(1-h_\theta(x)) & \text{if } y=0 \end{cases}$$


$$J(\theta) = \frac{1}{m} \sum_{i=1}^m \text{Cost}(x^i)$$



# What does this function look like?

$$\text{Cost}(x) = \begin{cases} -\log(h_\theta(x)) & \text{if } y=1 \\ -\log(1-h_\theta(x)) & \text{if } y=0 \end{cases}$$

$$J(\theta) = \frac{1}{m} \sum_{i=1}^m \text{Cost}(x^i)$$






# What does this function look like?

$$\text{Cost}(x) = \begin{cases} -\log(h_{\theta}(x)) & \text{if } y=1 \\ -\log(1-h_{\theta}(x)) & \text{if } y=0 \end{cases}$$

$$J(\theta) = \frac{1}{m} \sum_{i=1}^m \text{Cost}(x^i)$$



# Simplified notation

---

$$\text{Cost}(x) = \begin{cases} -\log(h_\theta(x)) & \text{if } y=1 \\ -\log(1-h_\theta(x)) & \text{if } y=0 \end{cases}$$



$$\text{Cost}(x) = -y \cdot \log(h_\theta(x)) - (1-y) \cdot \log(1-h_\theta(x))$$

$$J(\theta) = \frac{1}{m} \sum_{i=1}^m \text{Cost}(x^i)$$

# Simplified notation

---

$$\text{Cost}(x) = \begin{cases} -\log(h_\theta(x)) & \text{if } y=1 \\ -\log(1-h_\theta(x)) & \text{if } y=0 \end{cases}$$

$$J(\theta) = \frac{1}{m} \sum_{i=1}^m \text{Cost}(x^i)$$

↓

$$\text{Cost}(x) = -y \cdot \log(h_\theta(x)) - (1-y) \cdot \log(1-h_\theta(x))$$

↓  $y=0$

$$-0 \cdot \log(h_\theta(x)) - (1-0) \cdot \log(1-h_\theta(x))$$



# Simplified notation

$$\text{Cost}(x) = \begin{cases} -\log(h_\theta(x)) & \text{if } y=1 \\ -\log(1-h_\theta(x)) & \text{if } y=0 \end{cases}$$

$$J(\theta) = \frac{1}{m} \sum_{i=1}^m \text{Cost}(x^i)$$

$$\text{Cost}(x) = -y \cdot \log(h_\theta(x)) - (1-y) \cdot \log(1-h_\theta(x))$$

$$y = 0$$

$$-0 \cdot \log(h_\theta(x)) - (1-0) \cdot \log(1-h_\theta(x))$$

$$-\log(1-h_\theta(x))$$

# Simplified notation

$$\text{Cost}(x) = \begin{cases} -\log(h_\theta(x)) & \text{if } y=1 \\ -\log(1-h_\theta(x)) & \text{if } y=0 \end{cases}$$

$$J(\theta) = \frac{1}{m} \sum_{i=1}^m \text{Cost}(x^i)$$

$$\text{Cost}(x) = -y \cdot \log(h_\theta(x)) - (1-y) \cdot \log(1-h_\theta(x))$$

$$y = 1$$

$$-1 \cdot \log(h_\theta(x)) - (1-1) \cdot \log(1-h_\theta(x))$$

$$-\log(h_\theta(x))$$



# Putting it all together

---

$$\text{Cost}(x) = -y \cdot \log(h_\theta(x)) - (1-y) \cdot \log(1-h_\theta(x)) \quad J(\theta) = \frac{1}{m} \sum_{i=1}^m \text{Cost}(x^{(i)})$$

$$J(\theta) = \frac{1}{m} \sum_{i=1}^m -y^{(i)} \cdot \log(h_\theta(x^{(i)})) - (1-y^{(i)}) \cdot \log(1-h_\theta(x^{(i)}))$$



# Putting it all together

$$\text{Cost}(x) = -y \cdot \log(h_\theta(x)) - (1-y) \cdot \log(1-h_\theta(x)) \quad J(\theta) = \frac{1}{m} \sum_{i=1}^m \text{Cost}(x^i)$$

$$J(\theta) = \frac{1}{m} \sum_{i=1}^m \left[ -y^{(i)} \cdot \log(h_\theta(x^{(i)})) - (1-y^{(i)}) \cdot \log(1-h_\theta(x^{(i)})) \right]$$



$$J(\theta) = -\frac{1}{m} \sum_{i=1}^m \left[ y^{(i)} \cdot \log(h_\theta(x^{(i)})) + (1-y^{(i)}) \cdot \log(1-h_\theta(x^{(i)})) \right]$$

# Optimising the cost function

- Same form as for linear regression (only hypothesis function differs!)

$$\frac{\partial}{\partial \theta_j} J(\theta) = \frac{1}{m} \sum_{i=1}^m ((h_{\theta}(x^{(i)}) - y^{(i)}) \cdot x_j^{(i)})$$

$$\theta_j := \theta_j - \frac{\alpha}{m} \sum_{i=1}^m ((h_{\theta}(x^{(i)}) - y^{(i)}) \cdot x_j^{(i)})$$

$$h_{\theta}(x^{(i)}) = \frac{1}{1 + e^{-\theta^T * x}}$$



# Summary

---

- By using the sigmoid function as a transformation of normal regression and interpreting the output as a chance of being 0 or 1 we can do classification.
- Only the form of our hypothesis function is different
- Need a different cost function: should be smooth, and give logical values for large errors.



# Break for practical

---