

## **Week 3 – Tuesday session**

# **Stratified Case-control Analysis**

EPI202 – Epidemiologic Methods II

Murray A. Mittleman, MD, DrPH

Department of Epidemiology, Harvard TH Chan School of Public Health



**HARVARD T.H. CHAN**  
**SCHOOL OF PUBLIC HEALTH**

# Week 3: Discussion Topics

## 1. Stratified case-control data data

- Hypothesis tests
- Point and interval estimates – weighted averages
- Assumption of homogeneity
- Effect measure modification

## 2. Matching


- Matched cohort design
- Matched case-control design
- Appropriate, unnecessary and overmatching
- Matching ratio – impact on precision
- Matched case-control analysis
  - McNemar test and odds ratio estimator
  - Relationship to Mantel-Haenszel methods

# Matching Ratios

## Relative Efficiency of the Test of the Null Hypothesis

$H_A: RR =$	$Pr(A=1   Y=0)$	Number of Controls/Case						
		1	2	4	8	16	32	$\infty$
1	0-100%	50%	66%	80%	88%	95%	97%	100%
2	5%	30%	53%	71%	83%	92%	95%	100%
5	5%	20%	35%	54%	73%	84%	93%	100%

- Relative efficiency is the ratio of the variance of a particular design relative to having infinite controls (equivalent to conducting the full cohort analysis)
- Under the null, the efficiency of a particular matching ratio relative to the full cohort design is proportional to  $m/(m+1)$ , where  $m$  is the ratio of controls:cases.
- $Pr(A=1 | Y=0)$  is the exposure prevalence in the controls



# Hypothesis Test for Stratified Data (1)

- We first stratify the data on all confounding variables to form  $I$  strata and then calculate the **Mantel-Haenszel  $\chi^2$  test statistic**:

$$Z^2 = \frac{\left[ \sum_{i=1}^I X_i - \sum_{i=1}^I E_i(X_i | H_0) \right]^2}{\sum_{i=1}^I \text{Var}_i(X_i | H_0)} \sim \chi_1^2$$

# Summary Odds Ratio

## Point Estimate

- To calculate the estimate of the summary odds ratio, we use a weighted sum of the stratum-specific estimates:

$$\hat{OR} = \frac{\sum_{i=1}^I w'_i \hat{OR}_i}{\sum w'_i} = \sum_{i=1}^I w_i \hat{OR}_i$$

$$\text{Where } w_i = \frac{w'_i}{\sum w'_i} \quad \text{and} \quad \sum w_i = 1$$

- The exact value of the summary odds ratio depends upon the chosen weights.
- In the absence of effect measure modification, any weights will yield an unbiased estimate of the common odds ratio
- Weights are chosen to optimize the statistical precision of the estimate

# Mantel-Haenszel Estimator of the Odds Ratio

- Computational formula for the **Mantel-Haenszel summary odds ratio**:

$$\hat{OR}_{MH} = \frac{\sum_{i=1}^I w'_i \hat{OR}_i}{\sum_{i=1}^I w'_i} = \frac{\sum_{i=1}^I \frac{b_i c_i}{T_i} \frac{a_i d_i}{b_i c_i}}{\sum_{i=1}^I \frac{b_i c_i}{T_i}} = \frac{\sum_{i=1}^I \frac{a_i d_i}{T_i}}{\sum_{i=1}^I \frac{b_i c_i}{T_i}}$$

- A given stratum may contribute to the numerator, the denominator, both or neither

## 95% CI for $\ln(\text{OR}_{\text{MH}})$

$$X \pm Z_{1-\alpha/2} \sqrt{\hat{\text{Var}}(X)}$$

- Where:

- $X = \ln(\hat{\text{OR}}_{\text{MH}})$
- $Z_{1-\alpha/2} = 100(1-\alpha/2)^{\text{th}}$  percentile of the standard normal distribution
- $\hat{\text{Var}}(X) = \text{estimated variance of } \ln(\hat{\text{OR}}_{\text{MH}})$

**True or False? The Mantel-Haenszel odds ratio will always be a value between the stratum-specific odds ratio estimates.**

$$\hat{OR}_{MH} = \frac{\sum_{i=1}^I \frac{a_i d_i}{T_i}}{\sum_{i=1}^I \frac{b_i c_i}{T_i}} = \frac{\frac{4 * 224}{292} + \frac{9 * 390}{444} + \frac{4 * 330}{393} + \frac{6 * 362}{442}}{\frac{2 * 62}{292} + \frac{12 * 33}{444} + \frac{33 * 26}{393} + \frac{65 * 9}{442}} = 3.99$$

True

False

Do not know

Because the OR(MH) is a weighted average of stratum-specific estimated, it will lie between them.



**True or False? The crude odds ratio will always be a value between the stratum-specific odds ratio estimates.**

	1995		1996		1997		Combined	
Derek Jeter	12/48	.250	183/582	.314	190/654	.291	385/1284	.300
David Justice	104/411	.253	45/140	.321	163/495	.329	312/1046	.298

True

 False

Do not know


# Test for Homogeneity Odds Ratio

- The test for homogeneity of the odds ratio has the following form

$$H = \sum_{i=1}^I \frac{[\ln(\hat{OR}_i) - \ln \hat{OR}_{MH}]^2}{\hat{Var}_i [\ln(\hat{OR}_i)]} \sim \chi^2_{I-1}$$

- Where:
  - $\hat{OR}_i$  is the stratum specific odds ratio
  - $\hat{OR}_{MH}$  is the Mantel-Haenszel summary odds ratio
  - $\hat{Var}[\ln(\hat{OR}_i)]$  is the (large-sample) variance of the stratum specific  $\ln(OR)$ ;
- H is distributed  $\chi^2$  with I-1 degrees of freedom

**Based on the point estimates and 95% confidence intervals from a case-control study, do you think there is clear evidence of effect measure modification of the odds ratio for the association between OC use and myocardial infarction?**



Yes

No

Maybe ✓

The confidence intervals are wide and tend to overlap with each other considerably.

It is not clear whether the observed heterogeneity is more than one would expect due to random variability.

**If you reject the null hypothesis and conclude that there is statistically significant effect measure modification of the odds ratio, it is valid to report which of the following:**

$$H = \sum_{i=1}^I \frac{[\ln(\hat{OR}_i) - \ln \hat{OR}_{MH}]^2}{\hat{Var}_i[\ln(\hat{OR}_i)]}$$

The crude OR

The Mantel Haenszel OR

The stratum-specific ORs

The standardized OR

**HAVE A GOOD WEEK**