

# Neutron Stars in General Relativity

Diego García Tejada<sup>a,b</sup>

<sup>a</sup>Universidad Autónoma de Madrid, Cantoblanco, 28049, Madrid, Spain

<sup>b</sup>Instituto de Física Teórica UAM/CSIC, Cantoblanco, 28049, Madrid, Spain

December 2025

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

## 1 Introduction

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

## 2 Equations of stellar structure

In this section we will develop the equations of stellar structure for both Newtonian gravity and general relativity. Once derived we will compare their behavior by solving them for a toy model of a neutron star of constant density. By doing this we will see how, at such high densities, GR is necessary for the description of compact objects.

### 2.1 Newtonian equation

Let our star be spherically symmetric, and have mass  $M$  and radius  $R$ . The mass enclosed by a shell of radius  $r$  (for  $r \leq R$ ) is [1]:

$$m(r) = \int_0^r 4\pi r^2 \rho(r) dr \Rightarrow \frac{dm(r)}{dr} = 4\pi r^2 \rho(r), \quad (1)$$

where we have taken the limit of the energy density  $\rho$  being equal to the mass density  $\rho_0$  [2].

To calculate the equation of hydrostatic equilibrium we equate the forces generated by stars on gravity to the pressure generated by the fluid. The total pressure force generated on a shell of radius  $r$  and thickness  $dr$  is

$$F_P = -[p(r+dr) - p(r)] 4\pi r^2 = -4\pi r^2 \frac{dp}{dr} dr. \quad (2)$$

In order for the star to be in hydrostatic equilibrium, we need this force to be equal to that of gravity. By doing this we derive the equation of hydrostatic equilibrium<sup>1</sup> to be

$$4\pi r^2 \frac{dp}{dr} dr = -\frac{Gm(r)dm}{r^2} \Rightarrow \frac{dp}{dr} = -\frac{Gm(r)\rho(r)}{r^2}, \quad (3)$$

where in the second step we used (1) to rewrite  $dm$ .

### 2.2 TOV equation

In GR to find the equations of stellar structure we need to solve Einstein's equation for the interior of a solid object where  $T_{\mu\nu} \neq 0$ . We start again by considering a spherically symmetric static star. The general metric describing such a space is [3, 4]

$$ds^2 = -e^{2\Phi(r)} dt^2 + e^{2\Lambda(r)} dr^2 + r^2 d\Omega^2. \quad (4)$$

From this metric we compute the components of the Einstein tensor to be [3]

$$\begin{aligned} G_{tt} &= \frac{1}{r^2} e^{2(\Phi-\Lambda)} (2r\partial_r \Lambda - 1 + e^{2\Lambda}), \\ G_{rr} &= \frac{1}{r^2} (2r\partial_r \Phi + 1 - e^{2\Lambda}), \\ G_{\theta\theta} &= r^2 e^{-2\Lambda} [\partial_r^2 \Phi + (\partial_r \Phi)^2 - \partial_r \Phi \partial_r \Lambda \\ &\quad + \frac{1}{r} (\partial_r \Phi - \partial_r \Lambda)], \\ G_{\phi\phi} &= \sin^2 \theta G_{\theta\theta}. \end{aligned} \quad (5)$$

<sup>1</sup>Notice that (3) is  $\rho \frac{dv}{dr} = -\nabla p - \rho \nabla \Phi$ , when  $\mathbf{v} = 0$ .

Modeling the star as a perfect fluid, in its commoving frame, the energy-momentum tensor is

$$T_{\mu\nu} = (\rho - p)U_\mu U_\nu + pg_{\mu\nu}. \quad (6)$$

Normalizing the four-velocity to be timelike, we find that in this frame it becomes  $U_\mu = (e^\Phi, 0, 0, 0)$ . Inserting this into (6) we obtain that

$$\begin{aligned} T_{tt} &= \rho e^{2\Phi}, \\ T_{rr} &= p e^{2\Lambda}, \\ T_{\theta\theta} &= p r^2, \\ T_{\phi\phi} &= \sin^2 \theta T_{\theta\theta}. \end{aligned} \quad (7)$$

With (5) and (7) we find three independent components of Einstein's equation. the  $tt$  component

$$\frac{1}{r^2} e^{-2\Lambda} (2r \partial_r \Lambda - 1 + e^{2\Lambda}) = 8\pi G \rho, \quad (8)$$

the  $rr$  component

$$\frac{1}{r^2} e^{-2\Lambda} (2r \partial_r \Phi + 1 - e^{2\Lambda}) = 8\pi G p, \quad (9)$$

and the  $\theta\theta$  component

$$e^{-2\Lambda} [\partial_r^2 \Phi + (\partial_r \Phi)^2 - \partial_r \Phi \partial_r \Lambda + \frac{1}{r} (\partial_r \Phi - \partial_r \Lambda)] = 8\pi G p. \quad (10)$$

Motivated by Schwarzschild's exterior solution we make the following change of variable

$$m(r) = \frac{1}{2G} (r - r e^{-2\Lambda}) \Leftrightarrow e^{2\Lambda} = \left[ 1 - \frac{2Gm(r)}{r} \right]^{-1}. \quad (11)$$

Substituting this new definition onto equation (8) allows us to simplify it to

$$\frac{dm(r)}{dr} = 4\pi \rho(r) r^2. \quad (12)$$

We see that this equation is identical to (12), with the caveat that  $\rho \neq \rho_0$  outside the NR limit<sup>2</sup>. Using (11) in (9) we find that it can be rewritten as

$$\frac{d\Phi(r)}{dr} = \frac{4\pi G r^3 p(r) + Gm(r)}{r [r - 2Gm(r)]}. \quad (13)$$

From the conservation of Energy-Momentum ( $\nabla_\mu T^{\mu\nu} = 0$ ), where the only non null component will be that with  $\nu = r$  [3], we obtain the following relation

$$(\rho + p) \frac{d\Phi}{dr} = -\frac{dp}{dr}. \quad (14)$$

Inserting this result back onto (13) we finally obtain the equation of hydrostatic equilibrium for general relativity, also called the Tolman-Oppenheimer-Volkoff equation.

$$\frac{dp}{dr} = -(\rho + p) \frac{4\pi G r^3 p(r) + Gm(r)}{r [r - 2Gm(r)]}. \quad (15)$$

<sup>2</sup>See [3, 4] for the effects of this difference.

## 2.3 Constant density star

Now, in order to compare both formalisms let us assume that our star has mass  $M$ , radius  $R$  and a constant density  $\rho_\star$  for  $r \leq R$ . Under this assumptions both (1) and (12) result in

$$m(r) = \begin{cases} \frac{4}{3}\pi\rho_\star r^3 & \text{for } r \leq R \\ \frac{4}{3}\pi\rho_\star R^3 & \text{for } r > R \end{cases}. \quad (16)$$

With this, we now solve the differential equations (3) and (15). The integration is left as an exercise to the reader, its important to note that the boundary condition is  $p(R) = 0$ . This condition is needed in order to match the interior metric to Schwarzschild at  $r = R$ , but physically it also makes sense as the limit of the star will be where there is no pressure due to the gas. The solution for the Newtonian equation is

$$p(r) = \frac{2\pi}{3} G \rho_\star^2 (R^2 - r^2), \quad (17)$$

while the solution for the TOV equation (15) is

$$p(r) = \rho_\star \left( \frac{\sqrt{R^3 - 2GMR^2} - \sqrt{R^3 - 2GMr^2}}{\sqrt{R^3 - 2GMr^2} - 3\sqrt{R^3 - 2GMR^2}} \right). \quad (18)$$




Figure 1: Pressure profiles of various constant density stars obtained in both Newtonian gravity and General Relativity. The different line style indicates whether it was obtained from the Newtonian equation (dashed) or the TOV equation (solid). The mass of the star is given in units of solar masses.

While the equation governing the mass is the same for both models, therefore giving the same Mass-Radius values, the pressure profiles differ greatly. From figure 1 we can see that the general relativity solution gives higher pressure values for the same points  $r$ . This difference in pressure is greater as we increase  $r$  and we increase in  $M$ . This is due to the fact that equation 18 reaches a singularity for any  $M \geq 4R/9G$ . This is called Buchdahl's theorem [3] and is a general feature in GR.

When we solve the TOV equations for different stars we see that there is a maximum allowed mass before they collapse. This feature is not built into Newtonian gravity, we can set a limit of which the matter inside the star collapses but this would be due to matter degeneracy pressure where as this limit is entirely due to gravity.

This reason is why we need GR in order to describe Neutron Stars, and other compact objects, the pressures are

so high that the Newtonian approximation would not be valid. For the same masses and radii the pressure would be so low that we could not be able to explain how a neutron star can come to be in the first place [1–5].

### 3 Equations of State

In this section we will look at two possible Equations of State (EoSs) to describe Neutron Stars. The first model is the first ever considered to describe neutron stars [5], it consists on a completely degenerate ideal fermi gas (where the fermions are neutrons). The second model is a more realistic one obtained from chiral effective field theory [6], due to the complex nature of this, we will take it as god-given and only analyze the phenomenology that it brings.

#### 3.1 Free Fermi Gas

From statistical mechanics we know that the number density of a gas can be computed as

$$n = \int \frac{g f}{h^3} d^3 k, \quad (19)$$

where  $f$  is the distribution function,  $g$  indicates the degeneracy of spin  $g = 2S + 1$ ,  $k$  is the particles momenta and  $h$  is Planck's constant [1, 2, 7, 8].

The energy density will be [2, 8]

$$\rho = \int E \frac{g f}{h^3} d^3 k, \quad (20)$$

where  $E$  is the particle's energy ( $E^2 = k^2 + m^2$ ).

Finally the pressure is given by [1, 2, 7]

$$p = \frac{g}{3h^3} \int k v f d^3 k, \quad (21)$$

where  $v$  is the particles velocity ( $v = k/E$ ).

If we have a completely degenerate Fermi Gas<sup>3</sup> the distribution function becomes

$$f(E) = \begin{cases} 1, & E \leq E_F \\ 0, & E \geq E_F \end{cases}, \quad (22)$$

where  $E_F$  is the Fermi energy [8]. Substituting this into the previous expressions we get the following relations for: the number density

$$n = \int_0^{k_F} \frac{g}{h^3} 4\pi k^2 dk \Rightarrow n = \frac{8\pi k_F^3}{3h^3} = \frac{8\pi m_\chi^3 \xi_F^3}{3h^3}, \quad (23)$$

the energy density

$$\rho = \frac{8\pi}{h^3} \int_0^{k_F} \sqrt{m_\chi^2 + k^2} k^2 dk = \frac{\pi m_\chi^4}{h^3} \psi(\xi_F), \quad (24)$$

and the pressure

$$p = \frac{8\pi}{3h^3} \int_0^{k_F} \frac{k^4 dk}{\sqrt{k^2 + m_\chi^2}} = \frac{\pi m_\chi^4}{h^3} \phi(\xi_F), \quad (25)$$

where

$$\psi = \sqrt{1 + \xi_F^2} (2\xi_F^3 + \xi_F) - \ln \left( \xi_F + \sqrt{1 + \xi_F^2} \right), \quad (26)$$

$$\phi = \sqrt{1 + \xi_F^2} \left( \frac{2}{3} \xi_F^3 - \xi_F \right) + \ln \left( \xi_F + \sqrt{1 + \xi_F^2} \right). \quad (27)$$

<sup>3</sup>We can consider a NS to be degenerate as its temperature is smaller than the Fermi Temperature [4].

### 3.2 Nuclear Matter

Fusce mauris. Vestibulum luctus nibh at lectus. Sed bibendum, nulla a faucibus semper, leo velit ultricies tellus, ac venenatis arcu wisi vel nisl. Vestibulum diam. Aliquam pellentesque, augue quis sagittis posuere, turpis lacus congue quam, in hendrerit risus eros eget felis. Maecenas eget erat in sapien mattis porttitor. Vestibulum porttitor. Nulla facilisi. Sed a turpis eu lacus commodo facilisis. Morbi fringilla, wisi in dignissim interdum, justo lectus sagittis dui, et vehicula libero dui cursus dui. Mauris tempor ligula sed lacus. Duis cursus enim ut augue. Cras ac magna. Cras nulla. Nulla egestas. Curabitur a leo. Quisque egestas wisi eget nunc. Nam feugiat lacus vel est. Curabitur consectetur.

### 4 Solutions for the TOV equation

Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris.

#### 4.1 Numerical integration of the TOV equation

Suspendisse vel felis. Ut lorem lorem, interdum eu, tincidunt sit amet, laoreet vitae, arcu. Aenean faucibus pede eu ante. Praesent enim elit, rutrum at, molestie non, nonummy vel, nisl. Ut lectus eros, malesuada sit amet, fermentum eu, sodales cursus, magna. Donec eu purus. Quisque vehicula, urna sed ultricies auctor, pede lorem egestas dui, et convallis elit erat sed nulla. Donec luctus. Curabitur et nunc. Aliquam dolor odio, commodo pretium, ultricies non, pharetra in, velit. Integer arcu est, nonummy in, fermentum faucibus, egestas vel, odio.

#### 4.2 Fermionic Star

Sed commodo posuere pede. Mauris ut est. Ut quis purus. Sed ac odio. Sed vehicula hendrerit sem. Duis non odio. Morbi ut dui. Sed accumsan risus eget odio. In hac habitasse platea dictumst. Pellentesque non elit. Fusce sed justo eu urna porta tincidunt. Mauris felis odio, sollicitudin sed, volutpat a, ornare ac, erat. Morbi quis dolor. Donec pellentesque, erat ac sagittis semper, nunc dui lobortis purus, quis congue purus metus ultricies tellus. Proin et quam. Class aptent taciti sociosqu ad litora torquent per conubia nostra, per inceptos hymenaeos. Praesent sapien turpis, fermentum vel, eleifend faucibus, vehicula eu, lacus.

#### 4.3 Neutron Star

Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Donec odio elit, dictum in, hendrerit sit amet, egestas sed, leo. Praesent fe-

giat sapien aliquet odio. Integer vitae justo. Aliquam vestibulum fringilla lorem. Sed neque lectus, consectetur at, consectetur sed, eleifend ac, lectus. Nulla facilisi. Pellentesque eget lectus. Proin eu metus. Sed porttitor. In hac habitasse platea dictumst. Suspendisse eu lectus. Ut mi mi, lacinia sit amet, placerat et, mollis vitae, dui. Sed ante tellus, tristique ut, iaculis eu, malesuada ac, dui. Mauris nibh leo, facilisis non, adipiscing quis, ultrices a, dui.

## 5 Conclusions

Morbi luctus, wisi viverra faucibus pretium, nibh est placerat odio, nec commodo wisi enim eget quam. Quisque libero justo, consectetur a, feugiat vitae, porttitor eu, libero. Suspendisse sed mauris vitae elit sollicitudin malesuada. Maecenas ultricies eros sit amet ante. Ut venenatis velit. Maecenas sed mi eget dui varius euismod. Phasellus aliquet volutpat odio. Vestibulum ante ipsum primis in faucibus orci luctus et ultrices posuere cubilia Curae; Pellentesque sit amet pede ac sem eleifend consectetur. Nullam elementum, urna vel imperdiet sodales, elit ipsum pharetra ligula, ac pretium ante justo a nulla. Curabitur tristique arcu eu metus. Vestibulum lectus. Proin mauris. Proin eu nunc eu urna hendrerit faucibus. Aliquam auctor, pede consequat laoreet varius, eros tellus scelerisque quam, pellentesque hendrerit ipsum dolor sed augue. Nulla nec lacus.

Suspendisse vitae elit. Aliquam arcu neque, ornare in, ullamcorper quis, commodo eu, libero. Fusce sagittis erat at erat tristique mollis. Maecenas sapien libero, molestie et, lobortis in, sodales eget, dui. Morbi ultrices rutrum lorem. Nam elementum ullamcorper leo. Morbi dui. Aliquam sagittis. Nunc placerat. Pellentesque tristique sodales est. Maecenas imperdiet lacinia velit. Cras non urna. Morbi eros pede, suscipit ac, varius vel, egestas non, eros. Praesent malesuada, diam id pretium elementum, eros sem dictum tortor, vel consectetur odio sem sed wisi.

Sed feugiat. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Ut pellentesque augue sed urna. Vestibulum diam eros, fringilla et, consectetur eu, nonummy id, sapien. Nullam at lectus. In sagittis ultrices mauris. Curabitur malesuada erat sit amet massa. Fusce blandit. Aliquam erat volutpat. Aliquam euismod. Aenean vel lectus. Nunc imperdiet justo nec dolor.

Etiam euismod. Fusce facilisis lacinia dui. Suspendisse potenti. In mi erat, cursus id, nonummy sed, ullamcorper eget, sapien. Praesent pretium, magna in eleifend egestas, pede pede pretium lorem, quis consectetur tortor sapien facilisis magna. Mauris quis magna varius nulla scelerisque imperdiet. Aliquam non quam. Aliquam porttitor quam a lacus. Praesent vel arcu ut tortor cursus volutpat. In vitae pede quis diam bibendum placerat. Fusce elementum convallis neque. Sed dolor orci, scelerisque ac, dapibus nec, ultricies ut, mi. Duis nec dui quis leo sagittis commodo.

## References

[1] S. Chandrasekhar, *An Introduction to the Study of Stellar Structure*. Chicago: University of Chicago

Press, 1939.

- [2] S. L. Shapiro and S. A. Teukolsky, *Black Holes, White Dwarfs, and Neutron Stars: The Physics of Compact Objects*. New York: Wiley-Interscience, 1983.
- [3] S. M. Carroll, *Spacetime and Geometry: An Introduction to General Relativity*. Cambridge, UK: Cambridge University Press, 2 ed., 2019.
- [4] C. W. Misner, K. S. Thorne, and J. A. Wheeler, *Gravitation*. San Francisco, USA: W. H. Freeman, 1973.
- [5] J. R. Oppenheimer and G. M. Volkoff, “On massive neutron cores,” *Phys. Rev.*, vol. 55, pp. 374–381, Feb 1939.
- [6] K. Hebeler, J. M. Lattimer, C. J. Pethick, and A. Schwenk, “Equation of state and neutron star properties constrained by nuclear physics and observation,” *The Astrophysical Journal*, vol. 773, p. 11, jul 2013.
- [7] D. Clayton, *Principles of Stellar Evolution and Nucleosynthesis*. Astronomy / Astrophysics, University of Chicago Press, 1983.
- [8] R. K. Pathria and P. D. Beale, *Statistical Mechanics*. Oxford: Elsevier, 3rd ed., 2011.