

TABLE 1. Definition of the survey indicators available in the SEEDNet Library.

Indicator ^a	Numerator	Denominator
Access to Electricity	Households with electricity	All households
Access to Improved Water Source	Households whose main source of drinking water is an improved source	All households
Iodized Salt Intake in Household	Number of households with salt tested with iodized salt	Number of households where salt was tested for iodine content
Percentage of women with anemia	Non-pregnant women whose hemoglobin count is less than 12.0 grams per deciliter (g/dl) plus the number of pregnant women whose count is less than 11.0 g/dl	Women of reproductive age measured for anemia in households selected for anemia testing
Antenatal Care Visits (4+) during Pregnancy	Women who attended four or more antenatal care visits for their last most recent live birth or stillbirth	Women who have had a live birth in the two years preceding the survey
Low Birth Weight Prevalence	Live births in the two years preceding the survey who were weighed at birth and were reported as weighing less than 2.5 kg	Live births in the last two years who were weighed at birth
Exclusive Breastfeeding (0-5 months)	Children aged 0-5 months living with their mother who were breastfeeding and given nothing else the day preceding the interview	Children born 0-5 months preceding the survey who are living with their mother
BCG Immunization (12-23 months)	Living children age 12–23 months receiving specified vaccine according to vaccination card or mother’s report	Living children age 12–23 months
DPT3 Immunization (12-23 months)	Living children age 12–23 months receiving specified vaccine according to vaccination card or mother’s report	Living children age 12–23 months
Measles Immunization (12-23 months)	Living children age 12–23 months receiving specified vaccine according to vaccination card or mother’s report	Living children age 12–23 months
Children Slept u/ Bed Net (0-59 months)	Children age 0-59 months who slept under an insecticide-treated net (ITN)	Children age 0-59 months
Diarrhea Treatment with ORS (0-59 months)	Living children ;59 months with diarrhea during the two weeks preceding interview who were given fluid from oral rehydration salts packet or pre-packaged ORS fluid	Children 0-59 months with diarrhea in the two weeks preceding the interview
Stunting Prevalence (0-59 months)	Children ;59 months whose height-for-age z-score is below minus 2 (-2.0) standard deviations (SD) below the median on the WHO Child Growth Standards	Living children 0-59 months before the survey who have a valid non-flagged height for age z-scores
Wasting Prevalence (0-59 months)	Children ;59 months whose weight-for-height z-score is below minus 2 (-2.0) standard deviations (SD) below the median on the WHO Child Growth Standards	Living children 0-59 months before the survey who have a valid non-flagged weight for height z-scores
Prevalence of Anemia in Children (6-59 months)	Children whose hemoglobin count is less than 11 grams per deciliter (g/dl)	Children 6-59 months who were measured in households selected for anemia testing

Note: ^a Standard definitions are used in all Settlement-level datasets. For the most recent detailed definitions, we refer to the Guide to DHS Statistics DHS-8. Indicators that were not measured in the survey or that did not match the values published in the original DHS report or in StatCompiler are not reported. Anemia; Women, Anemia; Children, Wasting, Stunting, Diarrhoea Treatment; ORS, Water Source, Low Birth Weight, Exclusive Breastfeeding, BCG, DPT3 and Measles Immunization, Iodized Salt, Electricity, Antenatal Care.