

3D Magnetic modelling of ellipsoidal bodies

Diego Takahashi Tomazella¹ and Vanderlei C. Oliveira Jr¹

¹Department of Geophysics, Observatório Nacional, Rio de Janeiro, Brazil

Correspondence to: Diego Takahashi Tomazella (diego.takahashi@gmail.com)

Abstract.

TEXT

1 Introduction

TEXT

5 2 Methodology

2.1 Geometrical parameters and coordinate systems

Let (x, y, z) be a point referred to a Cartesian coordinate system with axes x , y and z pointing to, respectively, North, East and down. For convenience, we denominate this coordinate system as *main coordinate system*. Let us consider an ellipsoidal body with centre at the point (x_c, y_c, z_c) , semi-axes defined by positive constants a , b , c , where $a > b > c$, and orientation defined by
10 three angles α , β , and γ . The points (x, y, z) located on the surface of this ellipsoidal body satisfy the following equation:

$$(\mathbf{r} - \mathbf{r}_c)^T \mathbf{A} (\mathbf{r} - \mathbf{r}_c) = 1, \quad (1)$$

where $\mathbf{r} = [x \ y \ z]^T$, $\mathbf{r}_c = [x_c \ y_c \ z_c]^T$, \mathbf{A} is a positive definite matrix given by

$$\mathbf{A} = \mathbf{V} \begin{bmatrix} a^{-2} & 0 & 0 \\ 0 & b^{-2} & 0 \\ 0 & 0 & c^{-2} \end{bmatrix} \mathbf{V}^T, \quad (2)$$

and \mathbf{V} is an orthogonal matrix whose columns are defined by unit vectors \mathbf{v}_1 , \mathbf{v}_2 , and \mathbf{v}_3 .

15 The vectors \mathbf{v}_1 , \mathbf{v}_2 , and \mathbf{v}_3 depend on the orientation angles α , β , γ and are defined as follows (Clark et al., 1986):

$$\mathbf{v}_1 = \begin{bmatrix} -\cos \alpha \cos \delta \\ -\sin \alpha \cos \delta \\ -\sin \delta \end{bmatrix}, \quad (3)$$

$$\mathbf{v}_2 = \begin{bmatrix} \cos \alpha \cos \gamma \sin \delta + \sin \alpha \sin \gamma \\ \sin \alpha \cos \gamma \sin \delta - \cos \alpha \sin \gamma \\ -\cos \gamma \cos \delta \end{bmatrix}, \quad (4)$$

$$\mathbf{v}_3 = \begin{bmatrix} \sin \alpha \cos \gamma - \cos \alpha \sin \gamma \sin \delta \\ -\cos \alpha \cos \gamma - \sin \alpha \sin \gamma \sin \delta \\ \sin \gamma \cos \delta \end{bmatrix}. \quad (5)$$

5 For triaxial ellipsoids (i.e., $a > b > c$), the orthogonal matrix \mathbf{V} (Eq. 2) is calculated by using Eqs. 3, 4, and 5 as follows:

$$\mathbf{V} = \begin{bmatrix} \mathbf{v}_1 & \mathbf{v}_2 & \mathbf{v}_3 \end{bmatrix}. \quad (6)$$

Similarly, the matrix \mathbf{V} (Eq. 2) for prolate ellipsoids (i.e., $a > b = c$) is calculated according to Equation 6 by using Equations 3, 4, and 5, but with $\gamma = 0^\circ$ (Emerson et al., 1985). Finally, the matrix \mathbf{V} (Eq. 2) for oblate ellipsoids (i.e., $a < b = c$) is calculated by using Eqs. 3, 4, and 5, with $\gamma = 0^\circ$, as follows (Emerson et al., 1985):

$$10 \quad \mathbf{V} = \begin{bmatrix} \mathbf{v}_2 & \mathbf{v}_1 & -\mathbf{v}_3 \end{bmatrix}. \quad (7)$$

The orientation of the semi-axes a , b , and c are defined by the first, second, and third columns of the matrix \mathbf{V} given by Equation 6, in the case of a triaxial or prolate ellipsoid, or the matrix \mathbf{V} given by Equation 7, in the case of an oblate ellipsoid.

The magnetic modelling of an ellipsoidal body is commonly performed in a particular Cartesian coordinate system that is aligned with the body semi-axes and has the origin coincident with the body centre. For convenience, we denominate this
 15 particular coordinate system as *local coordinate system*. The relationship between the Cartesian coordinates $(\tilde{x}, \tilde{y}, \tilde{z})$ of a point in a local coordinate system and the Cartesian coordinates (x, y, z) of the same point in the main system is given by:

$$\tilde{\mathbf{r}} = \mathbf{V}^\top (\mathbf{r} - \mathbf{r}_c), \quad (8)$$

where $\tilde{\mathbf{r}} = [\tilde{x} \ \tilde{y} \ \tilde{z}]^\top$, \mathbf{r} and \mathbf{r}_c are defined in Equation 1 and the matrix \mathbf{V} is defined according to Eqs. 6 or 7, depending on the ellipsoid type.

20 2.2 Theoretical background

Based on the mathematical theory of the magnetic induction developed by Poisson (1824), Maxwell (1873) affirmed that, if V is the gravitational potential produced by any body with uniform density ρ and arbitrary shape at a point (x, y, z) , then $-\frac{\partial V}{\partial x}$ is the magnetic scalar potential produced at the same point by the same body if it has a uniform magnetization oriented along x with intensity ρ . Maxwell (1873) generalized this idea as a way of determining the magnetic scalar potential produced
 25 by any body uniformly magnetized in a given direction. By presuming that this uniform magnetization is due to induction, he postulated that the resulting magnetic field (intensity) at all points within the body must also be uniform and parallel the

magnetization, which results that the gravitational potential V at points within the body must be a quadratic function of the spatial coordinates. Apparently, Maxwell (1873) was the first one to affirm that the only finite bodies having a gravitational potential with this property and that, as a consequence, can be uniformly magnetized in the presence of a uniform and static magnetic field are the ones bounded by surfaces of second degree, which are ellipsoids.

- 5 Consider a magnetized ellipsoid immersed in a uniform magnetic field \mathbf{H}_0 (in Am^{-1}). This uniform field can be, for example, the main component of the Earth's magnetic field, which is usually assumed to be generated by the Earth's liquid core. In the absence of conduction currents, the total magnetic field $\mathbf{H}(\mathbf{r})$ at the position \mathbf{r} (Eqs. 2 and 8) of a point referred to the main coordinate system is defined as follows (Stratton, 2007):

$$\mathbf{H}(\mathbf{r}) = \mathbf{H}_0 - \nabla\phi(\mathbf{r}), \quad (9)$$

- 10 where the second term is the negative gradient of the magnetic scalar potential $\phi(\mathbf{r})$ given by:

$$\phi(\mathbf{r}) = -\frac{1}{4\pi} \iiint_V \mathbf{M}(\mathbf{r}')^\top \nabla \left(\frac{1}{\|\mathbf{r} - \mathbf{r}'\|} \right) dx' dy' dz'. \quad (10)$$

In this equation, $\mathbf{r}' = [x' \ y' \ z']^\top$ is the position vector of a point located within the volume V , the integral is conducted over the variables x' , y' and, z' representing the coordinates of a point located within the volume V of the ellipsoid, $\|\cdot\|$ denotes the Euclidean norm and $\mathbf{M}(\mathbf{r}')$ is the magnetization vector (in Am^{-1}). Equation 10 is valid anywhere, independently if the

- 15 position vector \mathbf{r} represents a point located inside or outside the magnetized body (DuBois, 1896).

Based on Maxwell's postulate, let us assume that the body has a uniform magnetization given by

$$\mathbf{M} = \mathbf{K} \mathbf{H}^*, \quad (11)$$

where \mathbf{H}^* is the resultant uniform magnetic field at any point within the body and \mathbf{K} is a constant and symmetrical 2nd-order tensor representing the magnetic susceptibility of the body. In this case, Equation 9 can be rewritten as follows:

$$20 \quad \mathbf{H}(\mathbf{r}) = \mathbf{H}_0 - \mathbf{N}(\mathbf{r}) \mathbf{K} \mathbf{H}^*, \quad (12)$$

where $\mathbf{N}(\mathbf{r})$ is a symmetrical matrix whose ij -element $n_{ij}(\mathbf{r})$ is given by

$$n_{ij}(\mathbf{r}) = -\frac{1}{4\pi} \frac{\partial^2 f(\mathbf{r})}{\partial r_i \partial r_j}, \quad i = 1, 2, 3, \quad j = 1, 2, 3, \quad (13)$$

$r_1 = x$, $r_2 = y$, $r_3 = z$ are the elements of the position vector \mathbf{r} (Eq. 1), and

$$f(\mathbf{r}) = \iiint_V \frac{1}{\|\mathbf{r} - \mathbf{r}'\|} dx' dy' dz'. \quad (14)$$

- 25 Notice that the scalar function $f(\mathbf{r})$ (Eq. 14) is proportional to the gravitational potential that would be produced by the ellipsoidal body with volume V if it had a uniform density equal to the inverse of the gravitational constant. It can be shown that the elements $n_{ij}(\mathbf{r})$ are finite whether \mathbf{r} is a point within or without the volume V (Peirce, 1902; Webster, 1904). The matrix $\mathbf{N}(\mathbf{r})$ (Eq. 12) is called *depolarization tensor* (Soliv  rez, 1981, 2008).

The following part of this paper moves on to describe the magnetic field $\mathbf{H}(\mathbf{r})$ (Eq. 12) at points located both within and without the volume V of the ellipsoidal body. However, the mathematical developments are conveniently performed in the local coordinate system related to the respective ellipsoidal body.

2.3 Coordinate transformation

- 5 To continue our description of the magnetic modelling of ellipsoidal bodies, it is convenient to perform two important coordinate transformations. The first one transforms the scalar function $f(\mathbf{r})$ (Eq. 14) from the main coordinate system into a new scalar function $\tilde{f}(\tilde{\mathbf{r}})$ referred to the local coordinate system. The function $\tilde{f}(\tilde{\mathbf{r}})$ was first presented by Dirichlet (1839) to describe the gravitational potential produced by homogeneous ellipsoids. Posteriorly, several authors also deduced and used this function for describing the magnetic and gravitational fields produced by triaxial, prolate, and oblate ellipsoids (Maxwell, 1873; Thomson and Tait, 1879; DuBois, 1896; Peirce, 1902; Webster, 1904; Kellogg, 1929; Stoner, 1945; Osborn, 1945; 10 Lowes, 1974; Peake and Davy, 1953; Chang, 1961; Clark et al., 1986; Tejedor et al., 1995; Stratton, 2007). It is convenient to use $\tilde{f}^*(\tilde{\mathbf{r}})$ and $\tilde{f}^\dagger(\tilde{\mathbf{r}})$ to define the function $\tilde{f}(\tilde{\mathbf{r}})$ evaluated, respectively, at points $\tilde{\mathbf{r}}$ inside and outside the volume V of the ellipsoidal body.

The scalar function $\tilde{f}^*(\tilde{\mathbf{r}})$ is given by

$$15 \quad \tilde{f}^*(\tilde{\mathbf{r}}) = \pi abc \int_0^\infty \left(1 - \frac{\tilde{x}^2}{a^2 + u} - \frac{\tilde{y}^2}{b^2 + u} - \frac{\tilde{z}^2}{c^2 + u} \right) \frac{1}{R(u)} du, \quad \tilde{\mathbf{r}} \in V, \quad (15)$$

where

$$R(u) = \sqrt{(a^2 + u)(b^2 + u)(c^2 + u)}. \quad (16)$$

- This function represents the gravitational potential that would be produced by the ellipsoidal body at points located within its volume V if it had a uniform density equal to the inverse of the gravitational constant. Notice that, in this case, the gravitational 20 potential is a quadratic function of the spatial coordinates \tilde{x} , \tilde{y} , and \tilde{z} , which supported the Maxwell's (1873) postulate about uniformly magnetized ellipsoids. In a similar way, the function $\tilde{f}^\dagger(\tilde{\mathbf{r}})$ is given by

$$\tilde{f}^\dagger(\tilde{\mathbf{r}}) = \pi abc \int_\lambda^\infty \left(1 - \frac{\tilde{x}^2}{a^2 + u} - \frac{\tilde{y}^2}{b^2 + u} - \frac{\tilde{z}^2}{c^2 + u} \right) \frac{1}{R(u)} du, \quad \tilde{\mathbf{r}} \notin V, \quad (17)$$

- where $R(u)$ is defined by Equation 16 and the parameter λ is defined according to the ellipsoid type as a function of the spatial coordinates \tilde{x} , \tilde{y} , and \tilde{z} (see Appendix B). For readers interested in additional information about the parameter λ , we 25 recommend Webster (1904, p. 234), Kellogg (1929, p. 184) and Clark et al. (1986).

The second important coordinate transformation is defined with respect to Equation 12. By properly using the orthogonality of matrix \mathbf{V} (Eqs. 6 and 7), the magnetic field $\mathbf{H}(\mathbf{r})$ (Eq. 12) can be transformed from the main coordinate system to a local coordinate system as follows:

$$\underbrace{\mathbf{V}^\top \mathbf{H}(\mathbf{r})}_{\tilde{\mathbf{H}}(\tilde{\mathbf{r}})} = \underbrace{\mathbf{V}^\top \mathbf{H}_0}_{\tilde{\mathbf{H}}_0} - \underbrace{\mathbf{V}^\top \mathbf{N}(\mathbf{r}) \mathbf{V}}_{\tilde{\mathbf{N}}(\tilde{\mathbf{r}})} \underbrace{\mathbf{V}^\top \mathbf{K} \mathbf{V}}_{\tilde{\mathbf{K}}} \underbrace{\mathbf{V}^\top \mathbf{H}^*}_{\tilde{\mathbf{H}}^*}, \quad (18)$$

where the superscript " \sim " denotes quantities referred to the respective local coordinate system.

In Equation 18, the transformed depolarization tensor $\tilde{\mathbf{N}}(\tilde{\mathbf{r}})$ is calculated as a function of the original depolarization tensor $\mathbf{N}(\mathbf{r})$ (Eq. 12). In this case, the elements of $\tilde{\mathbf{N}}(\tilde{\mathbf{r}})$ are calculated as a function of the second derivatives of the function $f(\mathbf{r})$ (Eq. 14), which is defined in the main coordinate system. It can be shown (Appendix A), however, that the elements $\tilde{n}_{ij}(\tilde{\mathbf{r}})$ of

5 $\tilde{\mathbf{N}}(\tilde{\mathbf{r}})$ can also be calculated as follows:

$$\tilde{n}_{ij}(\tilde{\mathbf{r}}) = -\frac{1}{4\pi} \frac{\partial^2 \tilde{f}(\tilde{\mathbf{r}})}{\partial \tilde{r}_i \partial \tilde{r}_j}, \quad i = 1, 2, 3, \quad j = 1, 2, 3, \quad (19)$$

where $\tilde{r}_1 = \tilde{x}$, $\tilde{r}_2 = \tilde{y}$, and $\tilde{r}_3 = \tilde{z}$ are the elements of the transformed vector $\tilde{\mathbf{r}}$ (Eq. 8) and $\tilde{f}(\tilde{\mathbf{r}})$ is given by Equation 15 or 17, depending if $\tilde{\mathbf{r}}$ represents a point located within or without the volume V of the ellipsoidal body.

2.4 Internal magnetic field and magnetization

10 By considering $\tilde{\mathbf{r}}$ as a point within the volume V of the ellipsoid and using the Maxwell's postulate about the uniformity of the magnetic field $\mathbf{H}(\mathbf{r})$ inside ellipsoidal bodies, we can use Equation 18 for defining the resultant uniform magnetic field $\tilde{\mathbf{H}}^*$ inside the ellipsoidal body as follows:

$$\tilde{\mathbf{H}}^* = \left(\mathbf{I} - \tilde{\mathbf{N}}^* \tilde{\mathbf{K}} \right)^{-1} \tilde{\mathbf{H}}_0, \quad (20)$$

where \mathbf{I} is the identity matrix and $\tilde{\mathbf{N}}^*$ represents the transformed depolarization tensor evaluated within the volume V of the
15 ellipsoidal body. The elements of $\tilde{\mathbf{N}}^*$ will be presented in a latter section.

Let us pre-multiply the uniform internal field $\tilde{\mathbf{H}}^*$ (Eq. 20) by the transformed susceptibility tensor $\tilde{\mathbf{K}}$ (Eq. 18) to obtain

$$\begin{aligned} \tilde{\mathbf{M}} &= \tilde{\mathbf{K}} \left(\mathbf{I} - \tilde{\mathbf{N}}^* \tilde{\mathbf{K}} \right)^{-1} \tilde{\mathbf{H}}_0 \\ &= \left(\mathbf{I} - \tilde{\mathbf{K}} \tilde{\mathbf{N}}^* \right)^{-1} \tilde{\mathbf{K}} \tilde{\mathbf{H}}_0, \end{aligned} \quad (21)$$

where $\tilde{\mathbf{M}}$ represents the transformed magnetization, as can be easily verified by using Eqs. 11 and 18. The matrix identity used for obtaining the second line of Equation 21 is given by Searle (1982, p. 151).

20 Equation 21 can be easily generalized for the case in which the ellipsoid has also a uniform remanent magnetization $\tilde{\mathbf{M}}_R$. Let us first consider that the uniform remanent magnetization satisfies the condition $\tilde{\mathbf{H}}_A = \tilde{\mathbf{K}}^{-1} \tilde{\mathbf{M}}_R$, where $\tilde{\mathbf{H}}_A$ represents a hypothetical uniform ancient field. Then, if we assume that $\tilde{\mathbf{H}}_0$, in Eqs. 20 and 21, is in fact the sum of the uniform magnetic field $\tilde{\mathbf{H}}_0$ and the hypothetical ancient field $\tilde{\mathbf{H}}_A$, we obtain the following generalized equation

$$\tilde{\mathbf{M}} = \left(\mathbf{I} - \tilde{\mathbf{K}} \tilde{\mathbf{N}}^* \right)^{-1} \left(\tilde{\mathbf{K}} \tilde{\mathbf{H}}_0 + \tilde{\mathbf{M}}_R \right). \quad (22)$$

25 Equation 22 is consistent with that given by Clark et al. (1986, Eq. 38).

2.5 External magnetic field and total-field anomaly

The magnetic field $\Delta\tilde{\mathbf{H}}(\tilde{\mathbf{r}})$ produced by an ellipsoid at external points is calculated from Eqs. 18 and 22 as the difference between the resultant field $\tilde{\mathbf{H}}(\tilde{\mathbf{r}})$ and the uniform field $\tilde{\mathbf{H}}_0$:

$$\Delta\tilde{\mathbf{H}}(\tilde{\mathbf{r}}) = -\tilde{\mathbf{N}}^\dagger(\tilde{\mathbf{r}})\tilde{\mathbf{M}}. \quad (23)$$

- 5 The elements of the transformed depolarization tensor $\tilde{\mathbf{N}}^\dagger(\tilde{\mathbf{r}})$ used in Equation 23 will be presented in the following section.

The $\Delta\tilde{\mathbf{H}}(\tilde{\mathbf{r}})$ can be interpreted as a uniformly magnetized body located in the crust while the uniform field $\tilde{\mathbf{H}}_0$ can be interpreted as the main component of the geomagnetic field on the study area, which is commonly assumed to be generated at the liquid part of the Earth's core. Equation 23 gives the magnetic field (in A m^{-1}) produced by an ellipsoid. However, in geophysics, the most widely used field is the magnetic induction (in nT). Fortunately, this conversion can be easily done by

- 15 multiplying Equation 23 by $k_m = 10^9 \mu_0$, where μ_0 represents the magnetic constant (in H m^{-1}).

For geophysical applications, it is preferable to calculate the total-field anomaly produced by the magnetic sources. This scalar quantity is defined as follows (Blakely, 1996):

$$\Delta\tilde{T}(\tilde{\mathbf{r}}) = \|\tilde{\mathbf{B}}_0 + \Delta\tilde{\mathbf{B}}(\tilde{\mathbf{r}})\| - \|\tilde{\mathbf{B}}_0\|, \quad (24)$$

where $\tilde{\mathbf{B}}_0 = k_m \tilde{\mathbf{H}}_0$ and $\Delta\tilde{\mathbf{B}}(\tilde{\mathbf{r}}) = k_m \Delta\tilde{\mathbf{H}}(\tilde{\mathbf{r}})$ (Eq. 23). In practical situations, however, $\|\tilde{\mathbf{B}}_0\| \gg \|\Delta\tilde{\mathbf{B}}(\tilde{\mathbf{r}})\|$ and, conse-

- 15 quently, the following approximation is valid (Blakely, 1996):

$$\Delta\tilde{T}(\tilde{\mathbf{r}}) \approx \frac{\tilde{\mathbf{B}}_0^\top \Delta\tilde{\mathbf{B}}(\tilde{\mathbf{r}})}{\|\tilde{\mathbf{B}}_0\|}. \quad (25)$$

2.6 Transformed depolarization tensors $\tilde{\mathbf{N}}(\tilde{\mathbf{r}})$

2.6.1 Depolarization tensor $\tilde{\mathbf{N}}^*$

- 20 The elements of the transformed depolarization tensor $\tilde{\mathbf{N}}^*$ used to compute the uniform magnetic field $\tilde{\mathbf{H}}^*$ (Eq. 20) and the magnetization $\tilde{\mathbf{M}}$ (Eqs. 21 and 22) are calculated according to Equation 19, with $\tilde{f}(\tilde{\mathbf{r}})$ given by $\tilde{f}^*(\tilde{\mathbf{r}})$ (Eq. 15). As we have already pointed out, the $\tilde{f}^*(\tilde{\mathbf{r}})$ (Eq. 15) is a quadratic function of the spatial coordinates \tilde{x} , \tilde{y} and \tilde{z} . Consequently, the off-diagonal elements $\tilde{n}_{ij}^*(\tilde{\mathbf{r}})$, $i = 1, 2, 3$, $j = 1, 2, 3$, $i \neq j$, are equal to zero. The remaining diagonal elements are given by (Stoner, 1945):

$$\tilde{n}_{ii}^*(\tilde{\mathbf{r}}) = \frac{abc}{2} \int_0^\infty \frac{1}{(e_i^2 + u)R(u)} du, \quad i = 1, 2, 3, \quad (26)$$

- 25 where $R(u)$ is defined by Equation 16 and $e_1 = a$, $e_2 = b$, and $e_3 = c$. These elements are commonly known as *demagnetizing factors*. Notice that, according to Equations 18 and A7,

$$\mathbf{N}(\mathbf{r}) = \mathbf{V}\tilde{\mathbf{N}}^*\mathbf{V}^\top, \quad (27)$$

where $\tilde{\mathbf{N}}^*$ (Eqs. 20, 21 and 22) is a diagonal matrix and \mathbf{V} (Eqs. 6 and 7) is an orthogonal matrix. This equation shows that, for the particular case in which \mathbf{r} and consequently $\tilde{\mathbf{r}}$ represent a point inside the volume V of the ellipsoid, the elements $\tilde{n}_{ii}^*(\tilde{\mathbf{r}})$ (Eq. 26) of $\tilde{\mathbf{N}}^*$ represent the eigenvalues while the columns of \mathbf{V} represent the eigenvectors of the original depolarization tensor $\mathbf{N}(\mathbf{r})$.

5 INTRODUIZIR OS ELEMENTOS $\tilde{n}_{ii}^*(\tilde{\mathbf{r}})$ PARA OS TRES TIPOS DE ELIPSOIDE

2.6.2 Depolarization tensor $\tilde{\mathbf{N}}^\dagger(\tilde{\mathbf{r}})$

The elements of the transformed depolarization tensor $\tilde{\mathbf{N}}^\dagger(\tilde{\mathbf{r}})$ used to compute the magnetic field $\Delta\tilde{\mathbf{H}}(\tilde{\mathbf{r}})$ produced by the ellipsoid (Eq. 23) and the total-field anomaly $\Delta\tilde{T}(\tilde{\mathbf{r}})$ (Eqs. 24 and 25) are calculated according to Equation 19, with $\tilde{f}(\tilde{\mathbf{r}})$ given by $\tilde{f}^\dagger(\tilde{\mathbf{r}})$ (Eq. 17).

10 3 Conclusions

TEXT

Appendix A: Relationship between the derivatives of the functions $f(\mathbf{r})$ and $\tilde{f}(\tilde{\mathbf{r}})$

Let $\tilde{f}(\tilde{\mathbf{r}})$ be the scalar function obtained by transforming $f(\mathbf{r})$ (Eq. 14) from the main coordinate system to a local coordinate system. For convenience, let us rewrite Equation 8 as follows:

$$15 \quad \tilde{r}_k = v_{k1} r_1 + v_{k2} r_2 + v_{k3} r_3 + c_k, \quad (\text{A1})$$

where \tilde{r}_k , $k = 1, 2, 3$, are the elements of the transformed position vector $\tilde{\mathbf{r}}$ (Eq. 8), r_j , $j = 1, 2, 3$, are the elements of the position vector \mathbf{r} (Eq. 1), v_{kj} , $j = 1, 2, 3$, are the elements of the matrix \mathbf{V} (Eq. 6 or 7), and c_k is a constant defined by the coordinates x_c , y_c , and z_c of the centre of the ellipsoidal body.

By considering the functions $f(\mathbf{r})$ (Eq. 14) and $\tilde{f}(\tilde{\mathbf{r}})$ evaluated at the same point, but on different coordinate systems, we
20 have:

$$\frac{\partial f(\mathbf{r})}{\partial r_j} = \frac{\partial \tilde{f}(\tilde{\mathbf{r}})}{\partial \tilde{r}_1} \frac{\partial \tilde{r}_1}{\partial r_j} + \frac{\partial \tilde{f}(\tilde{\mathbf{r}})}{\partial \tilde{r}_2} \frac{\partial \tilde{r}_2}{\partial r_j} + \frac{\partial \tilde{f}(\tilde{\mathbf{r}})}{\partial \tilde{r}_3} \frac{\partial \tilde{r}_3}{\partial r_j}, \quad j = 1, 2, 3,$$

which, from Equation A1, can be given by

$$\frac{\partial f(\mathbf{r})}{\partial r_j} = v_{j1} \frac{\partial \tilde{f}(\tilde{\mathbf{r}})}{\partial \tilde{r}_1} + v_{j2} \frac{\partial \tilde{f}(\tilde{\mathbf{r}})}{\partial \tilde{r}_2} + v_{j3} \frac{\partial \tilde{f}(\tilde{\mathbf{r}})}{\partial \tilde{r}_3}, \quad j = 1, 2, 3. \quad (\text{A2})$$

Now, by deriving $\frac{\partial f(\mathbf{r})}{\partial r_j}$ (Eq. A2) with respect to the i th element r_i of the position vector \mathbf{r} (Eq. 1), we obtain:

$$\begin{aligned}
\frac{\partial^2 f(\mathbf{r})}{\partial r_i \partial r_j} &= v_{j1} \frac{\partial}{\partial r_i} \left(\frac{\partial \tilde{f}(\tilde{\mathbf{r}})}{\partial \tilde{r}_1} \right) + v_{j2} \frac{\partial}{\partial r_i} \left(\frac{\partial \tilde{f}(\tilde{\mathbf{r}})}{\partial \tilde{r}_2} \right) + v_{j3} \frac{\partial}{\partial r_i} \left(\frac{\partial \tilde{f}(\tilde{\mathbf{r}})}{\partial \tilde{r}_3} \right) \\
&= v_{j1} \left(\frac{\partial^2 \tilde{f}(\tilde{\mathbf{r}})}{\partial \tilde{r}_1 \partial \tilde{r}_1} v_{i1} + \frac{\partial^2 \tilde{f}(\tilde{\mathbf{r}})}{\partial \tilde{r}_2 \partial \tilde{r}_1} v_{i2} + \frac{\partial^2 \tilde{f}(\tilde{\mathbf{r}})}{\partial \tilde{r}_3 \partial \tilde{r}_1} v_{i3} \right) + \\
&+ v_{j2} \left(\frac{\partial^2 \tilde{f}(\tilde{\mathbf{r}})}{\partial \tilde{r}_1 \partial \tilde{r}_2} v_{i1} + \frac{\partial^2 \tilde{f}(\tilde{\mathbf{r}})}{\partial \tilde{r}_2 \partial \tilde{r}_2} v_{i2} + \frac{\partial^2 \tilde{f}(\tilde{\mathbf{r}})}{\partial \tilde{r}_3 \partial \tilde{r}_2} v_{i3} \right) + \\
&+ v_{j3} \left(\frac{\partial^2 \tilde{f}(\tilde{\mathbf{r}})}{\partial \tilde{r}_1 \partial \tilde{r}_3} v_{i1} + \frac{\partial^2 \tilde{f}(\tilde{\mathbf{r}})}{\partial \tilde{r}_2 \partial \tilde{r}_3} v_{i2} + \frac{\partial^2 \tilde{f}(\tilde{\mathbf{r}})}{\partial \tilde{r}_3 \partial \tilde{r}_3} v_{i3} \right) \\
&= \begin{bmatrix} v_{j1} & v_{j2} & v_{j3} \end{bmatrix} \tilde{\mathbf{F}}(\tilde{\mathbf{r}}) \begin{bmatrix} v_{i1} \\ v_{i2} \\ v_{i3} \end{bmatrix}, \tag{A3}
\end{aligned}$$

where $\tilde{\mathbf{F}}(\tilde{\mathbf{r}})$ is a 3×3 matrix whose ij -th element is $\frac{\partial^2 \tilde{f}(\tilde{\mathbf{r}})}{\partial \tilde{r}_i \partial \tilde{r}_j}$. From Equation A3, we obtain

$$\mathbf{F}(\mathbf{r}) = \mathbf{V} \tilde{\mathbf{F}}(\tilde{\mathbf{r}}) \mathbf{V}^\top, \tag{A4}$$

- 5 where $\mathbf{F}(\mathbf{r})$ is a 3×3 matrix whose ij -th element is $\frac{\partial^2 f(\mathbf{r})}{\partial r_i \partial r_j}$ and \mathbf{V} is defined by Eqs. 6 or 7, depending on the ellipsoid type. As one may noticed, the matrices $\mathbf{F}(\mathbf{r})$ and $\tilde{\mathbf{F}}(\tilde{\mathbf{r}})$ represent the Hessians of the functions $f(\mathbf{r})$ (Eq. 14) and $\tilde{f}(\tilde{\mathbf{r}})$, respectively. Besides, the depolarization tensor $\mathbf{N}(\mathbf{r})$ (Eq. 12) can be rewritten by using the matrix $\mathbf{F}(\mathbf{r})$ as follows

$$\mathbf{N}(\mathbf{r}) = -\frac{1}{4\pi} \mathbf{F}(\mathbf{r}). \tag{A5}$$

By properly using the orthogonality of the matrix \mathbf{V} , we may rewrite Equation A4 as follows:

$$10 \quad \tilde{\mathbf{F}}(\tilde{\mathbf{r}}) = \mathbf{V}^\top \mathbf{F}(\mathbf{r}) \mathbf{V}. \tag{A6}$$

Finally, by multiplying both sides of Equation A6 by $-\frac{1}{4\pi}$ and using Equation A5, we conclude that

$$\tilde{\mathbf{N}}(\tilde{\mathbf{r}}) = \mathbf{V}^\top \mathbf{N}(\mathbf{r}) \mathbf{V}. \tag{A7}$$

Appendix B: Parameter λ and its spatial derivatives

- Here, we follow the reasoning presented by Webster (1904) for analysing the parameter λ which defines triaxial, prolate and
15 oblate ellipsoids.

B1 Parameter λ defining triaxial ellipsoids

Let us consider an ellipsoid with semi-axes a , b , c oriented along the \tilde{x} -, \tilde{y} -, and \tilde{z} -axis, respectively, of its local coordinate system, where $a > b > c > 0$. This ellipsoid is defined by the following equation:

$$\frac{\tilde{x}^2}{a^2} + \frac{\tilde{y}^2}{b^2} + \frac{\tilde{z}^2}{c^2} = 1. \tag{B1}$$

A quadric surface (e.g., ellipsoid, hyperboloid of one sheet or hyperboloid of two sheets) which is confocal with the ellipsoid defined in Equation B1 can be described as follows:

$$\frac{\tilde{x}^2}{a^2 + u} + \frac{\tilde{y}^2}{b^2 + u} + \frac{\tilde{z}^2}{c^2 + u} = 1, \quad (\text{B2})$$

where u is a real number. Equation B2 represents an ellipsoid for u satisfying the condition

$$5 \quad u + c^2 > 0. \quad (\text{B3})$$

Given a , b , c , and a u satisfying B3, we may use B2 for determining a set of points (x, y, z) lying on the surface of an ellipsoid which is confocal with that one defined in Equation B1. Now, consider the problem of determining the ellipsoid which is confocal with that one defined in B1 and pass through a particular point $(\tilde{x}, \tilde{y}, \tilde{z})$. This problem consists in determining the real number u that, given a , b , c , \tilde{x} , \tilde{y} , and \tilde{z} , satisfies Equation B2 and the condition expressed by Equation B3. By rearranging

10 Equation B2, we obtain the following cubic equation for u :

$$p(u) = (a^2 + u)(b^2 + u)(c^2 + u) - (b^2 + u)(c^2 + u)\tilde{x}^2 - (a^2 + u)(c^2 + u)\tilde{y}^2 - (a^2 + u)(b^2 + u)\tilde{z}^2. \quad (\text{B4})$$

This cubic equation shows that:

$$u = \begin{cases} d \rightarrow \infty & , \quad p(u) > 0 \\ -c^2 & , \quad p(u) < 0 \\ -b^2 & , \quad p(u) > 0 \\ -a^2 & , \quad p(u) < 0 \end{cases}. \quad (\text{B5})$$

Notice that, according to B5, the smaller, intermediate and largest roots of the cubic equation $p(u)$ (Eq. B4) are located, respectively, in the intervals $[-a^2, -b^2]$, $[-b^2, -c^2]$ and $[-c^2, \infty[$. Remember that we are interested in a u satisfying the condition expressed by Equation B3. Consequently, according to the signal analysis shown in Equation B5, we are interested in the largest root λ of the cubic equation $p(u)$ (Eq. B4).

From Equation B4, we obtain a simpler one given by

$$p(u) = u^3 + p_2 u^2 + p_1 u + p_0, \quad (\text{B6})$$

20 where

$$p_2 = a^2 + b^2 + c^2 - \tilde{x}^2 - \tilde{y}^2 - \tilde{z}^2, \quad (\text{B7})$$

$$p_1 = b^2 c^2 + a^2 c^2 + a^2 b^2 - (b^2 + c^2)\tilde{x}^2 - (a^2 + c^2)\tilde{y}^2 - (a^2 + b^2)\tilde{z}^2 \quad (\text{B8})$$

and

$$25 \quad p_0 = a^2 b^2 c^2 - b^2 c^2 \tilde{x}^2 - a^2 c^2 \tilde{y}^2 - a^2 b^2 \tilde{z}^2. \quad (\text{B9})$$

Finally, from Eqs. B7, B8 and B9, the largest root λ of $p(u)$ (Eq. B6) can be calculated as follows (Weisstein, 2017):

$$\lambda = 2\sqrt{-Q}\cos\left(\frac{\theta}{3}\right) - \frac{p_2}{3}, \quad (\text{B10})$$

where

$$\theta = \cos^{-1}\left(\frac{R}{\sqrt{Q^3}}\right), \quad (\text{B11})$$

5

$$Q = \frac{3p_1 - p_2^2}{9} \quad (\text{B12})$$

and

$$R = \frac{9p_1p_2 - 27p_0 - 2p_2^3}{54}. \quad (\text{B13})$$

B2 Parameter λ defining prolate and oblate ellipsoids

- 10 Let us now consider a prolate ellipsoid with semi-axes a, b, c oriented along the \tilde{x} -, \tilde{y} -, and \tilde{z} -axis, respectively, of its local coordinate system, where $a > b = c > 0$. In this case, the Equation defining the surface of the ellipsoid is obtained by substituting $c = b$ in Equation B1. Consequently, the equation defining the respective confocal quadric surface is given by

$$\frac{\tilde{x}^2}{a^2 + u} + \frac{\tilde{y}^2 + \tilde{z}^2}{b^2 + u} = 1 \quad (\text{B14})$$

and the new condition that must be fulfilled by the variable u so that Equation B14 represent an ellipsoid is

$$15 \quad u + b^2 > 0. \quad (\text{B15})$$

Similarly to the case of a triaxial ellipsoid presented in the previous subsection, we are interested in determining the real number u that, given $a, b, \tilde{x}, \tilde{y}$, and \tilde{z} , satisfies Equation B14 and the condition expressed by Equation B15. From Equation B14, we obtain the following quadratic equation for u :

$$p(u) = (a^2 + u)(b^2 + u) - (b^2 + u)\tilde{x}^2 - (a^2 + u)(\tilde{y}^2 + \tilde{z}^2). \quad (\text{B16})$$

- 20 This equation shows that

$$u = \begin{cases} d \rightarrow \infty & , \quad f(\rho) > 0 \\ -b^2 & , \quad f(\rho) < 0 \\ -a^2 & , \quad f(\rho) > 0 \end{cases} \quad (\text{B17})$$

and, consequently, that its two roots lie in the intervals $[-a^2, -b^2]$ and $[-b^2, \infty[$. Therefore, according to the condition established by Equation B15 and the signal analysis shown in Equation B17, we are interested in the largest root λ of the quadratic equation $p(u)$ (Eq. B16).

By properly manipulating Equation B16, we obtain a simpler one given by

$$p(u) = u^2 + p_1 u + p_0, \quad (\text{B18})$$

where

$$p_1 = a^2 + b^2 - \tilde{x}^2 - \tilde{y}^2 - \tilde{z}^2 \quad (\text{B19})$$

5 and

$$p_0 = a^2 b^2 - b^2 \tilde{x}^2 - a^2 (\tilde{y}^2 + \tilde{z}^2). \quad (\text{B20})$$

Finally, by using Eqs. B19 and B20, the largest root λ of $p(u)$ (Eq. B18) can be easily calculated as follows:

$$\lambda = \frac{-p_1 + \sqrt{p_1^2 - 4p_0}}{2}. \quad (\text{B21})$$

In the case of oblate ellipsoids, the procedure for determining the parameter λ is very similar to that one for prolate ellipsoids.

- 15 The semi-axes a, b, c of oblate ellipsoids are defined so that $b = c > a > 0$ and the condition that must be fulfilled by the variable u is $u + a^2 > 0$. In this case, the two roots of the resulting quadratic equation lie in the intervals $[-b^2, -a^2]$ and $[-a^2, \infty[$. Consequently, we are still interested in the largest root of the quadratic equation for the variable u , which is also calculated by using Equation B21.

B3 Spatial derivative of the parameter λ

- 15 The magnetic modelling of triaxial, prolate or oblate ellipsoids requires not only the parameter λ defined by Eqs. B10 and B21, but also its derivatives with respect to the spatial coordinates \tilde{x}, \tilde{y} , and \tilde{z} . Fortunately, the spatial derivatives of the parameter λ can be calculated in a very similar way for all ellipsoid types.

Let us first consider a triaxial ellipsoid. In this case, the spatial derivatives of λ are given by

$$\frac{\partial \lambda}{\partial \tilde{r}_j} = \frac{\frac{2\tilde{r}_j}{(e_j^2 + \lambda)}}{\left(\frac{\tilde{x}}{a^2 + \lambda}\right)^2 + \left(\frac{\tilde{y}}{b^2 + \lambda}\right)^2 + \left(\frac{\tilde{z}}{c^2 + \lambda}\right)^2}, \quad j = 1, 2, 3, \quad (\text{B22})$$

- 20 where $\tilde{r}_1 = \tilde{x}$, $\tilde{r}_2 = \tilde{y}$, $\tilde{r}_3 = \tilde{z}$, $e_1 = a$, $e_2 = b$, and $e_3 = c$. This equation can be determined directly from equation B2. The spatial derivatives of λ in the case of prolate or oblate ellipsoids can also be calculated by using Equation B22, but with semi-axes $a > b = c$ or $b = c > a$, respectively.

Author contributions. TEXT

Acknowledgements. TEXT

References

- Blakely, R. J.: Potential theory in gravity and magnetic applications, Cambridge University Press, 1996.
- Chang, H.: Fields external to open-structure magnetic devices represented by ellipsoid or spheroid, *British Journal of Applied Physics*, 12, 160, 1961.
- 5 Clark, D., Saul, S., and Emerson, D.: Magnetic and gravity anomalies of a triaxial ellipsoid, *Exploration Geophysics*, 17, 189–200, 1986.
- Dirichlet, L. P. G.: Sur un nouvelle methode pour la determination des integrales multiples, *Journal de mathématiques pures et appliquées*, 1, 164–168, http://sites.mathdoc.fr/JMPA/PDF/JMPA_1839_1_4_A11_0.pdf, 1839.
- DuBois, H.: *The Magnetic Circuit in Theory and Practice*, Longmans, Green, and Co., 1896.
- Emerson, D. W., Clark, D., and Saul, S.: Magnetic exploration models incorporating remanence, demagnetization and anisotropy: HP 41C
- 10 handheld computer algorithms, *Exploration Geophysics*, 16, 1–122, 1985.
- Kellogg, O. D.: *Foundations of Potential Theory*, Frederick Ungar Publishing Company, 1929.
- Lowes, F. J.: The Torque on a Magnet, *Proceedings of the Royal Society of London. Series A, Mathematical and Physical Sciences*, 337, 555–567, <http://www.jstor.org/stable/78532>, 1974.
- Maxwell, J. C.: *A treatise on electricity and magnetism*, vol. 2, Clarendon press, 1873.
- 15 Osborn, J.: Demagnetizing factors of the general ellipsoid, *Physical review*, 67, 351, 1945.
- Peake, H. J. and Davy, N.: The magnetic fields produced by uniformly magnetized ellipsoids of revolution, *British Journal of Applied Physics*, 4, 207, 1953.
- Peirce, B.: *Elements of the Theory of the Newtonian Potential Function*, Cornell University Library historical math monographs, Ginn, 1902.
- Poisson, S. D.: Second mémoire sur ta théorie du magnétism, *Mémoires de l'Académie des sciences de l'Institut de France*, pp. 488–533,
- 20 1824.
- Searle, S. R.: *Matrix Algebra Useful for Statistics*, Wiley-Interscience, 1982.
- Solivérez, C. E.: Magnetostatics of anisotropic ellipsoidal bodies, *IEEE Transactions on Magnetism*, 17, 1363–1364, doi:10.1109/TMAG.1981.1061200, 1981.
- Solivérez, C. E.: Campos eléctricos generados por elipsoides uniformemente polarizados, *Revista mexicana de física E*, 54, 203 – 207, 2008.
- 25 Stoner, E. C.: The demagnetizing factors for ellipsoids, *The London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science*, 36, 803–821, 1945.
- Stratton, J. A.: *Electromagnetic Theory*, Wiley-IEEE Press, 2007.
- Tejedor, M., Rubio, H., Elbaile, L., and Iglesias, R.: External fields created by uniformly magnetized ellipsoids and spheroids, *IEEE transactions on magnetism*, 31, 830–836, 1995.
- 30 Thomson, W. and Tait, P.: *Treatise on Natural Philosophy, Part II,* Cambridge University Press, Cambridge, 1879.
- Webster, A.: *The Dynamics of Particles and of Rigid, Elastic, and Fluid Bodies: Being Lectures on Mathematical Physics*, B. G. Teubners Sammlung von Lehrbüchern auf dem Gebiete der mathematischen Wissenschaften, B.G. Teubner, 1904.
- Weisstein, E. W.: Cubic Formula: From MathWorld – A Wolfram Web Resource, <http://mathworld.wolfram.com/CubicFormula.html>, [Online; accessed 2017-01-23], 2017.