

# Working Paper - Design and Analysis of Multivariate Experiments

Diego Vallarino

15/12/2021

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
1.1	Cargamos los datos . . . . .	1
1.2	Descriptive data analysis . . . . .	1
1.3	Linear Model for the Experiment . . . . .	2
1.4	Model Hypothesis . . . . .	3
1.5	Significant factors . . . . .	3
1.6	Variance components . . . . .	3
1.7	Perform multiple comparisons . . . . .	4
1.8	Confidence Intervals with 95% . . . . .	4

---

## 1 Introduction

Certain experimental conditions give rise to designs that contain crossed and nested factors. In this exercise we can identify a design with three factors, a fixed *Spectrometer* factor with 3 levels crossed with a random *Plots* factor with 3 levels, and a third *Samples* factor with 2 levels as well. random that is crossed with *Spectrometer* but nested with *Plots*.

Therefore, the analysis of the experiment is as follows:

### 1.1 Cargamos los datos

### 1.2 Descriptive data analysis

##	parcela	muestra	espectrometro	medida
##	Min. :1	Min. :1.0	Min. :1	Min. :3.700
##	1st Qu.:1	1st Qu.:1.0	1st Qu.:1	1st Qu.:3.778
##	Median :2	Median :1.5	Median :2	Median :3.860
##	Mean :2	Mean :1.5	Mean :2	Mean :3.841
##	3rd Qu.:3	3rd Qu.:2.0	3rd Qu.:3	3rd Qu.:3.881
##	Max. :3	Max. :2.0	Max. :3	Max. :3.988

```
## 'data.frame': 36 obs. of 4 variables:
## $ parcela : int 1 1 1 1 1 1 1 1 1 1 ...
## $ muestra : int 1 1 2 2 1 1 2 2 1 1 ...
## $ espectrometro: int 1 1 1 1 2 2 2 2 3 3 ...
## $ medida : num 3.83 3.87 3.82 3.85 3.93 ...

## vars n mean sd median trimmed mad min max range skew
## parcela* 1 36 2.00 0.83 2.00 2.00 1.48 1.0 3.00 2.00 0.00
## muestra* 2 36 1.50 0.51 1.50 1.50 0.74 1.0 2.00 1.00 0.00
## espectrometro* 3 36 2.00 0.83 2.00 2.00 1.48 1.0 3.00 2.00 0.00
## medida 4 36 3.84 0.08 3.86 3.84 0.12 3.7 3.99 0.29 0.15
## kurtosis se
## parcela* -1.58 0.14
## muestra* -2.05 0.08
## espectrometro* -1.58 0.14
## medida -0.98 0.01
```

### 1.3 Linear Model for the Experiment

The model is as follows:

```
modelo<-lm(medida ~ espectrometro + r(parcela) + espectrometro:r(parcela) + r(muestra)%in%r(parcela)
+ espectrometro:(r(muestra)%in%r(parcela)), datos)
```

A continuación se presentan los calculos:

```
##
## Shapiro-Wilk normality test
##
## data: modelo$residuals
## W = 0.88116, p-value = 0.001084

## Analysis of variance (unrestricted model)
## Response: medida
##
##          Mean Sq Sum Sq Df F value Pr(>F)
## espectrometro    0.0626  0.125  2    9.06 0.0327
## parcela          0.0175  0.035  2    2.14 0.2397
## espectrometro:parcela 0.0069  0.028  4    2.71 0.1326
## parcela:muestra    0.0038  0.011  3    1.50 0.3076
## espectrometro:parcela:muestra 0.0025  0.015  6    7.38 0.0004
## Residuals        0.0003  0.006 18      -    -
##
##          Err.term(s) Err.df  VC(SS)
## 1 espectrometro          (3)      4   fixed
## 2 parcela      (3) + (4) - (5)   3.74 0.000779
## 3 espectrometro:parcela          (5)      6 0.001089
## 4 parcela:muestra          (5)      6 0.000211
## 5 espectrometro:parcela:muestra (6)     18 0.001101
## 6 Residuals          -      - 0.000345
## (VC = variance component)
##
##          Expected mean squares
## espectrometro      (6) + 2 (5) + 4 (3) + 12 Q[1]
```

```

## parcela (6) + 2 (5) + 6 (4) + 4 (3) + 12 (2)
## espectrometro:parcela (6) + 2 (5) + 4 (3)
## parcela:muestra (6) + 2 (5) + 6 (4)
## espectrometro:parcela:muestra (6) + 2 (5)
## Residuals (6)

```

## 1.4 Model Hypothesis

For the Spectrometer Fixed factor, A:

$$H_0 : \alpha_1 = \alpha_2 = \alpha_3 = 0$$

$$H_1 : \alpha_i \neq 0$$

For Random Factor Plots, B

$$H_0 : \sigma_B^2 \neq 0$$

$$H_1 : \sigma_B^2 > 0$$

For the interaction Spectrometer and Plot, AB

$$H_0 : \sigma_{AB}^2 \neq 0$$

$$H_1 : \sigma_{AB}^2 > 0$$

For the random factor Plots nested to Samples, C(B):

$$H_0 : \sigma_C^2 \neq 0$$

$$H_1 : \sigma_C^2 > 0$$

For the Spectrometer  $\times$  Samples interaction nested to Plots, AC(B)

$$H_0 : \sigma_{AC}^2 \neq 0$$

$$H_1 : \sigma_{AC}^2 > 0$$

## 1.5 Significant factors

According to the evidence presented with the ANOVA analysis, only the fixed factor **Spectrometer** is significant, together with its *interaction* with the Plot.

## 1.6 Variance components

- a) Error variance: 0.0003
- b) Variance of the parcel factor: 0.000779
- c) Variance of the spectrometer:plot interaction: 0.001089
- d) Variance of the plot:sample interaction: 0.000211
- e) Variance of the spectrometer:plot:sample interaction: 0.001101

The estimate of the variance of the response variable is  $= 0.0003 + 0.000779 + 0.001089 + 0.000211 + 0.001101 = \mathbf{0.00348}$  and therefore, the percentage of the variance components are:

- a) Error variance: 9%
- b) Variance of the parcel factor: 22%
- c) Variance of the spectrometer:plot interaction: 31%
- d) Variance of plot:sample interaction: 6%
- e) Variance of the spectrometer:plot:sample interaction: 32%

We conclude that the *spectrometer:plot:sample interaction factor* is the main cause of variability in the response (32%), followed by the *spectrometer:plot factor* (31%).

## 1.7 Perform multiple comparisons

```
HSD.test(modelo, "fert", console = TRUE)
```

```
## Name: fert
## espectrometro parcela muestra
```

```
HSD.test(modelo,"espectrometro", group=TRUE,console=TRUE)
```

```
##
## Study: modelo ~ "espectrometro"
##
## HSD Test for medida
##
## Mean Square Error: 0.0003449722
##
## espectrometro, means
##
##      medida      std  r   Min   Max
## 1 3.802667 0.06176986 12 3.720 3.887
## 2 3.923917 0.04659976 12 3.872 3.988
## 3 3.795333 0.05199184 12 3.700 3.872
##
## Alpha: 0.05 ; DF Error: 18
## Critical Value of Studentized Range: 3.609304
##
## Minimum Significant Difference: 0.01935196
##
## Treatments with the same letter are not significantly different.
##
##      medida groups
## 2 3.923917      a
## 1 3.802667      b
## 3 3.795333      b
```

We can identify that **spectrometers 1 and 3** have similar performances, while spectrometer 2 has a different performance than the other two. Therefore, we can identify 2 homogeneous groups of spectrometers.

## 1.8 Confidence Intervals with 95%

```
HSD.test(modelo,"espectrometro", group=FALSE,console=TRUE)
```

```
##
## Study: modelo ~ "espectrometro"
##
## HSD Test for medida
##
## Mean Square Error: 0.0003449722
##
```

```

## espectrometro, means
##
##      medida      std  r   Min   Max
## 1 3.802667 0.06176986 12 3.720 3.887
## 2 3.923917 0.04659976 12 3.872 3.988
## 3 3.795333 0.05199184 12 3.700 3.872
##
## Alpha: 0.05 ; DF Error: 18
## Critical Value of Studentized Range: 3.609304
##
## Comparison between treatments means
##
##      difference pvalue signif.      LCL      UCL
## 1 - 2 -0.121250000 0.0000      *** -0.14060196 -0.10189804
## 1 - 3  0.007333333 0.6063           -0.01201862  0.02668529
## 2 - 3  0.128583333 0.0000      ***  0.10923138  0.14793529

```