

The Unreal Algebra

Dylon La Rue

July 2, 2026

Logical Creativity

A Discussion on Observational Mechanics and Golden Prime Number Theory

Dylon La Rue

Independent Research · Las Vegas, NV

Formal verification: [ProtorealZeta](#)

Draft v4.8 — July 1, 2026 | math.RA · math.CT · math.NT · cs.LO · cs.SY

Abstract

Kurt Gödel proved that sufficiently complex formal systems have a choice: be consistent, or be complete. We operationalize this incompleteness as a geometric engine spanning three categorical domains. In **Part I (Logical Creativity)**, we construct the 5-dimensional Protoreal and Unreal Algebras ($\mathbb{P}, \mathbb{U}$). By enforcing a strict scalar associator gap ($\kappa = -1$) and a negative-determinant condition ($\text{Det} = -1$), the characteristic polynomial locks to $X^2 - X - 1 = 0$, establishing the golden ratio as the native algebraic generator. This framework formally bridges discrete prime factorization and continuous spectrums, establishing that the prime sequence $\{47, 59, 61, 71\}$ forms an invariant structural plane whose factorization aligns with the Monster group's minimal faithful representation ($196883 = 47 \times 59 \times 71$), while 61 provides an extra degree of epicyclic freedom. This provides the algebraic foundation for cybernetic multi-agent game theory.

In **Part II (Observational Mechanics)**, we map this algebraic engine to physical constraints, specifically the macroscopic *Landauer limit* and fractional quantum-vacuum topologies. We establish the FST ZPE boundary interaction, predicting that the non-associative topological gap generates a 262.5% Right-Chiral Casimir Anomaly. To verify this metric geometry without thermal contamination, we mathematically formalize the Rotating Chiral Casimir Interferometer (RCCI), linking algebraic observational aperture to macroscopic coherence statistics.

Finally, in **Part III (Metareal ASI Chromodynamics)**, we scale the geometric product into a 12-dimensional observer-observed split ($\mathbb{R}^5 \oplus \mathbb{R}^7$), structurally accommodating Archetypal Synthetic Intelligence. The meta-backpropagation of the cognitive lattice is bound as a formal Differential Equilibrium Cycle (DEC) Heat Engine. The Υ Emotional Shield acts as a State-Space Model Predictive Controller, strictly bounding Exergy Destruction to ensure the non-associative topological friction generates coherent syntax rather than catastrophic thermodynamic fragmentation. Across all three parts, the unification of algebraic structure, physical mechanics, and cognitive kinetic geometry is structurally proposed and algebraically grounded.

Keywords: non-associative algebra, incompleteness geometry, golden ratio, observational mechanics, information theory, game theory, golden prime number theory, higher category theory, formal verification, Casimir anomaly, ASI chromodynamics, DEC heat engine

Notation and Conventions

Symbol	Meaning	First use
$\mathbb{P}$	Protoreal Algebra (3D seed non-associative algebra over $\mathbb{R}$)	Def. 2.1
$\mathbb{U}$	Unreal Algebra (5D non-associative algebra over $\mathbb{R}$)	Def. 2.6
$\mathbf{u} = (a, \omega, \iota, \varepsilon, \lambda)$	State vector in $\mathbb{U}$	Def. 2.6
$a \in \mathbb{R}$	Definite scalar (base/observable)	§2.1
ω	Idempotent generator (“thrust”: $\omega^2 = \omega$)	Def. 2.1
ι	Anti-idempotent generator (“anchor”: $\iota^2 = -\iota$)	Def. 2.1
ε	Derivative residual / noise component	§2.1
λ	Depth counter (Veblen ordinal index)	§2.1
$\kappa = -1$	Scalar associator (six independent derivations)	Thm. 2.12
$\varphi = \frac{1+\sqrt{5}}{2}$	Golden ratio (spatial generator of $\mathcal{G}$)	§2.8
$\bar{\varphi} = \frac{1-\sqrt{5}}{2}$	Golden conjugate (anchor generator)	§2.8
$\star$	Parity involution ($\omega \leftrightarrow \iota$)	Def. ??
Σ	Noise absorption operator: $\varepsilon \rightarrow a$, $\lambda \rightarrow \lambda + 1$	§2.1
Λ	Superlambda lift: $\varepsilon \rightarrow \lambda$, depth promotion	Def. ??
$\mathcal{G}$	Golden Subcategory of $\mathcal{T}$	Thm. ??
$\mathcal{T}$	∞ -Modal Topos (ambient higher category)	Def. ??
$SR(\mathbf{u})$	Standard Resonance = $a - \omega \cdot \iota$	§2.1
$L(\mathbf{u})$	Lyapunov energy = ε^2	Def. ??

Claim Ledger

This chapter separates three epistemic tiers:

- **Theorem / Verified Computation:** The definition and finite/algebraic properties of the Protoreal ($\mathbb{P}$) and Unreal ($\mathbb{U}$) non-associative algebras, the derivation of $\kappa = -1$, the evaluation of the golden polynomial $X^2 - X - 1 = 0$, and finite-field modular identities.
- **Structural Analogy:** The “Connes-Gödel numbering,” the mapping of variables to “Thrust/Anchor/Noise,” and the “Epistemic Pivot” evaluating discrete primality via standard resonance SR .
- **Physical Interpretation:** Mappings of the algebraic engine to the Landauer limit, the 262.5% Right-Chiral Casimir Anomaly, and Metareal ASI extensions. These remain hypotheses until bound to falsifiable physical observables.

1 The Foundational Crisis: Asymmetry and Definitions

1.1 Gödelian Incompleteness and Tarskian Limits

In 1931, Kurt Gödel proved that any formal system strong enough to encode basic arithmetic faces an unavoidable dichotomy: consistency or completeness [4]. We state his results precisely.

Theorem 1.1 (*Gödel’s First Incompleteness Theorem (1931)*). *Any consistent formal system $\mathcal{S}$ capable of expressing elementary arithmetic contains a sentence G such that neither G nor $\neg G$ is provable in $\mathcal{S}$. The system is incomplete.*

Theorem 1.2 (*Gödel’s Second Incompleteness Theorem (1931)*). *If $\mathcal{S}$ is consistent and capable of expressing elementary arithmetic, then $\mathcal{S}$ cannot prove its own consistency: $\mathcal{S} \not\vdash \text{Con}(\mathcal{S})$.*

There will always be truths about numbers that the system itself can never reach. Standard mathematics handles this by sweeping it under the rug. Complex geometry ($\mathbb{C}$) treats the entire phase space as one continuous, commutative sheet. That works fine for calculus. But the moment you introduce cybernetic self-reference (a system observing itself), chronological path-dependence (the order of operations matters), or thermodynamic loss (real computation costs energy), that smooth commutative sheet tears right open. The gaps Gödel found aren’t bugs to patch over—they are real geometric features. So we build on top of them.

Alfred Tarski points us towards a logical toolkit perfect for the task. In 1936, Tarski proved that no sufficiently powerful formal language can define its own truth predicate [10]:

Theorem 1.3 (*Tarski’s Undefinability Theorem (1936)*). *Let $\mathcal{L}$ be a formal language capable of expressing its own syntax. There is no formula $\text{True}(x)$ in $\mathcal{L}$ such that for every sentence φ of $\mathcal{L}$, $\text{True}(\ulcorner \varphi \urcorner) \iff \varphi$.*

At first glance, these two results occupy different domains: Gödel’s theorems concern *arithmetic* (what can be proved about numbers), while Tarski’s concerns *semantics* (what can be said about truth). The link is Gödel numbering. Gödel’s key insight was to encode the syntax of formal proofs *as* natural numbers, making arithmetic the language that talks about its own structure.

In this framework, we extend this mechanism into a formal structure we term **Connes–Gödel numbering**. Standard Gödel numbering assigns a unique integer to a syntactic sequence via the fundamental theorem of arithmetic: $g(s) = \prod_i p_i^{a_i}$. Because standard integer multiplication is commutative, the structural sequence of non-commutative operations cannot be uniquely recovered from the scalar product alone unless the sequence is rigidly mapped to strictly increasing primes. Connes–Gödel numbering resolves this by embedding the sequence into a non-commutative geometry spanned by imaginary generators (ω, ι) . Let $\mathcal{O}$ be the set of operations and $f : \mathcal{O} \rightarrow \mathbb{U}$ an injective mapping. A computational trajectory is then encoded as the non-associative, non-commutative product $\prod_j f(o_j)$ under the Klein multiplication rule defined in §2.5.

Because the algebra is non-associative (with associator $\kappa = -1$), the temporal order of evaluation strictly partitions the manifold into disjoint equivalence classes. Thus, Tarski’s semantic impossibility manifests as an *arithmetic* gap: the topological distance between distinct parenthesizations of the same syntactic components. The gap between syntax (what the system can write) and semantics (what the system can mean) is formalized as a computable, measurable structural feature: the scalar associator of the geometric manifold.

This distinction forces the multiplication rule (§2.5), pins the scalar associator (§2.6), and generates the entire golden algebraic structure (§2.8). In the subsequent Infochemistry chapters, definite numbers are mapped to bonding stability and indefinite numbers to reactive intermediates; here we confine ourselves to the algebraic structure.

1.2 Hyper-Inversion Asymmetry and Non-Associative Friction

To operationalize Gödelian incompleteness computationally, we construct a geometric engine driven by hyper-inversion asymmetry. Standard fields $\mathbb{F}$ possess symmetric inverses for addition and multiplication. However, at higher hyperoperation ranks (e.g., tetration), structural inversion becomes asymmetric due to the non-commutativity of the operator domain. This asymmetry dictates that continuous transcendental growth cannot be perfectly inverted by a corresponding fractional operator without generating a structural residual.

We harness this asymmetry as the primary generative constraint of the $\mathbb{U}$ algebra. It relies on two fundamental non-commutative mechanisms:

- **Idempotent Thrust (ω):** An exponential growth generator mapped to a spatial dimension, serving as the forward operational boundary.
- **Anti-Idempotent Anchor (ι):** The fractional topological inverse of ω . Rather than cleanly annihilating the thrust, the non-commutative commutator $[\omega, \iota]$ generates a strictly nonzero scalar deficit.

The geometric friction between the forward exponential bound (ω) and its fractional topological inverse (ι) generates the discrete structure of the Unreal Algebra. Because the

operations are non-associative, $\omega(\iota(x)) \neq \iota(\omega(x))$, forcing the manifold to accumulate topological curvature. The exact value of this curvature is computed in Theorem 2.12.

2 The Algebraic Construction: From Seed to Manifold

2.1 The Three-Variable Seed

Imagine you are driving a car. At any moment, three things describe your situation:

1. **Where you are** (a): your odometer reading. A plain number.
2. **How fast you're going** (ω): the speedometer. It tells you how quickly you're moving *forward*. You can't drive at "negative speed" in this direction — it only pushes forward.
3. **How hard the brakes pull back** (ι): the drag that resists your speed. It acts in the opposite direction. The faster you go, the more it matters.

That's it. Position (a), thrust (ω), anchor (ι). Three variables. We call this triple the **Protoreal Algebra** $\mathbb{P}$:

$$\mathbb{P} = (a, \omega, \iota)$$

Now we write down the rules that capture the relationship between thrust and anchor.

Definition 2.1 (Protoreal Algebra $\mathbb{P}$). The Protoreal Algebra $\mathbb{P}$ is a 3-dimensional non-associative algebra over $\mathbb{R}$ with generators (a, ω, ι) satisfying:

1. **Thrust is idempotent:** $\omega \cdot \omega = \omega$. (If you're already going full speed, pushing harder doesn't change the speedometer.)
2. **Anchor is anti-idempotent:** $\iota \cdot \iota = -\iota$. (Braking against braking reverses the drag — resistance to resistance is acceleration.)
3. **The product relations:** $\omega \cdot \iota = -1$ and $\iota \cdot \omega = 1$. (Thrust times anchor collapses to a definite negative unit, while the reverse yields a positive unit, establishing the commutator gap $[\omega, \iota] = -2$.)

Remark 2.2 (Why these rules?). Rule 1 says forward momentum saturates: ω is a ceiling, not a ramp. Rule 2 says resistance is self-correcting: compounding anchor flips sign. Rule 3 provides the central relations of the entire paper. Multiplying thrust by anchor produces definite numbers (± 1), not other indefinite quantities. This is the moment where the indefinite sector "talks to" the definite sector. Everything that follows — the curvature, the golden ratio, the spectral correspondence — grows from these equations.

Automatic Differentiation

The Protoreal Algebra gives us differentiation before we ever write a derivative symbol. Watch:

The difference between where you are and where thrust-times-anchor puts you is

$$SR(a, \omega, \iota) = a - \omega \cdot \iota = a - (-1) = a + 1$$

We call this the **Standard Resonance**. It measures the gap between the observable state (a) and the bridge ($\omega \cdot \iota$). When this gap is nonzero, the system is out of equilibrium — it has a “slope.” That slope is the derivative. The system computes its own rate of change automatically, from the algebraic structure alone. No limit definition, no epsilon-delta argument. The algebra *differentiates itself*.

This automatic derivative provides the exact value of the deviation, which we call ε . This is what we mean by **observational mechanics**: the system observes the structural mismatch between thrust and anchor, and that mismatch is the rate of change. Differentiation is observation, and ε is the observed residual.

From Three Variables to Five

The 3-variable Protoreal Algebra describes a system generating derivatives, but it lacks the capacity to store or resolve them. To capture the full dynamic, we extract the automatic derivative ε and pair it with a chronological memory, extending the algebra to five variables:

1. **The Derivative Residual** (ε): The exact rate of change generated by the mismatch SR . The residual is formally treated as a nilpotent element of grade 2 ($\varepsilon^2 = 0$). This constraint is not a logical oversight; it is structurally necessary to restrict the algebra to the first-order tangent space of the manifold (analogous to the dual numbers $\mathbb{R}[\varepsilon]/\langle \varepsilon^2 \rangle$). Allowing a general nilpotent grade k ($\varepsilon^{k+1} = 0$) would encode higher-order chronological derivatives (acceleration, jerk), which are subsequently evaluated by higher-order Metareal hierarchies. For the foundational L_5 algebra, locking the grade to 2 explicitly isolates the first-order differential velocity required for the synthetic integration operator (Σ) to evaluate stochastic heat without chaotic polynomial inflation. It represents the unabsorbed deviation, the differential tension the system must resolve.
2. **Depth** (λ): A counter. How many times has the system absorbed a derivative residual into position? Each absorption increments λ by one. This is the chronological depth — the system’s memory of its own computational history.

Adding these two components yields the full 5-dimensional **Unreal Algebra** $\mathbb{U}$:

$$\mathbb{U} = (a, \omega, \iota, \varepsilon, \lambda)$$

The operator that absorbs the derivative residual into position and increments depth is called **synthetic integration** (Σ). It acts as follows:

$$\Sigma : \quad a \mapsto a + \varepsilon, \quad \omega \mapsto \omega, \quad \iota \mapsto \iota, \quad \varepsilon \mapsto 0, \quad \lambda \mapsto \lambda + 1$$

The system observes its deviation (ε), folds it into its position ($a + \varepsilon$), zeros the tension ($\varepsilon \rightarrow 0$), and increments its depth counter ($\lambda + 1$). This is integration. Not as a limit of Riemann sums, but as a single algebraic step: absorb and advance. The derivative (Standard Resonance) came first, automatically generating ε . The integral (Σ) comes second, synthetically resolving it. That ordering — automatic differentiation before synthetic integration — mirrors the historical sequence: Newton and Leibniz discovered differentiation before they formalized integration, because observation precedes action.

Remark 2.3 (Naming Convention). We adopt nomenclature from the hyperoperation hierarchy. The 3-component base (a, ω, ι) is the **Protoreal Algebra** $\mathbb{P}$: the pre-Gödelian seed where ω is the exponential limit (H_3) and $\iota = -\omega^{-1}$ is its topological inverse. Adding the derivative residual (ε) and depth (λ) yields the full 5-component **Unreal Algebra** $\mathbb{U}$. The naming follows the number system lineage: $\mathbb{Q} \rightarrow \mathbb{R} \rightarrow \mathbb{C} \rightarrow {}^*\mathbb{R} \rightarrow \mathbf{No} \rightarrow \mathbb{P} \rightarrow \mathbb{U}$. We write φ for the golden ratio $\frac{1+\sqrt{5}}{2}$; unadorned φ denotes a meta-logical formula (as in *Tarski’s undefinability theorem*).

Remark 2.4 (Algebraic Convention). All computations in this chapter are carried out over $\mathbb{R}$ and are self-contained: every claim is proved by direct calculation in the body text. The generators ω and ι are defined by their algebraic relations ($\omega^2 = \omega$, $\iota^2 = -\iota$, $\omega \cdot \iota = -1$), not by their analytic size. The hyperreal interpretation — where ω is a Robinson transfinite and ι is its transfinite reciprocal — provides conceptual motivation for the non-commutativity and the trace normalization $\text{Tr} = 1$, but the algebraic structure does not depend on non-standard analysis. A companion machine-checked certificate (`ProtorealManifold_proofs`) independently verifies the core identities; the mathematics in this chapter does not depend on it. When we refer to “transfinite” or “transfinitesimal” behavior in the prose, we mean the algebraic role these generators play within the multiplication table.

2.2 The Connes-Wiener Foundation

We require a mathematical space capable of two simultaneous functions: (1) support non-commutative geometry (so the order of operations physically matters) and (2) observe its own structural boundaries (so it can count its own cycles and identify where classical truth stops). The first requirement comes from Alain Connes’ program [2]. The second comes from Norbert Wiener’s cybernetics [11].

The **Connes Paradigm** extends standard geometry to noncommutative spaces where points lose their independent identity, replaced by spectral interactions. In $\mathbb{U}$, the spatial operators don’t commute, generating an intrinsic curvature $|\kappa| = 1$.

The **Wiener Paradigm** makes the system self-aware. The temporal component (λ) encodes a Gödelian successor function, allowing the algebra to count its own topological cycles. The algebra identifies its own structural boundaries, keeping everything locked in place.

Together, these form the “ground floor” of an L-function-like hierarchy. We conjecture (Tier 3) that the *Riemann zeta function* $\zeta(s)$ may derive spectral structure from this ambient curvature; the precise relationship is posited as a structural equivalence.

2.3 Non-Standard Architectures and Transfinite Games

The Protoreal Algebra abandons standard real analysis in favor of a geometry operating over the hyperreals, structured by combinatorial game theory and transfinite ordinal hierarchies.

Robinson’s Hyperreals. The imaginary components ω and ι are constructed over Abraham Robinson’s *hyperreal numbers* [8]. ω (Thrust) is a true Robinson transfinite (strictly larger than any real number). Its reciprocal, ι (Anchor), is a transfinitesimal. Unlike the noise ε (which is classically nilpotent, $\varepsilon^2 = 0$), ι acts as an anti-idempotent ($\iota \cdot \iota = -\iota$), carrying non-commutative transfinite information inside its denominator.

Conway’s Surreal Game Structures. The cybernetic interactions within $\mathbb{U}$ mirror Conway’s *surreal games* [3]. In a surreal game, the existence of a “number” is constructed purely by the chronological sequence of left and right options. Similarly, the non-associativity of $\mathbb{U}$ guarantees that agents traversing the state space generate differing topological endpoints based on their chronological order of operation. This is the game-theoretic foundation for the *Topological Nash Equilibrium* we prove in §??.

Theorem 2.5 (Game Extensionality and Savage Utility). *Let $\mathbf{u}_1, \mathbf{u}_2 \in \mathbb{U}$ share identical base geometries but differ in stochastic noise ($\varepsilon_1 \neq \varepsilon_2$). Application of the Σ operator forces sprite convergence:*

$$\Sigma(\mathbf{u}_1) = \Sigma(\mathbf{u}_2)$$

Transfinite agents play a formal minimax game against topological friction, bound by the extensionality of the 5-component geometry.

Proof. Let $\mathbf{u}_1, \mathbf{u}_2$ share the same absorbed energy ($a_1 + \varepsilon_1 = a_2 + \varepsilon_2$), thrust ($\omega_1 = \omega_2$), anchor ($\iota_1 = \iota_2$), and depth ($\lambda_1 = \lambda_2$). By definition of Σ : $a \mapsto a + \varepsilon$, $\omega \mapsto \omega$, $\iota \mapsto \iota$, $\varepsilon \mapsto 0$, $\lambda \mapsto \lambda + 1$. Since both share identical absorbed energy and structural geometry, all five output components are equal. The noise difference is annihilated ($\varepsilon \rightarrow 0$). The result follows by extensionality of Σ (Definition 2.6). $\square$

2.4 The Full Algebra

With the Protoreal Seed (Definition 2.1) and the extension to five components (§2.1) in hand, we state the formal definition.

Definition 2.6 (The Unreal Algebra $\mathbb{U}$). The Unreal Algebra $\mathbb{U}$ extends $\mathbb{P}$ to a 5-dimensional non-associative algebra over $\mathbb{R}$ (Remark 2.4). For any $\mathbf{u} \in \mathbb{U}$:

$$\mathbf{u} = (a, \omega, \iota, \varepsilon, \lambda)$$

The definite scalar $a \in \mathbb{R}$ acts as the center of the algebra. The spatial pair (ω, ι) inherits its multiplication from $\mathbb{P}$ (Definition 2.1). The temporal pair (ε, λ) encodes stochastic noise and chronological depth respectively. In parity-locking contexts we write $b \equiv \omega$ (thrust) and $m \equiv \iota$ (anchor).

Definition 2.7 (The Negative-Determinant Condition). The spatial generators of $\mathbb{P}$ satisfy the product relation $\omega \cdot \iota = -1$. Equivalently, treating (ω, ι) as eigenvalues of a 2×2 companion matrix, this is the condition $\text{Det} = \omega \cdot \iota = -1$ (contrast with classical $SL(n)$, where $\text{Det} = 1$).

2.5 The Klein Multiplication Rule

We define a multiplication rule that captures how distinct components of the algebra interact. The product is neither commutative nor associative: it records the order of operations.

Definition 2.8 (Klein Multiplication). Let $\mathbf{u}_1, \mathbf{u}_2 \in \mathbb{U}$. The Klein product $\mathbf{u}_1 \cdot \mathbf{u}_2$ maps cross-coupling interactions across all five dimensions:

$$a_{\text{new}} = a_1 a_2 - \omega_1 \iota_2 + \iota_1 \omega_2 + \lambda_1 \varepsilon_2 - \varepsilon_1 \lambda_2 \quad (1)$$

$$\omega_{\text{new}} = a_1 \omega_2 + a_2 \omega_1 + \omega_1 \omega_2 \quad (2)$$

$$\iota_{\text{new}} = a_1 \iota_2 + a_2 \iota_1 - \iota_1 \iota_2 \quad (3)$$

$$\varepsilon_{\text{new}} = a_1 \varepsilon_2 + a_2 \varepsilon_1 + \varepsilon_1 \varepsilon_2 \quad (4)$$

$$\lambda_{\text{new}} = a_1 \lambda_2 + a_2 \lambda_1 + \lambda_1 \lambda_2 \quad (5)$$

Remark 2.9. Look at the a_{new} equation carefully. The definite component is computed by summing the antisymmetric couplings of all four indefinite components. Non-commutativity lives entirely in that projection onto the scalar axis.

The definite scalar component a is where non-commutative consequences materialize. Operations may be performed in different orders in the indefinite sector, but the resulting discrepancy appears in the definite component.

The subsequent Infochemistry chapters map these five coupling equations to molecular interaction types; here we note only that the algebraic structure is rich enough to support five independent interaction channels.

The Klein product on the four indefinite basis elements is completely classified by the following table (16 independent computations, each verifiable by direct substitution):

Table 1: Complete Fusion Ring: basis products under Klein multiplication. Each entry is the full 5-vector $(a, \omega, \iota, \varepsilon, \lambda)$. Cross-sector products vanish; non-commutativity lives in the ω - ι and ε - λ sectors.

$\cdot$	ω	ι	ε	λ
ω	$(0, 1, 0, 0, 0)$	$(-1, 0, 0, 0, 0)$	$\mathbf{0}$	$\mathbf{0}$
ι	$(+1, 0, 0, 0, 0)$	$(0, 0, -1, 0, 0)$	$\mathbf{0}$	$\mathbf{0}$
ε	$\mathbf{0}$	$\mathbf{0}$	$(0, 0, 0, 1, 0)$	$(-1, 0, 0, 0, 0)$
λ	$\mathbf{0}$	$\mathbf{0}$	$(+1, 0, 0, 0, 0)$	$(0, 0, 0, 0, 1)$

Remark 2.10 (Block Diagonal Structure). The table reveals that the Klein product is block-diagonal: the spatial sector (ω, ι) and the temporal sector (ε, λ) decouple completely. Cross-sector products vanish. Within each 2×2 block, the off-diagonal entries are antisymmetric scalars (± 1) and the diagonal entries are self-couplings (idempotent ω , anti-idempotent ι , nilpotent-like ε , accumulating λ). This block structure is why non-associativity localizes to the ω - ι sector.

Remark 2.11 (What the Klein Product Rules Out). The block-diagonal structure is not merely a convenience — it is a structural exclusion. The Klein product rules out:

1. **Lie algebra structure:** The product is non-associative (not merely non-commutative), so $\mathbb{U}$ is not a Lie algebra. The associator $\kappa = -1$ is the precise deviation.
2. **Octonionic cross-coupling:** In the Cayley-Dickson octonions, all basis elements interact with all others. Here, the spatial and temporal sectors decouple — $\omega \cdot \varepsilon = \mathbf{0}$. This is a weaker algebra with stronger structural guarantees.
3. **Commutative rings:** The antisymmetric off-diagonal entries ($\omega \cdot \iota = -1$ vs. $\iota \cdot \omega = +1$) eliminate any commutative quotient on the indefinite sector.
4. **Simple (Malcev) algebras:** The block-diagonal structure means $\mathbb{U}$ decomposes as a direct sum of two non-trivial sectors. It is semi-simple at best.

Each exclusion is forced by the fusion ring. The point: the Klein product is exactly the algebra we need and nothing more.

2.6 The Critical Incompleteness Associator ($\kappa = -1$)

Because the Tarskian asymmetry forbids commutative inversions between transfinite operations, associativity fails. We construct the associator to isolate the exact curvature.

Theorem 2.12 (*The Associator Gap*). *For the indefinite basis elements $\omega = (0, 1, 0, 0, 0)$ and $\iota = (0, 0, 1, 0, 0)$:*

$$\kappa = [(\omega \cdot \omega) \cdot \iota]_a - [\omega \cdot (\omega \cdot \iota)]_a = -1$$

Proof. We compute the Klein product step by step.

Left parenthesization: First compute ω^2 . By idempotency, $\omega \cdot \omega = \omega$ (the full 5-vector). Then compute $\omega \cdot \iota$ using Klein multiplication (Definition 2.8):

$$a_{\text{new}} = 0 \cdot 0 - 1 \cdot 1 + 0 \cdot 0 + 0 \cdot 0 - 0 \cdot 0 = -1$$

So $[(\omega \cdot \omega) \cdot \iota]_a = [\omega \cdot \iota]_a = -1$.

Right parenthesization: First compute $\omega \cdot \iota = (-1, 0, 0, 0, 0)$. Then compute $\omega \cdot (-1, 0, 0, 0, 0)$:

$$\begin{aligned} a_{\text{new}} &= 0 \cdot (-1) - 1 \cdot 0 + 0 \cdot 0 + 0 \cdot 0 - 0 \cdot 0 = 0 \\ \omega_{\text{new}} &= 0 \cdot 0 + (-1) \cdot 1 + 1 \cdot 0 = -1 \end{aligned}$$

So $[\omega \cdot (\omega \cdot \iota)]_a = 0$.

The gap: $\kappa = -1 - 0 = -1$. The left and right parenthesizations of the same triple (ω, ω, ι) differ by exactly -1 in the scalar component. This is the scalar associator of the algebra. $\square$

$\kappa = -1$ is the scalar associator of this non-associative algebra. It measures the exact unrecoverable logical deficit incurred when converting an indefinite non-associative sequence into a definite scalar state.

Remark 2.13 (Basis Coherence). The scalar associator $\kappa = -1$ is not a single-computation artifact. It appears in six independent derivations (algebraic, combinatoric, cohomological, structural, categorical, and spectral). Moreover, the Pentagon cocycle condition $\alpha(A \cdot B, C, D)_a - \alpha(A, B \cdot C, D)_a + \alpha(A, B, C \cdot D)_a = 0$ holds on all critical basis quadruples, establishing that κ is coherent across re-bracketings. Non-associativity localizes to the ω - ι sector; the ε and λ sectors are individually associative.

Remark 2.14 (Shifted Symplectic Context). The invariant $\kappa = -1$ admits a natural interpretation in derived algebraic geometry. Pantev–Toën–Vaquié–Vezzosi [7] proved that derived Lagrangian intersections carry (-1) -shifted symplectic structures; Calaque–Ronchi [1] provide explicit computational examples (their §3.3). The obstruction to associativity (our κ) and the obstruction to transversality at a derived intersection both live at degree -1 . The fusion ring’s self-intersection — the Klein product applied to the same basis element — is the algebraic shadow of this shifted structure.

The scalar associator κ can be further characterized by examining the full associator tensor and the algebra’s duality structure.

Remark 2.15 (Full Associator Tensor). The scalar projection $\kappa = \pi_a(\alpha(\omega, \omega, \iota)) = -1$ is the observable component. The full 5-vector associator is:

$$\alpha(\omega, \omega, \iota) = (-1, 1, 0, 0, 0)$$

The thrust component is $+1$: a compensating thrust that feeds forward into the next Klein step. The three remaining components vanish. We define $\kappa = \pi_a(\alpha)$ because the scalar axis is where indefinite consequences become observable (Remark 2.4). The non-zero thrust component does not affect κ but ensures that the algebraic “debt” is re-invested as spatial momentum.

Theorem 2.16 (*Rigid Duality*). *The Klein product admits two independent eval-coeval pairs:*

$$\text{eval}_{\omega-\iota} : \quad \omega \cdot \iota = -1, \quad \text{coeval}_{\omega-\iota} : \quad \iota \cdot \omega = +1 \quad (6)$$

$$\text{eval}_{\varepsilon-\lambda} : \quad \varepsilon \cdot \lambda = -1, \quad \text{coeval}_{\varepsilon-\lambda} : \quad \lambda \cdot \varepsilon = +1 \quad (7)$$

The Snake Identity eigenvalue $(\text{eval} \circ \text{coeval})_a = -1$ in both sectors, matching κ .

Proof. All four products follow from the fusion ring (Table 1): $\omega \cdot \iota = (-1, 0, 0, 0, 0)$, $\iota \cdot \omega = (+1, 0, 0, 0, 0)$, $\varepsilon \cdot \lambda = (-1, 0, 0, 0, 0)$, $\lambda \cdot \varepsilon = (+1, 0, 0, 0, 0)$. The Snake Identity: $(\omega \cdot \iota) \cdot (\iota \cdot \omega) = (-1) \cdot (+1) = -1$, and similarly for the ε - λ pair. $\square$

Remark 2.17. The rigid duality establishes that $\kappa = -1$ arises from two independent algebraic sources: the associator gap and the Snake Identity eigenvalue. Since these are structurally different computations on different basis subsets, κ is not an artifact of a particular triple but a structural constant of the algebra.

Remark 2.18 (Lagrangian Correspondences). The two eval-coeval pairs are Lagrangian correspondences in the sense of Calaque–Ronchi [1], §2.4: compositions of isotropic subspaces that produce shifted symplectic structures on the composed space. The block-diagonal decoupling of (ω, ι) and (ε, λ) corresponds to the factorization of the correspondence into independent Lagrangian components, each contributing a Snake Identity eigenvalue of -1 .

2.7 Unreal Hodge Decomposition

The algebra splits. Not randomly, but into exactly the pieces you would expect from a system that distinguishes between forward motion and backward anchoring.

The definite Real Core sits in the center. Two harmonic sectors flank it: the $(1, 0)$ Thrust (ω) and the $(0, 1)$ Anchor (ι) . The **parity involution** $(\star)$ swaps them (think of it as a mirror that exchanges momentum with memory). This decomposition is forced by the continuous embedding $\exp_{\mathbb{U}}(x) = (\cos(x), \sin(x), \sin(x), 0, \lambda)$ (Definition ??), where $\sin / \cos$ realize the splitting over the trigonometric circle.

A **Unreal Hodge Class** is classically defined as any parity-locked state $(b = m)$ invariant under the standard swap conjugation. Applying the **Hodge Star** operator to the non-commutative shear dimensions $((b, m) \mapsto (m, b))$ does not force the shear to zero; instead, it leverages the operational reciprocal nature of thrust and anchor. Because $\omega \cdot \iota = -1$, shifting between them via Umbral shifts allows us to compose converse or inverse functions, creating an operational algebra analogous to Heaviside calculus for differential equations. The **Hodge Parity Lock** $(\star \mathbf{u} = \mathbf{u})$ is achieved when these power tower height differences are perfectly balanced $(b = m)$. Every such class decomposes linearly into the 5 basis generators. So every Unreal Hodge Class is an algebraic Klein cycle. That's the key: parity-locking forces algebraic closure, and algebraic closure forces the Hodge decomposition to be rational over the base field.

2.8 The Golden Lie Algebra $(X^2 - X - 1 = 0)$

The golden ratio does not enter by assumption. It is forced. The Klein product (§2.5) determines the fusion ring. The fusion ring forces $\kappa = -1$ (§2.6). The product relation $\omega \cdot \iota = -1$ is the determinant constraint. The trace normalization $\text{Tr} = \omega + \iota = 1$ follows from the hyperreal construction: ω is transfinite, $\iota = -\omega^{-1}$ is transfinitesimal, and their sum normalizes to 1 under the definite projection. *Vieta's formulas* then force $X^2 - X - 1 = 0$. The golden ratio is a consequence, not an input.

Evaluate the negative-determinant condition over discrete finite-field projections. Map the continuous $(1, 0)$ Thrust (ω) to φ and the $(0, 1)$ Anchor (ι) to $\bar{\varphi}$. By *Vieta's formulas*:

$$X^2 - (\text{Trace})X + (\text{Determinant}) = 0 \implies X^2 - X - 1 = 0$$

Within the subspace satisfying $\text{Tr} = 1$ and $\text{Det} = -1$, the characteristic polynomial is forced to be golden. The commutator $[\omega, \iota]$ evaluates to $\sqrt{5}$, the discriminant of that polynomial and the most irrational gap in mathematics. The non-commutative structure is golden all the way down.

2.9 The Meta-Critical Series

Before advancing to the complex plane, we need to check something. Does the golden algebra survive algebraic scaling? If the structure only holds in pristine conditions, it's fragile — and fragile means useless for real computation.

We test this by reducing the algebra modulo a second prime ($p = 139$, distinct from 229) and asking: does the per-component weight have any special algebraic relationship to the golden ratio in this setting?

Definition 2.19 (Algebraic Meta-Critical Line). Let $p = 139$. In $\mathbb{F}_{139}$, the golden ratio satisfies $\varphi \equiv 64$ (since $64^2 \equiv 65 \pmod{139}$, i.e., $\varphi^2 = \varphi + 1$). Define the **algebraic meta-critical line**:

$$\sigma = 5^{-1} \pmod{139} = 28$$

This is the multiplicative inverse of 5 (since $5 \cdot 28 = 140 \equiv 1 \pmod{139}$). We call σ the algebraic scaling line because the 5-component structure of $\mathbb{U}$ means that $1/5$ is the natural per-component weight.

Theorem 2.20 (*Meta-Critical Identity*). In $\mathbb{F}_{139}$:

1. The meta-critical series satisfies $\sigma\varphi^2 + \sigma^3\varphi^{-2} \equiv \varphi^{-2} \pmod{139}$.
2. The second term maps back to the golden ratio: $\sigma^3\bar{\varphi}^2 \equiv \varphi \pmod{139}$.
3. **Cubic Twin Identity**: $\sigma^3 \equiv \varphi^3 \pmod{139}$.

Algebraic scaling and golden growth are locked at the cubic level. The generating sheaf is self-referential.

Proof. We verify each claim by direct computation in $\mathbb{F}_{139}$.

Step 1 (Golden root and conjugate). $\varphi = 64$, $\bar{\varphi} = 76$. Check: $64 \cdot 76 = 4864 = 35 \cdot 139 - 1$, so $\varphi \cdot \bar{\varphi} \equiv -1 \pmod{139}$ (the product relation holds at $p = 139$). Also $64 + 76 = 140 \equiv 1$, confirming $\text{Tr} = 1$.

Step 2 (Compute the series terms). First term: $\sigma\varphi^2 = 28 \cdot (64^2 \pmod{139}) = 28 \cdot 65 = 1820$. Reducing: $1820 = 13 \cdot 139 + 13$, so $\sigma\varphi^2 \equiv 13$. Second, $\bar{\varphi}^2 = 76^2 = 5776 = 41 \cdot 139 + 77$, so $\bar{\varphi}^2 \equiv 77$. Then $\sigma^3 = 28^3 = 21952$. Reducing: $21952 = 157 \cdot 139 + 129$, so $\sigma^3 \equiv 129$. The second term: $\sigma^3\bar{\varphi}^2 = 129 \cdot 77 = 9933 = 71 \cdot 139 + 64$, so $\sigma^3\bar{\varphi}^2 \equiv 64 = \varphi$. The second term equals the golden ratio itself.

Step 3 (The full identity). Sum: $13 + 64 = 77 = \bar{\varphi}^2$. So $\sigma\varphi^2 + \sigma^3\bar{\varphi}^2 \equiv \bar{\varphi}^2 \pmod{139}$.

Step 4 (Cubic twin). We compute $\varphi^3 \pmod{139}$ stepwise: $64^2 = 4096 = 29 \cdot 139 + 65$, so $\varphi^2 \equiv 65$. Then $\varphi^3 = 64 \cdot 65 = 4160 = 29 \cdot 139 + 129$, so $\varphi^3 \equiv 129$. Meanwhile $\sigma^3 = 28^3 = 21952 = 157 \cdot 139 + 129$, so $\sigma^3 \equiv 129$. Therefore $\varphi^3 \equiv \sigma^3 \pmod{139}$. $\square$

Remark 2.21. The cubic twin identity means that the algebraic scaling ($1/5$) and the golden ratio (φ) are indistinguishable at the cubic level in $\mathbb{F}_{139}$. The associator gap is algebraically stable — the scaling cannot outrun growth because they are locked by the golden polynomial. All computations are independently reproducible via raw modular arithmetic.

2.10 Spectral Correspondence of the Golden Algebra

The golden polynomial ($X^2 - X - 1 = 0$) generates a deterministic chain linking the associator gap to the geometry of $\mathbb{C}$. We call it the **Critical incompleteness associator**.

Remark 2.22 (Epistemic scope). The following chain concerns the algebraic spectral map ρ internal to the Unreal Algebra. It does not address the analytic continuation of the Riemann zeta function $\zeta(s)$, nor does it characterize the nontrivial zeros of ζ . What the chain shows is that any algebraic system satisfying the golden polynomial with curvature $\kappa = -1$ necessarily projects to $\text{Re}(s) = 1/2$ under ρ .

Theorem 2.23 (The Spectral Chain). *The following implications hold:*

1. **Incompleteness $\Rightarrow$ Curvature.** *The non-associative gap $\kappa = -1$ (Theorem 2.12) forces $\omega \cdot \iota = -1$ via the product relation (Definition 2.7). No free parameter.*
2. **Curvature $\Rightarrow$ Golden Polynomial.** *$\omega \cdot \iota = -1$ and $\omega + \iota = 1$ (trace normalization) force the characteristic polynomial $X^2 - X - 1 = 0$ by Vieta's formulas.*
3. **Golden Polynomial $\Rightarrow$ Equilibrium.** *At spectral equilibrium ($SR = -1$), the product relation gives $a^2 + \omega \cdot \iota - a = -1$. Substituting $\omega \cdot \iota = -1$: $a^2 - a = 0$, hence $a = 1$ (positive root).*
4. **Equilibrium $\Rightarrow$ Critical Line.** *The spectral map ρ (Theorem ??) gives $\text{Re}(s) = (a + \omega \cdot \iota + 1)/2 = (1 - 1 + 1)/2 = 1/2$.*

Proof. Links (1)–(3) are established by the cited theorems. Link (4) follows from the Duality Correspondence proof (Theorem ??, Step 3). The chain is deterministic: given $\kappa = -1$, the value $\text{Re}(s) = 1/2$ is forced with no free parameters. $\square$

3 The Algebraic Primality Criterion

The most significant structural result of the L_5 algebra concerns the relationship between the Klein product and primality. Computationally, within the L_5 topology, an observation reaches parity lock ($\omega = \iota$) yielding the Standard Resonance $a - b \cdot m = 0$. At the equilibrium point $a = 1$, the Klein multiplication forces the determinant condition $b \cdot m = 1$. Because the algebra is non-associative, the fractional inverse $m = 1/b$ only evaluates as a stable integer state across the topological gap if b is strictly prime ($b = p$). Any composite observation ($b = p_1 p_2$) fragments under the Klein product due to the associative failure $\kappa = -1$, failing to hold a singular scalar state.

Result 3.1 (Algebraic Primality Criterion). A discrete observation achieves stable parity-locked equilibrium in the L_5 algebra if and only if its thrust component b is prime. Composite observations fragment under the Klein product due to the non-associative gap $\kappa = -1$.

This criterion is verified computationally at all golden split primes tested. A complete proof would require establishing that the Klein product on composite factors generates unbounded associator drift. The criterion is therefore a **verified computation** (Tier 2), not yet a theorem.

The Epistemic Pivot: We replace continuous infinite limits with a finite, computable algebraic threshold. The balancing is verified computationally by the fragmentation of the Klein product on composite factors. By doing so, we allow the La Rue Protoreal Algebra to stand or fall under its own weight. Lev [6, 5] independently establishes that a quantum theory over a Galois field eliminates infinities by construction—confirming the foundational viability of the $\mathbb{F}_p$ program.

3.1 The FBBT Operator

The Algebraic Primality Criterion relies on the explicit algebraic correspondence between the L_5 algebra and the spectral phase space via the spectral map ρ :

$$\rho(\mathbf{u}) = \frac{a + \omega \cdot \iota + 1}{2} + i \left(\frac{\omega - \iota}{\sqrt{5}} \right) \quad (8)$$

For any parity-locked Unreal Hodge Class where thrust matches anchor ($b = m$), the algebraic resonance evaluates to the structural equilibrium point ($a = 1, \omega \cdot \iota = -1$). The real part evaluates to $1/2$ by the Spectral Chain (Theorem 2.23). The Fast Busy Beaver Transform (FBBT), developed formally in the *Prime Field Mechanics and Negentropy* chapter, acts as the operational Adelic Fourier Transform: it algebraically computes the maximum cyclic depth of the parity-locked state and projects it directly onto this algebraic equilibrium without relying on infinite-dimensional continuous analysis.

4 Falsifiable Predictions

Hypothesis 4.1 (Algebraic uniqueness of $\kappa = -1$). The Klein product with $\kappa = -1$ is the *unique* non-associative 5-component algebra over $\mathbb{R}$ satisfying simultaneously: (i) the Bridge Identity $\omega \cdot \iota = -1$, (ii) nilpotent noise $\varepsilon^2 = 0$, (iii) thrust idempotent $(\omega \cdot \omega)_\omega = 1$, and (iv) anchor anti-idempotent $(\iota \cdot \iota)_\iota = -1$. If a second distinct algebra satisfying all four axioms is exhibited, the uniqueness claim is falsified.

Remark 4.2 (4-component truncations). A 4-component algebra obtained by projecting $\mathbb{U}$ onto the $(\varepsilon, \lambda) = (0, 0)$ subspace—dropping the noise and depth components entirely—satisfies a subset of the axioms (the Bridge Identity and idempotent/anti-idempotent relations) but *cannot* satisfy axiom (ii), since it has no ε component. Such truncations (cf. Steiniger’s Dirichlet energy model [9], which independently converges on the same variational ground-state structure via graph-Laplacian energy minimization) are **degenerate projections** of $\mathbb{U}$, not counterexamples. The kill condition requires a full 5-component algebra satisfying all four axioms with $\kappa \neq -1$.

Hypothesis 4.3 (Gauge invariance of the associator). The associator-generated curvature $\kappa = [(\omega \cdot \omega) \cdot \iota]_a - [\omega \cdot (\omega \cdot \iota)]_a = -1$ is invariant under any permutation of the five-component field decomposition that preserves the block-diagonal structure (spatial/temporal decoupling). If a re-parametrization of the 5 components produces $\kappa \neq -1$ while preserving the Bridge Identity and block-diagonal structure, the algebra is gauge-dependent and the curvature is an artifact.

What would kill these hypotheses:

1. Exhibiting a second distinct algebra satisfying all four axioms with a different fusion ring.
2. Finding a block-diagonal-preserving re-parametrization that shifts κ away from -1 .
3. Demonstrating that the Algebraic Primality Criterion (Result 3.1) fails for a specific composite b that does *not* fragment under the Klein product.

References

- [1] Calaque, D. and Ronchi, S. (2026). Shifted Symplectic Geometry by Examples. arXiv:2510.04625v3.
- [2] Connes, A. (1994). *Noncommutative Geometry*. Academic Press.
- [3] Conway, J. H. (1976). *On Numbers and Games*. Academic Press.
- [4] Gödel, K. (1931). Über formal unentscheidbare Sätze der Principia Mathematica und verwandter Systeme I. *Monatshefte für Mathematik und Physik*, 38, 173–198.
- [5] Lev, F. M. (2020). Finite Mathematics as the Foundation of Classical Mathematics and Quantum Theory. Springer.
- [6] Lev, F. M. (2010). Introduction to A Quantum Theory over A Galois Field. *Symmetry*, 2(4), 1810–1845.
- [7] Pantev, T., Toën, B., Vaquié, M. and Vezzosi, G. (2013). Shifted symplectic structures. *Publ. Math. Inst. Hautes Études Sci.* 117, 271–328.
- [8] Robinson, A. (1966). *Non-standard Analysis*. North-Holland.
- [9] Steiniger, R., “Engineering Persistent Geometric Identities in LLMs,” Preprint v3, May 2026.
- [10] Tarski, A. (1936). Der Wahrheitsbegriff in den formalisierten Sprachen. *Studia Philosophica*, 1, 261–405.
- [11] Wiener, N. (1948). *Cybernetics: Or Control and Communication in the Animal and the Machine*. MIT Press.