

Statistical challenges in astroparticle physics

Pat Scott

Department of Physics, McGill University

Slides available from

<http://www.physics.mcgill.ca/~patscott>

Outline

- 1 Overview
 - New physics
 - Global fits
- 2 Statistical Challenges
 - Detailed astroparticle likelihoods
 - Which experiments can we trust?
 - Compute time, coverage and scanning
 - Systematisation: Parameter space → Theory space

My view of astroparticle physics

How to define astroparticle physics?

My view of astroparticle physics

How to define astroparticle physics?

Option A

Use of known **particle physics** to tell us something about **astronomical objects/processes**

- Cosmic rays
- Supernova interiors, remnants
- Blazar, pulsar/magnetar studies
- Gravitational wave astronomy

My view of astroparticle physics

How to define astroparticle physics?

Option A

Use of known **particle physics** to tell us something about **astronomical objects/processes**

- Cosmic rays
- Supernova interiors, remnants
- Blazar, pulsar/magnetar studies
- Gravitational wave astronomy

Option B

Use of **astronomical probes** to tell us new things about **particle/fundamental physics**

- Dark matter searches
- Cosmological neutrino probes
- Big Bang Nucleosynthesis
- Inflationary theory testing

My view of astroparticle physics

How to define astroparticle physics?

Option A

Use of known **particle physics** to tell us something about **astronomical objects/processes**

- Cosmic rays
- Supernova interiors, remnants
- Blazar, pulsar/magnetar studies
- Gravitational wave astronomy

Option B

Use of **astronomical probes** to tell us new things about **particle/fundamental physics**

- Dark matter searches
- Cosmological neutrino probes
- Big Bang Nucleosynthesis
- Inflationary theory testing

Both are legitimate – **B is what gets me going though**, so is what this talk will focus on

Outline

1 Overview

- New physics
- Global fits

2 Statistical Challenges

- Detailed astroparticle likelihoods
- Which experiments can we trust?
- Compute time, coverage and scanning
- Systematisation: Parameter space → Theory space

Searching for new physics

Many reasons to look for physics Beyond the Standard Model (BSM) of particle physics:

- Higgs mass (hierarchy problem + vacuum stability)
- Dark matter exists
- Baryon asymmetry
- Neutrino masses and mixings

Searching for new physics

Many reasons to look for physics Beyond the Standard Model (BSM) of particle physics:

- Higgs mass (hierarchy problem + vacuum stability)
- Dark matter exists
- Baryon asymmetry
- Neutrino masses and mixings

So what do we do about it?

- Make new particles at high- E colliders
- Study rare processes at high- L colliders
- Hunt for dark matter
- Look for kooky neutrino physics, BBN deviations, baryo/leptogenesis ideas, evidence of inflation

Searching for new physics

Many reasons to look for physics Beyond the Standard Model (BSM) of particle physics:

- Higgs mass (hierarchy problem + vacuum stability)
- Dark matter exists
- Baryon asymmetry
- Neutrino masses and mixings

So what do we do about it?

- Make new particles at high- E colliders
- Study rare processes at high- L colliders
- Hunt for dark matter
- Look for kooky neutrino physics, BBN deviations, baryo/leptogenesis ideas, evidence of inflation

Searching for new physics


Many reasons to look for physics Beyond the Standard Model (BSM) of particle physics:

- Higgs mass (hierarchy problem + vacuum stability)
- Dark matter exists
- Baryon asymmetry
- Neutrino masses and mixings

So what do we do about it?

- Make new particles at high- E colliders
- Study rare processes at high- L colliders
- Hunt for dark matter
- Look for kooky neutrino physics, BBN deviations, baryo/leptogenesis ideas, evidence of inflation

- Direct detection – nuclear collisions and recoils – CDMS, XENON, DAMA, CRESST, CoGeNT, etc




- Direct detection – nuclear collisions and recoils – CDMS, XENON, DAMA, CRESST, CoGeNT, etc
- Direct production – missing E_T or otherwise – LHC, Tevatron




Dark matter searches

- Direct detection – nuclear collisions and recoils – CDMS, XENON, DAMA, CRESST, CoGeNT, etc
- Direct production – missing E_T or otherwise – LHC, Tevatron
- Indirect detection – annihilations producing (Weniger *JCAP*, 1204.2797)
 - gamma-rays – *Fermi*, HESS, CTA
 - anti-protons – PAMELA, AMS
 - anti-deuterons – GAPS
 - neutrinos – IceCube, ANTARES
 - e^+e^- – PAMELA, *Fermi*, ATIC, AMS
→ secondary radiation: Compton⁻¹, synchrotron, bremsstrahlung
 - secondary impacts on the CMB




Dark matter searches

- Direct detection – nuclear collisions and recoils – CDMS, XENON, DAMA, CRESST, CoGeNT, etc
- Direct production – missing E_T or otherwise – LHC, Tevatron
- Indirect detection – annihilations producing (Weniger *JCAP*, 1204.2797)
 - gamma-rays – *Fermi*, HESS, CTA
 - anti-protons – PAMELA, AMS
 - anti-deuterons – GAPS
 - neutrinos – IceCube, ANTARES
 - e^+e^- – PAMELA, *Fermi*, ATIC, AMS
→ secondary radiation: Compton⁻¹, synchrotron, bremsstrahlung
 - secondary impacts on the CMB
- Dark stars – JWST, VLT



Dark matter searches

- Direct detection – nuclear collisions and recoils – CDMS, XENON, DAMA, CRESST, CoGeNT, etc
- Direct production – missing E_T or otherwise – LHC, Tevatron
- Indirect detection – annihilations producing (Weniger *JCAP*, 1204.2797)
 - **gamma-rays** – *Fermi*, *HESS*, *CTA*
 - anti-protons – PAMELA, AMS
 - anti-deuterons – GAPS
 - **neutrinos** – *IceCube*, ANTARES
 - e^+e^- – PAMELA, *Fermi*, ATIC, AMS
→ secondary radiation: Compton⁻¹, synchrotron, bremsstrahlung
 - **secondary impacts on the CMB**
- Dark stars – JWST, VLT



Outline

- 1 Overview
 - New physics
 - Global fits
- 2 Statistical Challenges
 - Detailed astroparticle likelihoods
 - Which experiments can we trust?
 - Compute time, coverage and scanning
 - Systematisation: Parameter space → Theory space

BSM Model Scanning

Goals:

- 1 given a particular theory, determine which parameter combinations fit all experiments, and how well
- 2 given multiple theories, determine which fit the data better, and quantify how much better

BSM Model Scanning

Goals:

- 1 given a particular theory, determine which parameter combinations fit all experiments, and how well
⇒ parameter estimation
- 2 given multiple theories, determine which fit the data better, and quantify how much better ⇒ model comparison

Putting it all together: global fits

Issue 1: Combining fits to different experiments

Relatively easy – composite likelihood ($\mathcal{L}_1 \times \mathcal{L}_2 \equiv \chi_1^2 + \chi_2^2$ for simplest \mathcal{L})

- dark matter relic density from WMAP
- direct detection
- indirect detection
- LHC searches
- precision electroweak tests at LEP
- LEP limits on sparticle masses
- B -factory data (rare decays, $b \rightarrow s\gamma$)
- muon anomalous magnetic moment (“ $g - 2$ ”)

Putting it all together: global fits

Issue 2: Including the effects of uncertainties in input data
Easy – treat them as *nuisance parameters*

Issue 3: Finding the points with the best likelihoods
Tough – MCMCs, nested sampling, genetic algorithms, etc

Issue 4: Comparing theories
Depends – Bayesian model comparison, p values
(TS distribution? \rightarrow coverage???)

Putting it all together: global fits

Issue 2: Including the effects of uncertainties in input data
Easy – treat them as *nuisance parameters*

Issue 3: Finding the points with the best likelihoods
Tough – MCMCs, nested sampling, genetic algorithms, etc

Issue 4: Comparing theories
Depends – Bayesian model comparison, p values
(TS distribution? → coverage???)

The state of play

Yesterday

Experimentalists:

Toy models

Real constraints

Theorists:

Real models

Toy constraints

The state of play

Yesterday

Experimentalists:
Toy models
Real constraints

Theorists:
Real models
Toy constraints



Today

Real models
+
Real constraints
=
Real model
parameter space
constraints

The state of play

Yesterday

Experimentalists:
Toy models
Real constraints

Theorists:
Real models
Toy constraints

Today

Real models
+
Real constraints
=
Real model
parameter space
constraints

Tomorrow

Theory space
constraints

The state of play

Yesterday

Experimentalists:
Toy models
Real constraints

Theorists:
Real models
Toy constraints

Today

Real models
+
Real constraints
=
Real model
parameter space
constraints

Tomorrow

Theory space
constraints

1 Detailed astroparticle likelihoods for arbitrary models

The state of play

Yesterday

Experimentalists:
Toy models
Real constraints

Theorists:
Real models
Toy constraints

Today

Real models
+
Real constraints
=
Real model
parameter space
constraints

Tomorrow

Theory space
constraints

- 1 Detailed astroparticle likelihoods for arbitrary models
- 2 Which experiments can we trust?
- 3 Compute time, coverage and scanning

The state of play

Yesterday

Experimentalists:
Toy models
Real constraints

Theorists:
Real models
Toy constraints

Today

Real models
+
Real constraints
=
Real model
parameter space
constraints

Tomorrow

Theory space
constraints

- 1 Detailed astroparticle likelihoods for arbitrary models
- 2 Which experiments can we trust?
- 3 Compute time, coverage and scanning
- 4 **Systematisation: Parameter space \rightarrow Theory space**

Outline

- 1 Overview
 - New physics
 - Global fits
- 2 Statistical Challenges
 - Detailed astroparticle likelihoods
 - Which experiments can we trust?
 - Compute time, coverage and scanning
 - Systematisation: Parameter space \rightarrow Theory space

Two different approaches to including astro data in BSM scans

- 1 Just use the published limits on $\langle\sigma v\rangle$ (or $\sigma_{\text{SI,SD}}$)
 - Fast – can cover large parameter spaces
 - Not so accurate – experimental limits are invariably based on theoretical assumptions, e.g. $b\bar{b}$ spectrum
 - Full likelihood function almost never available
- 2 Use the data points directly in BSM scans
 - Slow – requires full treatment of instrument profile for each point
 - Accurate – can test each point self-consistently
 - Allows marginalisation over theoretical assumptions
 - Allows construction of full multi-dimensional likelihood function

Two different approaches to including astro data in BSM scans

- 1 Just use the published limits on $\langle\sigma v\rangle$ (or $\sigma_{\text{SI,SD}}$)
 - Fast – can cover large parameter spaces
 - Not so accurate – experimental limits are invariably based on theoretical assumptions, e.g. $b\bar{b}$ spectrum
 - Full likelihood function almost never available
- 2 Use the data points directly in BSM scans
 - Slow – requires full treatment of instrument profile for each point
 - Accurate – can test each point self-consistently
 - Allows marginalisation over theoretical assumptions
 - Allows construction of full multi-dimensional likelihood function

Two different approaches to including astro data in BSM scans

- 1 Just use the published limits on $\langle\sigma v\rangle$ (or $\sigma_{\text{SI,SD}}$)
 - Fast – can cover large parameter spaces
 - Not so accurate – experimental limits are invariably based on theoretical assumptions, e.g. $b\bar{b}$ spectrum
 - Full likelihood function almost never available
- 2 Use the data points directly in BSM scans
 - Slow – requires full treatment of instrument profile for each point
 - Accurate – can test each point self-consistently
 - Allows marginalisation over theoretical assumptions
 - Allows construction of full multi-dimensional likelihood function
- 3 (indirect only: use just flux upper limits)

Gamma-rays


Gamma-ray annihilation searches have been added to the global fits:

Fermi-LAT

Satellite pair conversion telescope

Dwarf galaxy Segue 1

(PS, Conrad et al *JCAP*, 0909.3300)



- Full binned Poissonian likelihood (no χ^2 approximation)
- Full treatment of PSF *and* energy dispersion (with fast convolution library FLATlib)
- Marginalisation over systematic error on effective area
- Diffuse BG from Fermi-LAT Galprop fits
- Isotropic BG best-fit isotropic power law
- J -factor from Martinez et al (*JCAP*, 0902.4715; best at the time)

Gamma-rays


Gamma-ray annihilation searches have been added to the global fits:

HESS

Air Čerenkov telescope

Milky Way+Carina+Sculptor+Sag dwarf

(Ripken, Conrad & PS *JCAP*, 1012.3939)



- χ^2 -based analysis using public flux limits
- ‘Milky Way’ = halo just beyond GC (45–150 pc)
- Virtual internal bremsstrahlung from co-annihilation strip models caught at high- E by HESS
- but: J -factors for Sag dwarf rather uncertain



Advanced IceCube Likelihood

Full unbinned likelihood; number (\mathcal{L}_{num}), spectral ($\mathcal{L}_{\text{spec}}$) & angular (\mathcal{L}_{ang}) parts:

$$\mathcal{L} = \mathcal{L}_{\text{num}}(n|\theta_{\text{signal+BG}}) \prod_{i=1}^n \mathcal{L}_{\text{spec},i} \mathcal{L}_{\text{ang},i} \quad (1)$$

Mock signal reconstruction with IceCube-DeepCore (86-string)

(PS, Savage, Edsjö & IceCube JCAP, 1207.0810)



Advanced IceCube Likelihood

Full unbinned likelihood; number (\mathcal{L}_{num}), spectral ($\mathcal{L}_{\text{spec}}$) & angular (\mathcal{L}_{ang}) parts:



$$\mathcal{L} = \mathcal{L}_{\text{num}}(n|\theta_{\text{signal}+\text{BG}}) \prod_{i=1}^n \mathcal{L}_{\text{spec},i} \mathcal{L}_{\text{ang},i} \quad (1)$$

Limits: IceCube 22-string data

vs.

IceCube-DeepCore (86-string)

(PS, Savage, Edsjö & IceCube JCAP, 1207.0810)




Advanced IceCube Likelihood

Full unbinned likelihood; number (\mathcal{L}_{num}), spectral ($\mathcal{L}_{\text{spec}}$) & angular (\mathcal{L}_{ang}) parts:

$$\mathcal{L} = \mathcal{L}_{\text{num}}(n|\theta_{\text{signal+BG}}) \prod_{i=1}^n \mathcal{L}_{\text{spec},i} \mathcal{L}_{\text{ang},i} \quad (1)$$

Limits: IceCube 22-string data


(PS, Savage, Edsjö & IceCube JCAP, 1207.0810)



vs. IceCube-DeepCore (86-string)

(Silverwood, PS, Danninger et al JCAP, 1210.0844)

\rightarrow IC86 has unique access to pts in more general MSSM



Advanced IceCube Likelihood

Full unbinned likelihood; number (\mathcal{L}_{num}), spectral ($\mathcal{L}_{\text{spec}}$) & angular (\mathcal{L}_{ang}) parts:

$$\mathcal{L} = \mathcal{L}_{\text{num}}(n|\theta_{\text{signal+BG}}) \prod_{i=1}^n \mathcal{L}_{\text{spec},i} \mathcal{L}_{\text{ang},i} \quad (1)$$

The examples here are CMSSM & MSSM-25 – but this about a framework, applicable to any model.

All methods are available implemented in DarkSUSY v5.0.6 and later:
www.darksusy.org

All IceCube data used are available at
<http://icecube.wisc.edu/science/data/ic22-solar-wimp> (and in DarkSUSY, for convenience)

Generalised DM CMB likelihood functions

Simple CMB likelihood function, for

- Any combination of annihilation or decay channels
- Any dark matter mass
- Any decay lifetime/annihilation cross-section

\rightarrow just requires interpolating one number in a table.

Cline & PS, *JCAP*, 1301.5908, using

- CMB energy deposition from Slatyer, 1211.0283 and Finkbeiner et al, 1109.6322
- PYTHIA annihilation/decay spectra of Cirelli et al, 1012.4515.

Generalised DM CMB likelihood functions

Simple CMB likelihood function, for

- Any combination of annihilation or decay channels
- Any dark matter mass
- Any decay lifetime/annihilation cross-section

\rightarrow just requires interpolating one number in a table.

f_{eff} for annihilation:

$$\ln \mathcal{L}(\langle \sigma v \rangle | m_\chi, r_i) = -\frac{1}{2} f_{\text{eff}}^2(m_\chi, r_i) \lambda_1 c_1^2 \left(\frac{\langle \sigma v \rangle}{2 \times 10^{-27} \text{cm}^3 \text{s}^{-1}} \right)^2 \left(\frac{\text{GeV}}{m_\chi} \right)^2 \quad (2)$$

Generalised DM CMB likelihood functions

Simple CMB likelihood function, for

- Any combination of annihilation or decay channels
- Any dark matter mass
- Any decay lifetime/annihilation cross-section

\rightarrow just requires interpolating one number in a table.

f_{eff} for annihilation:

$$\ln \mathcal{L}(\langle \sigma v \rangle | m_\chi, r_i) = -\frac{1}{2} f_{\text{eff}}^2(m_\chi, r_i) \lambda_1 c_1^2 \left(\frac{\langle \sigma v \rangle}{2 \times 10^{-27} \text{cm}^3 \text{s}^{-1}} \right)^2 \left(\frac{\text{GeV}}{m_\chi} \right)^2 \quad (2)$$

η for decay:



$$\ln \mathcal{L}(\tau | m_\chi, r_i) = -\frac{1}{2} \left(\frac{\delta \Omega}{\Omega_{\text{DM}} \tau} \right)^2 \eta^2(\tau, m_\chi, r_i) \quad (3)$$

Generalised DM CMB likelihood functions

Simple CMB likelihood function, for

- Any combination of annihilation or decay channels
- Any dark matter mass
- Any decay lifetime/annihilation cross-section

\rightarrow just requires interpolating one number in a table.



Direct detection likelihoods

Much solid work done in implementing detailed direct detection likelihoods in pheno analyses – details coming up this afternoon



Akrami, Savage et al 1011.4318, 1011.4297

Pato et al 1012.3458, 1106.0743, 1211.7063

Arina et al 1105.5121, 1210.4011

Strege et al 1107.1715, 1112.4192, 1201.3631, 1212.2636

Pato et al 1211.7063



Outline

- 1 Overview
 - New physics
 - Global fits
- 2 **Statistical Challenges**
 - Detailed astroparticle likelihoods
 - **Which experiments can we trust?**
 - Compute time, coverage and scanning
 - Systematisation: Parameter space \rightarrow Theory space

Conflicting experimental results


In many cases we have apparently inconsistent experimental results: anomalies, claimed detections, deviations, etc

Conflicting experimental results

In many cases we have apparently inconsistent experimental results: anomalies, claimed detections, deviations, etc

Direct Detection

DAMA vs. CoGeNT vs. CRESST-II vs. Exclusions




Conflicting experimental results

In many cases we have apparently inconsistent experimental results: anomalies, claimed detections, deviations, etc


Gamma-ray lines

130 GeV line - to bump or not to bump

(Weniger *JCAP*, 1204.2797)



A. Albert for *Fermi*-LAT Collab., LAT Symposium 2012



Conflicting experimental results

In many cases we have apparently inconsistent experimental results: anomalies, claimed detections, deviations, etc

What do we do with the suspect data?

Conflicting experimental results

In many cases we have apparently inconsistent experimental results: anomalies, claimed detections, deviations, etc

What do we do with the suspect data?

- 1 Keep it (remember to hold your nose)

Conflicting experimental results

In many cases we have apparently inconsistent experimental results: anomalies, claimed detections, deviations, etc

What do we do with the suspect data?

- 1 Keep it (remember to hold your nose)
- 2 Toss it out

Conflicting experimental results

In many cases we have apparently inconsistent experimental results: anomalies, claimed detections, deviations, etc

What do we do with the suspect data?

- 1 Keep it (remember to hold your nose)
- 2 Toss it out
- 3 Inflate the systematic errors

Conflicting experimental results

In many cases we have apparently inconsistent experimental results: anomalies, claimed detections, deviations, etc

What do we do with the suspect data?

- 1 Keep it (remember to hold your nose)
- 2 Toss it out
- 3 Inflate the systematic errors
- 4 Give it some arbitrary down-weighting (prior)

Conflicting experimental results

In many cases we have apparently inconsistent experimental results: anomalies, claimed detections, deviations, etc

What do we do with the suspect data?

- 1 Keep it (remember to hold your nose)
- 2 Toss it out
- 3 Inflate the systematic errors
- 4 Give it some arbitrary down-weighting (prior)
- 5 Try to do something a bit more objective (\rightarrow David's talk)

Outline

- 1 Overview
 - New physics
 - Global fits
- 2 **Statistical Challenges**
 - Detailed astroparticle likelihoods
 - Which experiments can we trust?
 - **Compute time, coverage and scanning**
 - Systematisation: Parameter space \rightarrow Theory space

Compute time example: The LHC likelihood monster

Time per point:

$\mathcal{O}(\text{minute})$ in **best** cases

Compute time example: The LHC likelihood monster

Time per point:

$\mathcal{O}(\text{minute})$ in **best** cases

Time per point for global fits to converge:

$\mathcal{O}(\text{seconds})$ in **worst** cases

Compute time example: The LHC likelihood monster

Time per point:

$\mathcal{O}(\text{minute})$ in **best** cases

Time per point for global fits to converge:

$\mathcal{O}(\text{seconds})$ in **worst** cases

Challenge:

About 2 orders of magnitude too slow to actually include LHC data in global fits properly

Taming the LHC monster

First Order Response:

“Test if things depend on the other parameters (hope not),
re-simulate published exclusion curve”

Taming the LHC monster

First Order Response:

“Test if things depend on the other parameters (hope not),
re-simulate published exclusion curve”

Not that great, but OK in some cases

- At least have some sort of likelihood this time
- Still a bit screwed if things do depend a lot on other parameters, but
- allows (potentially shaky) extrapolation, also to non-CMSSM models

Fittino, Mastercode

Taming the LHC monster

Second Order Response:

“That’s ridiculous. I’ve never met a calculation I can’t speed up.
There must be some way to have my cake and eat it too”

Taming the LHC monster

Second Order Response:

“That’s ridiculous. I’ve never met a calculation I can’t speed up.
There must be some way to have my cake and eat it too”

Maybe – this is the challenge.

- Interpolated likelihoods (how to choose nodes?)
- Neural network functional approximation (how to train accurately?)
- Some sort of smart reduction based on event topology?
- Something else?

Coverage

We don't **really** know the distribution of our test statistic in BSM global fits, as it is too expensive to Monte Carlo

- coverage is rarely spot-on unless mapping from parameters to data-space is linear

(Akrami, Savage, PS et al *JCAP*, 1011.4297, Bridges et al *JHEP*, 1011.4306, Strece et al *PRD*, 1201.3631)

- p -value assessments of goodness of fit should be viewed with scepticism (\rightarrow MasterCode)

Coverage

We don't *really* know the distribution of our test statistic in BSM global fits, as it is too expensive to Monte Carlo

- coverage is rarely spot-on unless mapping from parameters to data-space is linear

(Akrami, Savage, PS et al *JCAP*, 1011.4297, Bridges et al *JHEP*, 1011.4306, Strece et al *PRD*, 1201.3631)

- p -value assessments of goodness of fit should be viewed with scepticism (\rightarrow MasterCode)

What do we do for a CL/ p -value when we can't obtain the distribution of our test statistic??

Scanning and Convergence

Convergence remains an issue, especially for profile likelihood

Messy likelihood \implies best-fit point can be (and often is) easily missed (Akrami, PS et al *JHEP*, 0910.3950, Feroz et al *JHEP*, 1101.3296)

- frequentist CLs are often off, as isolikelihood levels are chosen incorrectly
- can impact coverage (overcoverage, or masking of undercoverage due to non- χ^2 TS distribution)
- need to use multiple priors and scanning algorithms (one optimised for profile likelihoods?)

Scanning and Convergence

Convergence remains an issue, especially for profile likelihood



Messy likelihood \implies best-fit point can be (and often is) easily missed
(Akrami, PS et al *JHEP*, 0910.3950, Feroz et al *JHEP*, 1101.3296)

- frequentist CLs are often off, as isolikelihood levels are chosen incorrectly
- can impact coverage (overcoverage, or masking of undercoverage due to non- χ^2 TS distribution)
- need to use multiple priors and scanning algorithms (one optimised for profile likelihoods?)

\rightarrow Differential Evolution
may help

DEIS algorithm
(Differential Evolution with
Importance Sampling)

Roebber, PS, Putze & Holder, in prep



Outline

- 1 Overview
 - New physics
 - Global fits
- 2 Statistical Challenges
 - Detailed astroparticle likelihoods
 - Which experiments can we trust?
 - Compute time, coverage and scanning
 - Systematisation: Parameter space \rightarrow Theory space

CMSSM, SMS \neq BSM

(SMS = Simplified Model Spectrum)

Want to do model comparison to actually work out which theory is right. . .

Challenge:

How do I easily adapt a global fit to different BSM theories?

CMSSM, SMS \neq BSM

(SMS = Simplified Model Spectrum)

Want to do model comparison to actually work out which theory is right. . .

Challenge:

How do I easily adapt a global fit to different BSM theories?

Somehow, we must recast things quickly to a new theory

- data
- likelihood functions
- scanning code 'housekeeping'
- even predictions

\Rightarrow a new, very abstract global fitting framework

Hitting the wall

Issues with current global fit codes:

- Strongly wedded to a few theories (e.g. constrained MSSM / mSUGRA)
- Strongly wedded to a few theory calculators
- All datasets and observables basically hardcoded
- Rough or non-existent treatment of most experiments (astroparticle + collider especially)
- Sub-optimal statistical methods / search algorithms
- \implies *already hitting the wall on theories, data & computational methods*

Closing remarks

- Robust analysis of astroparticle data in the context of new physics requires multi-messenger global fits

Closing remarks

- Robust analysis of astroparticle data in the context of new physics requires multi-messenger global fits
- Detailed astroparticle likelihoods for BSM searches are progressing well

Closing remarks

- Robust analysis of astroparticle data in the context of new physics requires multi-messenger global fits
- Detailed astroparticle likelihoods for BSM searches are progressing well
- Our decision-making process for including/excluding datasets needs improvement

Closing remarks

- Robust analysis of astroparticle data in the context of new physics requires multi-messenger global fits
- Detailed astroparticle likelihoods for BSM searches are progressing well
- Our decision-making process for including/excluding datasets needs improvement
- Coverage, scanning and general compute time all mess with each other – some progress, but more work needed

Closing remarks

- Robust analysis of astroparticle data in the context of new physics requires multi-messenger global fits
- Detailed astroparticle likelihoods for BSM searches are progressing well
- Our decision-making process for including/excluding datasets needs improvement
- Coverage, scanning and general compute time all mess with each other – some progress, but more work needed
- The move to theory space has begun – but requires all the rest to be solved too!