

Lectures in Astroparticle Phenomenology

I. Particle Cosmology

Pat Scott

McGill University / Imperial College London

Slides available from
www.physics.mcgill.ca/~patscott



Lecture Plan

Today: Particle Cosmology

- Λ CDM
- Power spectra of cosmological perturbations
- Reheating, Big Bang nucleosynthesis, cosmic strings

Tomorrow (SlfA Redfern): Dark Matter

- Theories
- Production
- Direct + indirect detection

Thursday (back here again): Global Fits

- Techniques, status and coming developments



Other cool stuff I won't be covering directly

(but that we can chat about at the end of one of lectures if you like)

- Neutrino mass models / GUTs and their observables
- Cosmic ray production + propagation
- Baryogenesis / leptogenesis
- Reionisation



Outline of Lecture 1

- 1 Cosmological Models
 - General
 - Λ CDM
- 2 Power spectra of cosmological perturbations
 - Background
 - Middle Universe observables
 - Rare objects
- 3 Specific particle/field processes (optional)
 - BBN
 - Cosmic strings
 - Phase transitions and reheating



Outline of Lecture 1

- 1 Cosmological Models
 - General
 - Λ CDM
- 2 Power spectra of cosmological perturbations
 - Background
 - Middle Universe observables
 - Rare objects
- 3 Specific particle/field processes (optional)
 - BBN
 - Cosmic strings
 - Phase transitions and reheating



Setting up a framework for cosmology

Step 1.

Assume the Universe to be isotropic and homogeneous

\implies Friedmann-Robertson Walker (FRW) metric:

$$g_{\mu\nu}x^\mu x^\nu = dt^2 + R(t)^2 \left(\frac{dr^2}{1 - kr^2} + r^2 d\Omega^2 \right). \quad (1)$$

- This is our distance measure in spacetime.
- Everything but $R(t)$ is determined by isotropy & homogeneity.



Setting up a framework for cosmology

Step 1.

Assume the Universe to be isotropic and homogeneous

\implies Friedmann-Robertson Walker (FRW) metric:

$$g_{\mu\nu}x^\mu x^\nu = dt^2 + R(t)^2 \left(\frac{dr^2}{1 - kr^2} + r^2 d\Omega^2 \right). \quad (1)$$

- This is our distance measure in spacetime.
- Everything but $R(t)$ is determined by isotropy & homogeneity.

Step 2.

Take the Einstein Field Equations from General Relativity:

$$G_{\mu\nu} = 8\pi G T_{\mu\nu} - \Lambda g_{\mu\nu}. \quad (2)$$



Setting up a framework for cosmology

Step 1.

Assume the Universe to be isotropic and homogeneous

⇒ Friedmann-Robertson Walker (FRW) metric:

$$g_{\mu\nu}x^\mu x^\nu = dt^2 + R(t)^2 \left(\frac{dr^2}{1 - kr^2} + r^2 d\Omega^2 \right). \quad (1)$$

- This is our distance measure in spacetime.
- Everything but $R(t)$ is determined by isotropy & homogeneity.

Step 2.

Take the Einstein Field Equations from General Relativity:

$$\text{geometry} \longrightarrow G_{\mu\nu} = 8\pi G T_{\mu\nu} - \Lambda g_{\mu\nu}. \quad (2)$$



Setting up a framework for cosmology

Step 1.

Assume the Universe to be isotropic and homogeneous
 \implies Friedmann-Robertson Walker (FRW) metric:

$$g_{\mu\nu}x^\mu x^\nu = dt^2 + R(t)^2 \left(\frac{dr^2}{1 - kr^2} + r^2 d\Omega^2 \right). \quad (1)$$

- This is our distance measure in spacetime.
- Everything but $R(t)$ is determined by isotropy & homogeneity.

Step 2.

Take the Einstein Field Equations from General Relativity:

geometry \longrightarrow $G_{\mu\nu} = 8\pi G T_{\mu\nu} - \Lambda g_{\mu\nu}$ energy (2)





Setting up a framework for cosmology

Step 1.

Assume the Universe to be isotropic and homogeneous

⇒ Friedmann-Robertson Walker (FRW) metric:

$$g_{\mu\nu}x^\mu x^\nu = dt^2 + R(t)^2 \left(\frac{dr^2}{1 - kr^2} + r^2 d\Omega^2 \right). \quad (1)$$

- This is our distance measure in spacetime.
- Everything but $R(t)$ is determined by isotropy & homogeneity.

Step 2.

Take the Einstein Field Equations from General Relativity:

$$\text{geometry} \longrightarrow G_{\mu\nu} = 8\pi G T_{\mu\nu} - \Lambda g_{\mu\nu}. \quad \text{energy} \quad (2)$$

→ Tells us how geometry of space adjusts to mass/energy



Setting up a framework for cosmology

Step 3.

Solve $\mu = 0, \nu = 0$ of Einstein Eq. \implies *Friedmann Equation*:

$$H(t)^2 \equiv \left(\frac{\dot{R}(t)}{R(t)} \right)^2 = \frac{8\pi G}{3} \rho(t) - \frac{k}{R(t)^2}. \quad (3)$$



Setting up a framework for cosmology

Step 3.

Solve $\mu = 0, \nu = 0$ of Einstein Eq. \implies *Friedmann Equation*:

$$H(t)^2 \equiv \left(\frac{\dot{R}(t)}{R(t)} \right)^2 = \frac{8\pi G}{3} \rho(t) - \frac{k}{R(t)^2}. \quad (3)$$

Solving this gives Hubble parameter $H(t)$ for some

- relationship between energy density and the scalefactor
 $R(t) = f[\rho(t)]$
- curvature $k \in \{+1, 0, -1\}$



Setting up a framework for cosmology

Step 3.

Solve $\mu = 0, \nu = 0$ of Einstein Eq. \implies *Friedmann Equation:*

$$H(t)^2 \equiv \left(\frac{\dot{R}(t)}{R(t)} \right)^2 = \frac{8\pi G}{3} \rho(t) - \frac{k}{R(t)^2}. \quad (3)$$

Solving this gives Hubble parameter $H(t)$ for some

- relationship between energy density and the scalefactor

$$R(t) = f[\rho(t)]$$

- curvature $k \in \{+1, 0, -1\}$

$H(t)$ encodes the dynamic evolution of the Universe



Setting up a framework for cosmology

Step 3.

Solve $\mu = 0, \nu = 0$ of Einstein Eq. \implies *Friedmann Equation*:

$$H(t)^2 \equiv \left(\frac{\dot{R}(t)}{R(t)} \right)^2 = \frac{8\pi G}{3} \rho(t) - \frac{k}{R(t)^2}. \quad (3)$$

Solving this gives Hubble parameter $H(t)$ for some

- relationship between energy density and the scalefactor

$$R(t) = f[\rho(t)]$$

- curvature $k \in \{+1, 0, -1\}$

$H(t)$ encodes the dynamic evolution of the Universe

Critical density:

For a flat Universe $k = 0$. This defines

$$\text{critical density:} \quad \rho_c = \frac{3H(t)}{8\pi G}$$

$$\text{cosmological density:} \quad \Omega_x \equiv \frac{\rho_x}{\rho_c}$$



Equations of state

Equations of state:

1st law of thermodynamics ($\mu = 0$ in conservation of $T_{\mu\nu}$) is

$$d(\rho R^3) = -pd(R^3), \quad \text{i.e. } \Delta E = -p\Delta V \quad (4)$$

with a constant equation of state $\rho = wp$, we get energy density-scalefactor relations

$$\rho \propto R^{-3(1+w)} \quad (5)$$

For different types of energy:

Matter: $w = 0 \implies \rho \propto R^{-3}$

Radiation: $w = 1/3 \implies \rho \propto R^{-4}$

Vacuum (Λ): $w = -1 \implies \rho \propto \text{constant}$

This is basically enough to solve the Friedmann Equation.



Ingredients of Λ CDM

Ingredients required for a cosmological model

A theory of gravity
+ associated assumptions

Types of energy

their equations of state

their (self-)interactions

An initial spectrum
of perturbations

Choices in Λ CDM

GR

+isotropy, homogeneity

radiation, matter, vacuum/dark energy

$w = 1/3, 0, -1/\text{other}$

photons, baryonic (SM) matter
+cold dark matter (CDM), ??

approximately scale invariant
on large scales



Ingredients of Λ CDM

Ingredients required for a cosmological model

A theory of gravity
+ associated assumptions

Types of energy

their equations of state

their (self-)interactions

An initial spectrum
of perturbations

Choices in Λ CDM

GR

+isotropy, homogeneity

radiation, **matter**, vacuum/dark energy

$w = 1/3, 0, -1$ /other

photons, baryonic (SM) matter
+ **cold dark matter (CDM)**, ??

approximately scale invariant
on large scales



Cosmological probes & 'concordance cosmology'

Joint fit to multiple
cosmological observables
gives a consistent set of
parameter values:


$$\Omega_{\Lambda} \approx 0.73$$

$$\Omega_{\text{matter}} \approx 0.27$$


=

$$\Omega_{\text{CDM}} \approx 0.23 + \Omega_{\text{baryons}} \approx 0.04$$

→ Λ CDM



Kowalski et al *ApJ* 2008



But what about inflation?

Question

Isn't inflation part of the Λ CDM model?



But what about inflation?

Question

Isn't inflation part of the Λ CDM model?

Answer

Not really, no.



But what about inflation?

Question

Isn't inflation part of the Λ CDM model?

Answer

Not really, no.

Approximately scale-invariant spectrum of perturbations to start with, on CMB scales (small wavenumber k)? **Yes.**

Due to inflation by definition? **No.**

$$\mathcal{P}_\delta(k) \propto \mathcal{P}_\mathcal{R}(k) \propto k^{n_s-1} \quad (6)$$



But what about inflation?

Question

Isn't inflation part of the Λ CDM model?

Answer

Not really, no.

Approximately scale-invariant spectrum of perturbations to start with, on CMB scales (small wavenumber k)? **Yes.**

Due to inflation by definition? **No.**

$$\mathcal{P}_\delta(k) \propto \mathcal{P}_\mathcal{R}(k) \propto k^{n_s-1+\alpha \log k/k_0} \quad (6)$$



But what about inflation?

Question

Isn't inflation part of the Λ CDM model?

Answer

Not really, no.

Approximately scale-invariant spectrum of perturbations to start with, on CMB scales (small wavenumber k)? **Yes.**

Due to inflation by definition? **No.**

$$\mathcal{P}_\delta(k) \propto \mathcal{P}_\mathcal{R}(k) \propto k^{n_s-1+\alpha \log k/k_0} \quad (6)$$

Λ CDM does not *demand* inflation, just as it does not *demand* any particular CDM

- inflation is just an idea for getting the required spectrum on CMB scales
- any particular DM model is just an idea for getting CDM



Outline of Lecture 1

- 1 Cosmological Models
 - General
 - Λ CDM
- 2 Power spectra of cosmological perturbations
 - Background
 - Middle Universe observables
 - Rare objects
- 3 Specific particle/field processes (optional)
 - BBN
 - Cosmic strings
 - Phase transitions and reheating



Generation of perturbations

During inflation (or its alternative), quantum fluctuations seed energy/density perturbations

- on all length scales, e.g.

$$\mathcal{P}_\delta(k) \propto k^{n_s-1} \quad (7)$$



Generation of perturbations

During inflation (or its alternative), quantum fluctuations seed energy/density perturbations

- on all length scales, e.g.

$$\mathcal{P}_\delta(k) \propto k^{n_s-1+\alpha \log k/k_0} \quad (7)$$



Generation of perturbations

During inflation (or its alternative), quantum fluctuations seed energy/density perturbations

- on all length scales, e.g.

$$\mathcal{P}_\delta(k) \propto k^{n_s-1+\alpha \log k/k_0} \quad (7)$$

- with some distribution of amplitudes – often assumed to be Gaussian:

$$\text{pdf}(\delta) = \frac{1}{\sqrt{2\pi}\sigma_{\chi,H}(z_X, R)} \exp\left(-\frac{\delta^2}{2\sigma_{\chi,H}^2(z_X, R)^2}\right) \quad (8)$$



Generation of perturbations

During inflation (or its alternative), quantum fluctuations seed energy/density perturbations

- on all length scales, e.g.

$$\mathcal{P}_\delta(k) \propto k^{n_s-1+\alpha \log k/k_0} \quad (7)$$

- with some distribution of amplitudes – often assumed to be Gaussian:

$$\text{pdf}(\delta) = \frac{1}{\sqrt{2\pi}\sigma_{\chi,H}(z_X, R)} \exp\left(-\frac{\delta^2}{2\sigma_{\chi,H}^2(z_X, R)^2}\right) \quad (8)$$

Universe inflates \rightarrow causes perturbations to exit the horizon
(move out of causal contact with each other)



Generation of perturbations

During inflation (or its alternative), quantum fluctuations seed energy/density perturbations

- on all length scales, e.g.

$$\mathcal{P}_\delta(k) \propto k^{n_s-1+\alpha \log k/k_0} \quad (7)$$

- with some distribution of amplitudes – often assumed to be Gaussian:

$$\text{pdf}(\delta) = \frac{1}{\sqrt{2\pi}\sigma_{\chi,H}(z_X, R)} \exp\left(-\frac{\delta^2}{2\sigma_{\chi,H}^2(z_X, R)^2}\right) \quad (8)$$

Universe inflates \rightarrow causes perturbations to exit the horizon
(move out of causal contact with each other)

Universe stops inflating, keeps expanding \rightarrow catches up with the perturbations



Generation of perturbations

During inflation (or its alternative), quantum fluctuations seed energy/density perturbations

- on all length scales, e.g.

$$\mathcal{P}_\delta(k) \propto k^{n_s-1+\alpha \log k/k_0} \quad (7)$$

- with some distribution of amplitudes – often assumed to be Gaussian:

$$\text{pdf}(\delta) = \frac{1}{\sqrt{2\pi}\sigma_{\chi,H}(z_X, R)} \exp\left(-\frac{\delta^2}{2\sigma_{\chi,H}^2(z_X, R)^2}\right) \quad (8)$$

Universe inflates \rightarrow causes perturbations to exit the horizon
(move out of causal contact with each other)

Universe stops inflating, keeps expanding \rightarrow catches up with the perturbations

\Rightarrow first out, last in (bigger-scale perturbations re-enter later)



Growth of perturbations

Post-inflation, Universe quickly becomes **radiation-dominated**

⇒ baryons + photons coupled by electromagnetism

⇒ growth of perturbations damped by radiation
free-streaming

⇒ growth of perturbations is **logarithmic only**

$$\delta \propto \log R \quad (9)$$

At $z \sim 3000$, baryons kinetically decouple as Universe becomes **matter-dominated**

⇒ damping relieved

⇒ perturbations grow linearly, structure growth begins

$$\delta \propto R \quad (10)$$



Observation of perturbations

Essentially all cosmological observables depend on 2 things:

- 1 The initial spectrum of perturbations
= distribution of amplitudes over scales: $\text{pdf}(\delta, k)$, $\mathcal{P}_\delta(k)$
→ this (mostly) comes from your theory of inflation
- 2 How the perturbations + their consequences are processed
→ the geometry of the Universe over time: $H(t)$
→ the specific content of the Universe: $\mathcal{L}_{\text{SM}+\text{BSM}}$
 - new particles
 - exotic objects (e.g. cosmic strings)
 - specific processing events associated with the content (BBN, phase transitions, etc)



⇒ can use CMB, large scale structure, etc to test specific particle theories

(**not** just how much stuff with each w and its impact on $H(t)$)




The cosmic microwave background (CMB)

Key is to look at amount of power on different scales for info on primordial spectrum and processing physics



The CMB – inflation-like examples

Power Spectra allowed by Planck




- ① Λ CDM + $dn_s/d\ln k + d^2(n_s)/d\ln k^2$, *Planck*+WMAP polarisation (WP)
- ② Λ CDM + $r + dn_s/d\ln k$, *Planck*+WP
- ③ Λ CDM + r , *Planck*+WP
- ④ Λ CDM, *Planck*+WP
- ⑤ Λ CDM + $dn_s/d\ln k$, *Planck*+WP
- ⑥ Λ CDM + $r + dn_s/d\ln k$, *Planck*+WP+BAO

Shandera, Erickcek, PS & Yana Galarza *Phys. Rev. D* 2013



Large scale structure (LSS)

- Density perturbations \equiv sound waves \rightarrow matter density oscillations
- Seen in CMB temperature, polarisation anisotropy at $z \sim 1100$
- Eventually grow to form galaxies, etc at $z \lesssim 20$



Some imprint of scales of density oscillations (=primordial spectrum) retained \rightarrow **Baryon Acoustic Oscillations**


SDSS-III Data Release 9 *MNRAS* 2012



Large scale structure – neutrino mass example

Neutrinos are warm dark matter

- ⇒ large free-streaming length, depends on m_ν
 - escape from small collapsing perturbations
- ⇒ Suppression in small-scale matter (processed) power spectrum



SNOWMASS 2013, arXiv:1309.5383



Rare objects: primordial black holes (PBHs)

Question

What is a *primordial* black hole?

- If density perturbations are big enough ($\delta > 0.3$), when they enter the horizon the whole thing collapses
→ immediate black hole
 - PBH mass is horizon mass at re-entry
- ⇒ number of PBHs with M_{PBH} maps directly to amplitude of perturbations on some scale k
- For some distribution of perturbations $\text{pdf}(\delta)$

$$\beta_{\text{PBH}} \equiv \Omega_{\text{PBH}}/\Omega_0 = \int_{\delta_{\min}}^{\infty} \text{pdf}(\delta) d\delta \quad (11)$$

Can repeat at different k to get limits on primordial spectrum
→ limits on inflationary theories



Rare objects: ultracompact minihalos (UCMHs)

Question

What is an *ultracompact* minihalo (UCMH)?



Rare objects: ultracompact minihalos (UCMHs)

Question

What is an *ultracompact* minihalo (UCMH)?

Answer

A DM halo that collapses shortly after matter-radiation equality
 \implies from a large amplitude density perturbation



Rare objects: ultracompact minihalos (UCMHs)

Question

What is an *ultracompact* minihalo (UCMH)?

Answer

A DM halo that collapses shortly after matter-radiation equality
 \Rightarrow from a large amplitude density perturbation

'Shortly' means z_{collapse} is $O(100)$ or more

\Rightarrow isolated collapse

\Rightarrow formation by radial infall

\Rightarrow very steep density profile $\rightarrow \rho \propto r^{-9/4}$

\Rightarrow **excellent indirect detection targets**

PS & Sivertsson
Phys. Rev. Lett. 2009
Lacki & Beacom *ApJL* 2010

Also good lensing prospects Ricotti & Gould *ApJ* 2009; Li et al *Phys. Rev. D* 2012



Rare objects: UCMH formation

Conditions for formation

- Seeded well before matter-radiation equality
- Requires $\delta \gtrsim \mathcal{O}(10^{-3})$
(compare with normal inflationary perturbations $\delta \sim 10^{-5}$)
- \rightarrow much more likely than PBH formation ($\delta \gtrsim 0.3$)

Usefulness


- Like PBHs, UCMH mass set by horizon scale at time of horizon entry
- \Rightarrow specific UCMH mass \equiv specific cosmological scale
- \Rightarrow limit on abundance of specific mass halo \equiv limit on power on specific scale k



Rare objects: observational limits


PBHs: energetic particles from evaporation, lensing, binary disruption

UCMHs: energetic particles from DM annihilation, lensing



Josan et al, *Phys. Rev. D* 2009


Carr et al, *Phys. Rev. D* 2010



Bringmann, PS & Akrami, *Phys. Rev. D* 2012

Rare objects: comparative limits on power spectrum



Limits on \mathcal{P}_δ from UCMHs ~ 5 orders better than from PBHs
 \Rightarrow strong limits on inflationary models



Bringmann, PS & Akrami, *Phys. Rev. D* 2012



Implications for inflation – slow-roll reconstruction



Impacts on slow-roll reconstruction

grey: original dashed: $z_c = 200$ colours: $z_c = 50$
(but beware extrapolation of α from WMAP scales)




Outline of Lecture 1

- 1 Cosmological Models
 - General
 - Λ CDM
- 2 Power spectra of cosmological perturbations
 - Background
 - Middle Universe observables
 - Rare objects
- 3 Specific particle/field processes (optional)
 - BBN
 - Cosmic strings
 - Phase transitions and reheating



Big Bang Nucleosynthesis (BBN)



locco et al (*Phys. Repts.* 2009)


- Light elements (H, He, Li, B, Be) form when Universe cools to $T \sim$ a few MeV
- Relative amounts are sensitive to baryon-to-photon ratio η
 \Rightarrow sensitive to Ω_b
- Can be messed up by additional energy injection from e.g.
 - Late-decaying or annihilating particles
 - Evaporation of PBHs

\Rightarrow stringent bounds on new physics



Cosmic strings

- Nothing (necessarily) to do with string theory
- 1D topological defect caused by field transition to different vacua in causally disconnected regions
- Breaking of *any* $U(1)$ symmetry in the early Universe should produce cosmic strings
- Searches for presence of strings can constrain particle theories up to the GUT scale
- Crucial quantity is string tension
 $G\mu \propto \text{symmetry-breaking scale}^2$
- Observational limits from
 - CMB position-space maps
 - 21 cm maps
 - pulsar timing
 - UCMH searches



Phase transitions and reheating

- As universe cools, vacuum goes through various phase transitions as symmetries break
 - Electroweak phase transition $\mathcal{O}(200 \text{ GeV})$
 - QCD/chiral symmetry breaking $\mathcal{O}(200 \text{ MeV})$
 - Breaking of symmetries associated with new physics
- Phase transitions may or may not produce:
 - defects like cosmic strings (depends on groups involved)
 - strong density perturbations at a particular k (depends on order of transition)
 - impacts on concurrent processes like kinetic decoupling of dark matter (more tomorrow)
- Drastic changes in field content have similar character
 - Reheating: mass in inflaton field at end of inflation converted into other particles, heating Universe
 - Particle genesis (baryo/lepto), creating matter asymmetry



Take-home points

- Cosmological observables are sensitive to
 - Initial distribution of density perturbations
 - Content of the Universe

⇒ they can be used to test

- Theories for inflation
- Theories for new symmetries + particles beyond the Standard Model
- Inflation has few other observables to correlate this with
- ...but many concrete Beyond Standard Model theories lead to correlated signals elsewhere → next 2 lectures

