

Ndvi2Gif: A Python Package for Multi-Seasonal Remote Sensing Analysis via Google Earth Engine

Diego García Díaz ¹

¹ Estación Biológica de Doñana (EBD-CSIC), Spain

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

Editor: [Open Journals](#) 

Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

License

Authors of papers retain copyright
and release the work under a
Creative Commons Attribution 4.0
International License ([CC BY 4.0](#))

Summary

Google Earth Engine (GEE) is a cloud-based platform for planetary-scale geospatial analysis (Gorelick et al., 2017). Ndvi2Gif is a Python package built on top of GEE and geemap (Wu, 2020) that simplifies the generation and analysis of multi-temporal composite images from satellite and climate reanalysis data. The package provides unified access to 7 satellite platforms (Sentinel-1/2/3, Landsat 4–9, MODIS) and climate reanalysis data (ERA5-Land, CHIRPS), with 52 predefined spectral and radar indices plus 48 climate variables. Beyond compositing, the package includes modules for SAR preprocessing, time series and phenology analysis, machine learning land cover classification, and hydroperiod analysis of surface water dynamics.

The core concept of Ndvi2Gif is the creation of **multi-seasonal statistical composites**. The `get_year_composite()` method returns an `ee.ImageCollection` where each image represents one year, with bands for each temporal period (e.g., `periods=12` generates bands named `january`, `february`, ..., `december`). This band-based temporal organization enables queries such as: *What was the maximum flood extent detected in January across 40 years of Landsat (1982–2024)?* by simply calling `collection.select('january').max()`. Built-in zonal statistics extraction via `get_stats()` allows direct computation of statistics per polygon without external GIS processing.

Statement of need

Generating multi-temporal composites in GEE requires 30–40 lines of boilerplate per sensor for collection filtering, cloud masking, scaling to reflectance, and temporal aggregation. Using Ndvi2Gif, this reduces to:

```
from ndvi2gif import NdviSeasonality

ndvi = NdviSeasonality(roi=roi, start_year=2013, end_year=2013,
                       sat='Landsat', index='ndvi', periods=4)
collection = ndvi.get_year_composite()
```

This simplification extends across 7 satellite platforms, 52 spectral and radar indices, 48 climate variables, multiple temporal aggregation methods, and flexible region of interest specifications—all through a unified API that inherits geemap's interactive mapping capabilities. The architectural decisions and design philosophy that enable this simplification are detailed in the sections below. [Figure 1](#) provides an overview of the package's workflow and capabilities.




Figure 1: Ndv2Gif workflow architecture and capabilities. Color coding: Blue — input data sources and ROI specification; Green — processing operations; Purple — main analysis classes; Orange — download-ready output products.

32 **State of the Field**

33 Remote sensing analysis using Google Earth Engine (GEE) has been widely facilitated by several
34 Python-based libraries. Packages such as **geemap** provide interactive mapping and user-friendly
35 access to Earth Engine data, while **eeomont** focuses on preprocessing utilities and spectral
36 index computation while preserving the native ee.ImageCollection temporal structure. **Wxee**
37 emphasizes interoperability with the scientific Python ecosystem by exporting Earth Engine
38 data into xarray objects for sequential time series analysis.

39 While these tools significantly lower the entry barrier to GEE, they are primarily designed
40 as general-purpose interfaces or extensible utility layers. As a result, researchers conducting
41 long-term ecological or phenological studies often need to implement substantial custom code
42 to ensure consistent temporal alignment, multi-seasonal comparability, and reproducibility
43 across decades of observations.

44 Ndvi2Gif addresses this gap by adopting a distinct design philosophy centered on **band-based**
45 **temporal organization**, where each image represents a temporal unit (typically a year) and
46 bands encode fixed intra-annual periods (e.g., months or seasons). This approach enables
47 direct multi-decadal queries across consistent seasonal windows, which is difficult to achieve
48 efficiently using the native collection-based paradigm.

49 Contributing this functionality to existing packages would require fundamental changes to their
50 core temporal abstractions and design goals. Ndvi2Gif therefore complements, rather than
51 replaces, existing GEE Python tools by providing a specialized framework tailored to multi-
52 seasonal ecological analysis, phenology validation, and operational environmental monitoring.

53 Key features

54 Multi-platform data access

55 Ndvi2Gif provides unified access to optical, SAR, and climate reanalysis datasets through a
56 consistent interface (Table 1).

Table 1: Supported satellite and reanalysis datasets.

Platform	Collection	Resolution	Coverage	Variables
Sentinel-2	S2 SR HARMONIZED	10/20m	2015–present	40+ optical
Sentinel-3	OLCI	300m	2016–present	36 optical
Landsat	L4–L9 Col. 2	30m	1982–present	40+ optical
MODIS	Terra/Aqua	500m	2000–present	40+ optical
Sentinel-1	GRD ARD	10m	2014–present	7 SAR
ERA5-Land	Hourly/Monthly	11km	1950–present	47 climate
CHIRPS	Daily	5.5km	1981–present	Precipitation

57 Flexible region of interest

58 The library supports multiple ROI specification formats, enabling seamless integration with
59 existing workflows and the eLTER research network (Wohner et al., 2019).

Table 2: Supported region of interest formats.

Format	Example	Description
Shapefile	'study_area.shp'	Local vector file
GeoJSON	'boundaries.geojson'	GeoJSON file
DEIMS ID	'deimsid:8eda49e9...'	eLTER site
S2 tile	's2:29SQB'	MGRS tile code
Landsat WRS	'wrs:200,32'	Path and Row
ee.Geometry	ee.Geometry.Point()	GEE geometry
geemap	Map.draw_features	Interactive selection

Advanced SAR processing

The S1ARDProcessor module implements a complete Analysis Ready Data (ARD) pipeline for Sentinel-1 imagery, based on Vollrath et al. (2020). This addresses a critical limitation of the default Sentinel-1 products available in Google Earth Engine, which do not include radiometric terrain correction or advanced speckle filtering.

The ARD workflow includes:

- **Radiometric terrain correction** using an angular-based approach with configurable digital elevation models (Copernicus and SRTM), essential for reducing topographic effects in mountainous or heterogeneous terrain.
- **Speckle filtering** using multiple algorithms (Refined Lee, Lee, Gamma MAP, Lee Sigma, Boxcar), allowing users to balance noise reduction and edge preservation depending on the application.
- **Scattering model selection**, supporting both volume scattering (vegetation-dominated surfaces) and surface scattering (bare soil or water).

This preprocessing enables the reliable use of SAR-derived indices for vegetation structure monitoring, surface moisture assessment, and land cover analysis in areas where optical data are limited by cloud cover.

Time series and phenology analysis

The TimeSeriesAnalyzer module extracts, analyses, and visualises temporal dynamics from multi-seasonal composite stacks, operating on temporally aligned bands within each yearly image to ensure consistency across years and reduce sensitivity to irregular acquisition frequency.

The module supports:

- **Trend detection and significance testing**, including non-parametric Mann–Kendall tests, Sen's slope estimation, and linear regression with confidence intervals.
- **Phenological metrics extraction**, such as Start of Season (SOS), End of Season (EOS), Peak of Season (POS), season length, amplitude, and growth and senescence rates, derived from smoothed seasonal trajectories.
- **Seasonal and interannual comparison**, enabling direct assessment of variability and long-term change across fixed temporal windows.
- **Integrated visual diagnostics**, producing multi-panel dashboards that combine time series plots, seasonal distributions, trend summaries, autocorrelation, and data quality indicators.

This design allows researchers to move seamlessly from data extraction to exploratory and quantitative temporal analysis within a single, reproducible workflow, and is particularly suited to long-term ecological monitoring, phenology validation, and climate–vegetation interaction studies.

Machine learning classification

The LandCoverClassifier module provides supervised and unsupervised land cover classification workflows based on multi-temporal composite stacks generated by NdviSeasonality. Supported supervised algorithms include Random Forest, Support Vector Machines (SVM), CART, and Gradient Tree Boost, while unsupervised approaches include K-means and Cascade K-means clustering.

The module supports feature stack generation from multi-seasonal indices, optional normalization, training and validation dataset handling, and accuracy assessment through confusion matrices and standard performance metrics. Feature importance analysis is also available for supervised classifiers, enabling interpretation of which temporal periods or indices contribute most strongly to classification results.

107 Hydroperiod analysis

108 The HydroperiodAnalyzer module computes surface water dynamics and flooding metrics
109 from multi-temporal optical composites. Working directly on NdviSeasonality outputs, it
110 derives pixel-level hydroperiod statistics without requiring external hydrological data.

111 The module computes:

- 112 ■ **Flood days** — total number of days classified as flooded within a period
- 113 ■ **Valid days** — observation count after cloud masking, enabling data quality assessment
- 114 ■ **Normalized flood days** — flood days relative to valid observations, correcting for gaps in
115 coverage
- 116 ■ **First and last day flooded** — timing metrics for flood onset and recession
- 117 ■ **Anomalies** — deviation from the long-term mean hydroperiod, useful for detecting
118 exceptional flood or drought years. The reference mean can be computed either from
119 the years selected in the analysis or from all available cycles in the chosen dataset, giving
120 users control over the baseline period

121 By default, hydrological cycles run from 1 September to 31 August, following the standard
122 hydrological year convention, but the cycle boundaries are fully configurable to accommodate
123 other definitions (e.g., calendar year, wet season onset).

124 This functionality is particularly suited to Mediterranean wetlands and seasonally inundated
125 ecosystems where surface water dynamics are highly variable and cloud cover frequently limits
126 optical observation.

127 Software Design

128 Ndvi2Gif preserves the native `ee.ImageCollection` abstraction but implements a **hierarchical**
129 **band-based approach**: each collection element corresponds to a year, and bands encode fixed
130 intra-annual sub-periods (months, biweekly intervals, or seasons). A 40-year monthly analysis
131 produces 40 images with 12 bands each, rather than 480 individual images.

132 This design enables queries such as: *What is the mean cyanobacteria index (NDCI)*
133 *detected from Sentinel-2 during August across the last 10 years?* by simply calling
134 `collection.select('august').mean()`, or *In which week of the year does chlorophyll*
135 *detected from Sentinel-3 peak?* by computing `collection.max().bandNames()`.

136 This architecture trades increased per-image band dimensionality for drastically simpler,
137 more reproducible seasonal queries. This trade-off is well suited to long-term ecological
138 and phenological analysis, but less appropriate for applications focused on high-frequency
139 pixel-level forecasting.

140 Beyond temporal organization, `ndvi2gif` integrates complete Sentinel-1 ARD preprocessing,
141 automated phenology metrics extraction, machine learning classification, zonal statistics, and
142 `geemap`-based visualization within a unified API.

143 Research Impact Statement

144 Ndvi2Gif has demonstrated realized research impact through peer-reviewed publications,
145 operational research infrastructure, and downstream software integration since its initial
146 release in May 2020. The package has been applied in calibration and validation studies
147 of Earth Observation products for phenology and surface water monitoring at the Doñana
148 protected area (Díaz-Delgado & García-Díaz, 2024).

149 Ndvi2Gif serves as a foundational component of *geeltermap*, a Python mapping application
150 providing operational environmental monitoring tools for the eLTER network (Díaz-Delgado et

151 al., 2024). Geeltermap integrates PhenoApp for phenology monitoring (García-Díaz & Díaz-
152 Delgado, 2023), FloodApp for flood extent analysis, and LSTApp for land surface temperature
153 assessment.

154 The package is currently used as a core analytical tool in the eLTER network infrastructure
155 and the SUMHAL biodiversity monitoring initiative in Spanish protected areas, demonstrating
156 readiness for reproducible multi-temporal remote sensing analysis.

157 AI Usage Disclosure

158 Ndvi2Gif was initially developed in May 2020 and underwent sustained development through
159 2024 before current-generation AI coding assistants became widely available. The core scientific
160 methodology, architectural design decisions, and algorithmic implementations represent the
161 author's original intellectual contributions. AI tools are currently used as development assistants
162 for refactoring, documentation, debugging, and test implementation, while all strategic decisions
163 regarding software architecture and scientific approach remain under the author's direction.

164 Acknowledgements

165 The author thanks the Google Earth Engine team and community, and especially acknowledges
166 Qiusheng Wu for geemap, which serves as the primary foundation of Ndvi2Gif, and for his
167 continued contributions to open-source geospatial software.

168 References

- 169 Díaz-Delgado, R., & García-Díaz, D. (2024). *In-situ validation of land surface phenology,*
170 *land surface temperature and surface water derived from earth observation products:*
171 *Doñana protected area as a potential cal/val supersite.* eLTER Plus Project. <https://doi.org/10.62880/rars25009>
172
- 173 Díaz-Delgado, R., García-Díaz, D., Marangi, C., Silver, M., Oggioni, A., Tagliolato Acquaviva
174 D'Aragona, P., Karnieli, A., Peterseil, J., Vicario, S., Anttila, S., Böttcher, K., Bolton,
175 W., Wohner, C., & Minic, V. (2024). *eLTER PLUS deliverable D4.4 - piloting of eLTER*
176 *specific downstream services contributing to eLTER standard observation variables.* eLTER
177 PLUS Project. <https://doi.org/10.5281/zenodo.14889212>
- 178 García-Díaz, D., & Díaz-Delgado, R. (2023). PhenoApp: A google earth engine based tool for
179 monitoring phenology. *Revista de Teledetección*, 61, 73–81. [https://doi.org/10.4995/raet.](https://doi.org/10.4995/raet.2023.18767)
180 [2023.18767](https://doi.org/10.4995/raet.2023.18767)
- 181 Gorelick, N., Hancher, M., Dixon, M., Ilyushchenko, S., Thau, D., & Moore, R. (2017).
182 Google earth engine: Planetary-scale geospatial analysis for everyone. *Remote Sensing of*
183 *Environment*, 202, 18–27. <https://doi.org/10.1016/j.rse.2017.06.031>
- 184 Vollrath, A., Mullissa, A., & Reiche, J. (2020). Angular-based radiometric slope correction
185 for sentinel-1 on google earth engine. *Remote Sensing*, 12(11), 1867. [https://doi.org/10.](https://doi.org/10.3390/rs12111867)
186 [3390/rs12111867](https://doi.org/10.3390/rs12111867)
- 187 Wohner, C., Peterseil, J., Oggioni, A., Bertrand, N., & Schentz, H. (2019). DEIMS-SDR—a
188 web portal to document research sites and their associated data. *Ecological Informatics*,
189 51, 135–140. <https://doi.org/10.1016/j.ecoinf.2019.01.005>
- 190 Wu, Q. (2020). Geemap: A python package for interactive mapping with google earth engine.
191 *Journal of Open Source Software*, 5(51), 2305. <https://doi.org/10.21105/joss.02305>