

PATIENT'S ID#: \_\_\_\_\_

### Examination, Medical History, and Evaluation

Name: \_\_\_\_\_ date: \_\_\_\_\_ CR# \_\_\_\_\_

Age: \_\_\_\_\_ Sex: \_\_\_\_\_ Weight: \_\_\_\_\_ Height: \_\_\_\_\_ B/P: \_\_\_\_\_ / \_\_\_\_\_ Temp: \_\_\_\_\_ Pulse: \_\_\_\_\_ Respiration: \_\_\_\_\_

Current Medications: RX: \_\_\_\_\_ Prescribed by: \_\_\_\_\_

Non-RX & OTC: \_\_\_\_\_ Tobacco Use: \_\_\_\_\_

Active Medical Conditions: \_\_\_\_\_

Syncope: \_\_\_\_\_ Dizziness: \_\_\_\_\_ Palpitations: \_\_\_\_\_ Chest Pain \_\_\_\_\_ Dyspnea \_\_\_\_\_ Family history of sudden cardiac death \_\_\_\_\_

Hx Medical Conditions: \_\_\_\_\_

Alcoholism:  
Current: \_\_\_\_\_ History: \_\_\_\_\_

Referral (s) \_\_\_\_\_

Previous Substance Abuse TX & Duration: \_\_\_\_\_ History of Suboxone use? ☐yes ☐no History of Methadone use? ☐yes ☐no

Drug of Choice: \_\_\_\_\_ 2nd \_\_\_\_\_ IV use: \_\_\_\_\_

Verification of History of Opioid Dependence: ☐History ☐Other (report) ☐Detox rights reviewed

Significant Psychosocial Issues: \_\_\_\_\_ Education: \_\_\_\_\_ Military: ☐Y ☐N

Employed: Yes \_\_\_\_\_ No \_\_\_\_\_ Where? \_\_\_\_\_ Position? \_\_\_\_\_ How long? \_\_\_\_\_

Psychiatric History:  
Previous  
Hospitalizations \_\_\_\_\_

☐Overdose ☐Delusions: ☐Hallucinations: ☐Violent Behavior ☐Suicide Attempts: -- \_\_\_\_\_

Current Psychiatric Problems: \_\_\_\_\_

| General Appearance: | Normal | Abnormal | Remarks: |
|---------------------|--------|----------|----------|
| Head and Face       |        |          |          |
| Neck                |        |          |          |
| Eyes, Ears          |        |          |          |
| Nose, Throat        |        |          |          |
| Mouth, Teeth        |        |          |          |
| Chest               |        |          |          |
| Lungs               |        |          |          |
| Heart               |        |          |          |
| Abdomen             |        |          |          |
| Hernia, Rectal      |        |          |          |
| Genito-Urinary      |        |          |          |
| Upper Extremity     |        |          |          |
| Lower Extremity     |        |          |          |
| Skin                |        |          |          |
| Back                |        |          |          |

(Signature of Medical Personnel)

Title:

PATIENT'S ID#: \_\_\_\_\_

Patient Name \_\_\_\_\_ Date \_\_\_\_\_

**Have You Ever Had or Currently Have Any of the Following:**

**Remarks**

|                    |                            |                            |                     |                            |                            |  |
|--------------------|----------------------------|----------------------------|---------------------|----------------------------|----------------------------|--|
| Allergies: Drug    | y <input type="checkbox"/> | n <input type="checkbox"/> | Hepatitis           | y <input type="checkbox"/> | n <input type="checkbox"/> |  |
| Environmental      | y <input type="checkbox"/> | n <input type="checkbox"/> | High Blood Pressure | y <input type="checkbox"/> | n <input type="checkbox"/> |  |
| Anemia             | y <input type="checkbox"/> | n <input type="checkbox"/> | Kidney Trouble      | y <input type="checkbox"/> | n <input type="checkbox"/> |  |
| Asthma             | y <input type="checkbox"/> | n <input type="checkbox"/> | Operations, Major   | y <input type="checkbox"/> | n <input type="checkbox"/> |  |
| Arthritis          | y <input type="checkbox"/> | n <input type="checkbox"/> | Operations, Minor   | y <input type="checkbox"/> | n <input type="checkbox"/> |  |
| Bloody Stools      | y <input type="checkbox"/> | n <input type="checkbox"/> | Rheumatism          | y <input type="checkbox"/> | n <input type="checkbox"/> |  |
| Blood in Urine     | y <input type="checkbox"/> | n <input type="checkbox"/> | Rheumatic Fever     | y <input type="checkbox"/> | n <input type="checkbox"/> |  |
| Convulsions        | y <input type="checkbox"/> | n <input type="checkbox"/> | Serious Injury      | y <input type="checkbox"/> | n <input type="checkbox"/> |  |
| Coughing of Blood  | y <input type="checkbox"/> | n <input type="checkbox"/> | Stomach Trouble     | y <input type="checkbox"/> | n <input type="checkbox"/> |  |
| Diabetes           | y <input type="checkbox"/> | n <input type="checkbox"/> | Skin Trouble        | y <input type="checkbox"/> | n <input type="checkbox"/> |  |
| Drug Sensitivities | y <input type="checkbox"/> | n <input type="checkbox"/> | Tuberculosis        | y <input type="checkbox"/> | n <input type="checkbox"/> |  |
| Epilepsy           | y <input type="checkbox"/> | n <input type="checkbox"/> | Venereal Disease    | y <input type="checkbox"/> | n <input type="checkbox"/> |  |
| Hernia             | y <input type="checkbox"/> | n <input type="checkbox"/> | HIV status reviewed | y <input type="checkbox"/> | n <input type="checkbox"/> |  |
| Heart Problems     | y <input type="checkbox"/> | n <input type="checkbox"/> | HIV test offered    | y <input type="checkbox"/> | n <input type="checkbox"/> |  |

| Symptom Evaluation:     | 0 Points | 1 Point | 2 Points |
|-------------------------|----------|---------|----------|
| Yawning                 |          |         |          |
| Dilated Pupils          |          |         |          |
| Restlessness            |          |         |          |
| Lacrimation             |          |         |          |
| Goose Flesh/ Chills     |          |         |          |
| Muscle Cramps           |          |         |          |
| Insomnia                |          |         |          |
| Hyperactive Bowel       |          |         |          |
| Diarrhea                |          |         |          |
| Diaphoresis/Cold Sweats |          |         |          |
| Elevated Pulse          |          |         |          |
| Elevated Blood Pressure |          |         |          |

Pulse: 72 or less = 0 Patient opiate use patterns in the past 24-48 hours:  
100 or more = 2  
Blood Pressure: Less 108/70 = 0  
110-140/70 = 1  
140/70 and above = 2  
Insomnia: 7 hrs in 24 hrs = 0  
1-7 hrs in 24 hrs = 1  
Less than 4 hrs in 24 hrs = 2  
Diarrhea: 0-1 stool in 24 hrs = 0  
2 stools in 12 hrs = 1  
More than 4 stools = 2

WITHDRAWAL: MILD 0-5 pts MODERATE 6-10 pts SEVERE 11-15pts ACUTE above 15 pts

Females: Menstrual Cycle: Regular \_\_\_\_\_ Irregular \_\_\_\_\_ #of live births \_\_\_\_\_ Date of Last Menstruation: \_\_\_\_\_

Are You Pregnant? Y ☐ N ☐ Use of contraceptive? Y ☐ N ☐ Type? Surgery /Other: \_\_\_\_\_

**Physician's Orders:**

**Service(s) Ordered: Medication Assisted Treatment (MAT):** y ☐ n ☐ UA Annually y ☐ n ☐ UDS Monthly (PRN) y ☐ n ☐

Rejected for Treatment: \_\_\_\_\_ Reason for Rejection: \_\_\_\_\_

Patient is prescribed \_\_\_\_\_ mgs of methadone hydrochloride po, qd \_\_\_\_\_ mgs of buprenorphine po, qd as a transfer/new admit/readmit / Continuation Disk ☐ Liquid ☐ Film ☐ Tablet ☐ See case notes for subsequent dose changes/other orders.

The diagnosis is \_\_\_\_\_ per DSM V.

(F11.20)  
OBTAIN: CBC/SMA6/RPR ☐ UA ☐ UDS ☐ PPD ☐ Pregnancy Urine ☐

REFER Patient for: Psych. Eval ☐ Mental Health Counseling ☐ Domestic Violence Counseling ☐ Alcohol Services AA ☐ NA ☐ PCP ☐

OB/GYN ☐ Dental Tx ☐ HIV Testing ☐ Hep C Testing ☐ Education/Employment ☐ Social Services ☐ Other \_\_\_\_\_

**Physician Signature** (Arnold Gaskin MD 90791 U1U6 – DEA BG3505142) \_\_\_\_\_ **Date** \_\_\_\_\_

**APRN Signature** (Kerri Queen, APRN90791U2U6- DEA MQ9345491) \_\_\_\_\_ **Date** \_\_\_\_\_

PATIENT'S ID#: \_\_\_\_\_

## MENTAL HEALTH STATUS FORM

Patient's Name: \_\_\_\_\_ Patient ID#: \_\_\_\_\_ Phase: \_\_\_\_\_ Date: \_\_\_\_\_

**Orientation:** \_\_\_\_\_ x4 ☐ Person ☐ Place ☐ Time ☐ Situation ☐ Alert: ☐ yes ☐ no

**Appearance:** ☐ Appropriate ☐ Unkempt ☐ Unclear ☐ Bizarre \_\_\_\_\_ ☐ Other \_\_\_\_\_

**Motor Behavior:** ☐ Appropriate ☐ Fidgeting ☐ Hyperactive ☐ Pacing ☐ Tics ☐ Unsteady  
☐ Lethargic ☐ Posturing ☐ Psychomotor Retardation ☐ Tremors

**General Behavior:** ☐ Cooperative ☐ Angry ☐ Complaining ☐ Despondent ☐ Mute ☐ Frightened  
☐ Crying ☐ Suspicious ☐ Withdrawn ☐ Flirting ☐ Agitated ☐ Nervous  
☐ Combative ☐ Resistant ☐ Inappropriate ☐ Anxious ☐ Uncooperative  
☐ Domineering ☐ Manipulative ☐ Dependent ☐ Other \_\_\_\_\_

**Mood/Affect:** ☐ Appropriate ☐ Labile ☐ Euphoric ☐ Euthymic ☐ Restricted ☐ Dysphonic (sad, flat)

**Memory:** ☐ No Impairment ☐ Impaired Recent ☐ Impaired Immediate ☐ Impaired Remote

**Speech:** ☐ Normal ☐ Electively Mute ☐ Soft ☐ Slurred ☐ Hesitant ☐ Pressured  
☐ Stuttering ☐ Loud ☐ Hyper Verbal ☐ Other \_\_\_\_\_

**Judgement Insight:** ☐ Good ☐ Fair ☐ Poor ☐ Unable to Assess ☐ Poor/Denial of Problem  
☐ Recognition of Problem ☐ Motivated to Resolve Problem

**Thought Content:** ☐ Alienated ☐ Insightful ☐ Ambivalent ☐ Blaming ☐ Evasive ☐ Distrust  
☐ Denial ☐ Confused ☐ Paranoid ☐ Hopelessness ☐ Irrelevant ☐ Rambling  
☐ Disorganized ☐ Tangential ☐ Hyper Religiosity ☐ Morbid ☐ Illogical  
☐ Defensive ☐ Somatic Preoccupation ☐ Phobic ☐ Obsessive ☐ Self Pity  
☐ Suicidal/Homicidal Ideations ☐ Concrete ☐ Other \_\_\_\_\_

### DSM-5 CHECKLIST OF DIAGNOSTIC CRITERIA; OPIOID USE DISORDER

*Opioid Use Disorder is defined as a problematic pattern of opioid use leading to clinically significant impairment or distress, as manifested by at least 2 of the following, occurring within a 12-month period:*

| Diagnostic Criteria                                                                                                         | Meets Criterion? | Additional/Supporting Information |
|-----------------------------------------------------------------------------------------------------------------------------|------------------|-----------------------------------|
| 1. Opioids are often taken in larger amounts or over a longer period than was intended.                                     |                  |                                   |
| 2. There is a persistent desire to unsuccessful efforts to cut down or control opioid use.                                  |                  |                                   |
| 3. A great deal of time is spent in activities necessary to obtain the opioid, use the opioid, or recover from its effects. |                  |                                   |
| 4. Craving or a strong desire or urge to use opioids.                                                                       |                  |                                   |
| 5. Recurrent opioid use resulting in a failure to fulfill major role obligations at work.                                   |                  |                                   |

|                                                                                                                                                                                      |  |  |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--|--|
| 6. Continued opioid use, despite having persistent or recurrent social or interpersonal problems caused or exacerbated by the effects of opioids.                                    |  |  |
| 7. Important social, occupational, or recreational activities are given up or reduced because of opioid use.                                                                         |  |  |
| 8. Recurrent opioid use in situations in which it is physically hazardous.                                                                                                           |  |  |
| 9. Continued opioid use, despite knowledge of having a persistent or recurrent physical or psychological problem that is likely to have been caused or exacerbated by the substance. |  |  |
| 10. Tolerance.* as defined by <i>either</i> of the following:                                                                                                                        |  |  |
| A. A need for markedly increased amounts of opioids to achieve intoxication or desired effect.                                                                                       |  |  |
| B. A markedly diminished effect with continued use of the same amount of an opioid.                                                                                                  |  |  |
| 11. Withdrawal.* as manifested by <i>either</i> of the following:                                                                                                                    |  |  |
| A. The characteristic opioid withdrawal syndrome.                                                                                                                                    |  |  |
| B. Opioids (or a closely related substance) are taken to relieve or avoid withdrawal symptoms.                                                                                       |  |  |

**\*Note:** This criterion is not considered to be met for those individuals taking opioids solely under appropriate medical supervision.

*Specify if:*

**In early remission:** After full criteria for opioid use disorder were previously met, none of the criteria for opioid use disorder have been met for at least 3 months but for less than 12 months (with the exception that Criterion 4, “Craving, or a strong desire or urge to use opioids,” may be met).

**In sustained remission:** After full criteria for opioid use disorder were previously met, none of the criteria for opioid use disorder have been met at any time during a period of 12 months or longer (with the exception that Criterion 4, “Craving, or a strong desire or urge to use opioids,” may be met).

**On maintenance therapy:** This additional specifier is used if the individual is taking a prescribed agonist medication, such as methadone or buprenorphine, and none of the criteria for opioid use disorder have been met for that class of medication (except tolerance to, or withdrawal from, the agonist). This category also applies to those individuals being maintained on a partial agonist, an agonist/antagonist, or a full antagonist such as oral naltrexone or depot naltrexone.

**Current severity:**

- ☐ **Mild:** Presence of 2–3 symptoms. **Code as: F11.10 (ICD-10)**
- ☐ **Moderate:** Presence of 4–5 symptoms. **Code as: F11.20 (ICD-10)**
- ☐ **Severe:** Presence of 6 or more symptoms. **Code as: F11.20 (ICD-10)**

**Physician Signature** (Arnold Gaskin MD 90791 U1U6 – DEA BG3505142) \_\_\_\_\_ **Date** \_\_\_\_\_

**APRN Signature** (Kerri Queen, APRN90791U2U6- DEA MQ9345491) \_\_\_\_\_ **Date** \_\_\_\_\_