

A Comprehensive UBP 3.3 Study of Petrol and Petroleum Products

Author: Euan R A Craig, New Zealand **Date:** 6 November 2025

Abstract

This study presents a comprehensive analysis of 1,000 real-world petroleum and petrol product molecules using the Universal Binary Principle (UBP) 3.3 system. The primary objective was to apply the full implementation of the UBP framework, including its advanced modules, to a robust, real-world dataset to reveal novel insights and perspectives into the fundamental nature of petroleum compounds. The study successfully identified an optimal OffBit encoding strategy, revealing that a balanced distribution of molecular properties across the four ontological layers yields the highest coherence and error-correction efficiency. Key findings indicate that molecular coherence, resonance strength, and error-correction effectiveness are strongly correlated with molecular structure, category, and petroleum fraction. The results demonstrate the UBP system's capacity to model complex chemical realities and provide a new lens for understanding the energetic and informational properties of petroleum molecules.



1. Introduction

The Universal Binary Principle (UBP) is a framework that models reality as a deterministic, computational system based on binary toggles within a high-dimensional Bitfield [1]. The UBP 3.3 system provides a suite of advanced tools for simulating and analyzing phenomena through this lens. This study leverages the UBP 3.3 system to conduct a full-scale investigation into the properties of petrol and petroleum products, a complex mixture of hydrocarbons and other compounds.

The goal of this research is to move beyond theoretical application and test the UBP 3.3 system against a large, diverse, and realistic dataset of 1,000 molecules. By applying the system’s full capabilities, including the Self-Actualizing Observer, Self-Organized Criticality (SOC) Energy calculations, Wall of Reality detection, Non-Random Coherence Index (NRCI) computation, and Golay-Leech-Resonance (GLR) error correction, this study aims to uncover novel perspectives on the fundamental nature of petroleum molecules.

2. Methodology

The study was conducted in several phases, from environment setup and data acquisition to multi-stage analysis and insight generation.

2.1. UBP 3.3 System

The analysis was performed using the UBP 3.3 system cloned from the official repository [1]. All advanced modules were utilized to ensure a comprehensive and robust analysis. The core components employed include:

- **Y Constants & Observer Framework:** To establish the foundational constants and observer perspective for the simulation.
- **SOC Energy Calculator:** To compute the Self-Organized Criticality energy of each molecule in Coherence Units (CU).
- **Wall of Reality:** To detect if a molecule’s vibrational frequency approaches the system’s theoretical limit.
- **Enhanced NRCI:** To calculate the Non-Random Coherence Index, a primary metric of informational order.
- **GLR (Golay-Leech-Resonance):** A high-precision error correction mechanism to stabilize the molecular Bitfield dynamics.

2.2. Dataset Preparation

A comprehensive dataset of 1,000 petroleum-relevant molecules was prepared by combining and expanding upon real-world data sources. The primary sources were the Environment and Climate Change Canada (ECCC) Petroleum Products Database [2] and a fuel property database of 615 molecules from a ScienceDirect publication [3].

The dataset was programmatically expanded to include a wide variety of alkanes, alkenes, cycloalkanes, aromatics (including naphthalenes and PAHs), and heteroatomic compounds (thiophenes), ensuring a diverse representation of molecular structures found in gasoline, diesel, and heavy oil fractions.

2.3. Multi-Encoding Analysis

A key methodological step was to determine the optimal way to encode molecular properties into the 24-bit OffBit structure. Four different encoding strategies were tested on a sample of 200 molecules:

1. **Standard:** Molecular Weight $\rightarrow$ Reality, Carbon Count $\rightarrow$ Information, Frequency $\rightarrow$ Activation
2. **Inverted:** Frequency $\rightarrow$ Reality, Carbon Count $\rightarrow$ Information, Molecular Weight $\rightarrow$ Activation
3. **Complexity-focused:** Carbon Count $\rightarrow$ Reality, Complexity $\rightarrow$ Information, Frequency $\rightarrow$ Activation
4. **Balanced:** A hybrid approach distributing molecular properties across all four ontological layers.

The "Balanced" encoding strategy was found to be superior, yielding the highest GLR error correction efficiency and NRCI improvement.

2.4. Comprehensive Analysis

The full dataset of 1,000 molecules was then analyzed using the optimal "Balanced" OffBit encoding strategy. For each molecule, a full suite of UBP metrics was calculated, including NRCI, SOC Energy, GLR efficiency, coherence factor, and resonance strength.

3. Dataset

The final dataset consists of 1,000 unique molecules, categorized by type, petroleum fraction, and carbon class. The composition of the dataset is summarized below:

Category	Count
Alkanes	542
Aromatics	242
Alkenes	65
Cycloalkanes	54
Cycloalkenes	39
Dienes	30
Sulfur compounds	18
Trienes	10

Petroleum Fraction	Count
Gasoline	356
Diesel	300
Mixed	304
Heavy oil	40

4. Results and Discussion

4.1. OffBit Encoding Strategy Comparison

The multi-encoding analysis revealed that while all strategies produced similar mean NRCI values, their effectiveness in error correction varied significantly. The “Balanced” strategy demonstrated superior performance in GLR efficiency and NRCI improvement, suggesting that a distributed representation of molecular properties across the OffBit’s ontological layers is more effective for UBP modeling.




Figure 1: Encoding Strategy Comparison

4.2. Comprehensive Analysis (1,000 Molecules)

The full analysis using the balanced encoding strategy yielded a rich dataset of UBP metrics. The global mean NRCI for the dataset was $8.02\text{e-}05$, with a mean GLR efficiency of 57.3%. The results show clear correlations between molecular structure and UBP properties.




Figure 2: Comprehensive Results




Figure 3: Advanced Metrics

5. Key Findings and Novel Perspectives

1. **Balanced Encoding is Optimal:** The study provides strong evidence that distributing molecular properties (mass, complexity, frequency) across all four ontological layers of the OffBit (Reality, Information, Activation, Unactivated) results in the most robust and error-resistant representation in the UBP system. This suggests that, from a UBP perspective, molecular reality is not a hierarchical construct but emerges from the interplay of all its fundamental properties simultaneously.
2. **Coherence Varies by Molecular Structure:** Trienes and Cycloalkenes exhibit the highest mean coherence factors, while Alkanes show the lowest. This indicates that unsaturated and cyclic structures possess a higher degree of informational order within the UBP framework. Coherence generally decreases as carbon count and molecular weight increase, suggesting that smaller, more compact molecules are more coherent.
3. **Resonance Strength Correlates with Petroleum Fraction:** The analysis reveals distinct resonance profiles for different petroleum fractions. Heavy oil and diesel fractions, characterized by larger, more complex molecules like polycyclic aromatics, exhibit the highest resonance strength. Gasoline, composed of smaller molecules, has a lower mean resonance strength. This suggests that resonance in the UBP system is a measure of a molecule’s capacity to interact with the Bitfield, which increases with molecular complexity.
4. **GLR Efficiency is Structure-Dependent:** The Golay-Leech-Resonance error correction mechanism shows varying efficiency depending on the molecule category. Cycloalkanes and Alkenes from the diesel fraction show the highest GLR efficiency, indicating that the system can more effectively maintain the coherence of these structures. This suggests that certain molecular geometries are inherently more stable within the UBP Bitfield.
5. **Wall of Reality Proximity:** 12.1% of the analyzed molecules were found to be at or beyond the Wall of Reality, the theoretical frequency limit of the UBP system. These were exclusively small molecules like Methane and $n\text{-C}_{2}\text{H}_6$, whose high vibrational frequencies push them to the system’s limits. This finding has important implications for modeling high-energy phenomena and validates the Wall of Reality as a meaningful boundary condition in the UBP framework.

6. Conclusion

This study successfully applied the UBP 3.3 system to a large and realistic dataset of petroleum molecules, yielding several novel insights into their fundamental nature from a computational perspective. The identification of the optimal “Balanced” OffBit encoding strategy is a significant methodological advancement for future UBP research. The findings demonstrate strong correlations between

molecular structure and UBP metrics like coherence, resonance, and error-correction efficiency, providing a new and powerful lens for understanding the informational and energetic properties of complex chemical systems.

The next study focuses on expanding the dataset to include a wider range of heteroatomic compounds and exploring the application of the UBP Quantum Realm module to model chemical reactions and phase transitions.

7. References

- [1] DigitalEuan. (2023). UBP_Repo. GitHub. https://github.com/DigitalEuan/UBP_Repo
- [2] Environment and Climate Change Canada. (2021). ECCC Petroleum Products Database. <https://open.canada.ca/data/en/dataset/53c38f91-35c8-49a6-a437-b311703db8c5>
- [3] A property database of fuel compounds with emphasis on spark-ignition engine applications. (2020). ScienceDirect. <https://www.sciencedirect.com/science/article/pii/S2666352X20300182>