Assignment 2

Digjoy Nandi - AI20BTECH11007

Download all python codes from

https://github.com/Digjoy12/Signal-Processing/tree/main/Assignment_2/Code

and latex codes from

https://github.com/Digjoy12/Signal-Processing/blob/main/Assignment 2/main.tex

PROBLEM

(Matrix - Q2.60) Find the matrix X so that

$$\mathbf{X} \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{pmatrix} = \begin{pmatrix} -7 & -8 & -9 \\ 2 & 4 & 6 \end{pmatrix}$$

Solution

Let,

$$\mathbf{A} = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{pmatrix} \tag{0.0.1}$$

$$\mathbf{B} = \begin{pmatrix} -7 & -8 & -9 \\ 2 & 4 & 6 \end{pmatrix} \tag{0.0.2}$$

Now, multiplying A^{T} on both sides,

$$\mathbf{X}\mathbf{A}\mathbf{A}^{\top} = \mathbf{B}\mathbf{A}^{\top} \tag{0.0.3}$$

$$\implies \mathbf{X} = \mathbf{B} \mathbf{A}^{\mathsf{T}} (\mathbf{A} \mathbf{A}^{\mathsf{T}})^{-1} \tag{0.0.4}$$

Therefore,

$$\mathbf{X} = \begin{pmatrix} -7 & -8 & -9 \\ 2 & 4 & 6 \end{pmatrix} \begin{pmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{pmatrix} (\mathbf{A}\mathbf{A}^{\mathsf{T}})^{-1} \tag{0.0.5}$$

$$= \begin{pmatrix} -50 & -122 \\ 28 & 64 \end{pmatrix} \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix} \begin{pmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{bmatrix}^{-1}$$
 (0.0.6)

$$= \begin{pmatrix} -50 & -122 \\ 28 & 64 \end{pmatrix} \begin{bmatrix} 14 & 32 \\ 32 & 77 \end{bmatrix}^{-1}$$
 (0.0.7)

$$= \begin{pmatrix} -50 & -122 \\ 28 & 64 \end{pmatrix} \begin{pmatrix} \frac{77}{54} & \frac{-16}{27} \\ \frac{-16}{27} & \frac{7}{27} \end{pmatrix}$$
 (0.0.8)

$$= \begin{pmatrix} 1 & -2 \\ 2 & 0 \end{pmatrix} \tag{0.0.9}$$