

Discrete Structures (Monsoon 2022)

Ashok Kumar Das

Associate Professor

IEEE Senior Member

Center for Security, Theory and Algorithmic Research
International Institute of Information Technology, Hyderabad (IIIT Hyderabad)

E-mail: *ashok.das@iiit.ac.in*

URL: <http://www.iiit.ac.in/people/faculty/ashokkdas>
<https://sites.google.com/view/iitkgpakdas/>

Recap: Functions

Definition


A function or mapping or map or transformation is defined by two sets X and Y , and a rule (relation) f which assigns to each element of X to exactly one element of Y .

In other words, a (binary) relation f from X to Y is called a function from X to Y , if each element of X is related to exactly one element of Y .

- The set X is called the *domain* and Y the *co-domain (range)* of the function f .
- The image $y \in Y$ (y in Y) of an element $x \in X$ is denoted by $y = f(x)$.
- For a function f from set X to set Y is $f : X \rightarrow Y$, if $y \in Y$, then a pre-image of y is an element $x \in X$ for which $f(x) = y$.
- The set of all elements in Y which have at least one pre-image is called the *image* of f , denoted by $Im(f)$.

Function

- Consider the sets $X = \{a, b, c\}$ and $Y = \{1, 2, 3, 4\}$, and the relation (rule) f from X to Y defined as $f(a) = 2, f(b) = 4, f(c) = 1$.



- The pre-image of 2 is a .
- Note that 3 does not have any pre-image.
- The image of f is $Im(f) = \{1, 2, 4\}$.
- $f(X) = \text{Image of } f = Im(f) = \{f(x) | x \in X\} \subseteq Y$

NOTE: All functions are RELATIONS; however, a relation may or may not be a FUNCTION

Definition (Partial Function)

A **partial function** $f : X \rightarrow Y$ is a rule which assigns to every element $x \in D$ (D is a proper subset of X , that is, $D \subset X$) a unique value in Y .

Types of Functions

Definition (One-to-One Function)

A function $f : X \rightarrow Y$ is **1-1 (one-to-one) or injective** if each element in the co-domain Y is the image of at most one element in the domain X .

In other words, $f : X \rightarrow Y$ is 1-1 if distinct elements in the domain X have distinct images in the co-domain Y , i.e., if $a, b \in X$ such that $a \neq b$, then $f(a) \neq f(b)$ or, equivalently, if $f(a) = f(b)$, then $a = b$.

If a function $f : X \rightarrow Y$ is NOT 1-1, it is called **many-one** function.

Definition (Onto Function)

A function $f : X \rightarrow Y$ is **onto or surjective**, if each element in the co-domain Y is the image of at least one element in the domain X . In other words, $f : X \rightarrow Y$ is called onto if $Im(f) = Y$.

Definition (Bijective Function)

A function $f : X \rightarrow Y$ is **bijective**, if it is both 1-1 and onto.

Theorem

If a function $f : X \rightarrow Y$ is 1-1, then $f : X \rightarrow \text{Im}(f)$ is a bijection.

Theorem

If a function $f : X \rightarrow Y$ is 1-1, and X and Y are finite sets of the same size, that is, $|X| = |Y|$, then $f : X \rightarrow Y$ is a bijection.

- Let $f : X \rightarrow Y$ is a function with $|X| = m$ and $|Y| = n$. Then
 - The total number of functions from X to Y is n^m
 - The total number of injective (1-1) functions from X to Y with $m < n$ is ${}^nC_m \cdot m!$
 - The total number of surjective (onto) functions from X to Y with $m > n$ is

$$n!S(m, n)$$

where the Stirling number is given by

$$S(m, m) = S(m, 1) = 1$$

$$S(m, n) = n \cdot S(m-1, n) + S(m-1, n-1)$$

- The total number of bijective functions from X to Y with $m = n$ is $n!$

Permutations


Definition (Permutation)

Let S be a finite set of elements. A permutation p on S is a bijection from S to itself (i.e., $p : S \rightarrow S$).

Example: Let $S = \{1, 2, 3, 4, 5, 6, 7, 8, 9\}$. A permutation $p : S \rightarrow S$ is defined as follows:

$$p(1) = 3, p(2) = 5, p(3) = 9, p(4) = 4, p(5) = 7, p(6) = 1,$$

$$p(7) = 2, p(8) = 6, p(9) = 8$$



- A permutation $p : S \rightarrow S$ on a finite set $S = \{a_1, a_2, \dots, a_n\}$ is displayed as an array:

$$p = \begin{pmatrix} a_1 & a_2 & \cdots & a_n \\ p(a_1) & p(a_2) & \cdots & p(a_n) \end{pmatrix}$$

where $p(a_i)$ is the p -image of a_i .

Definition (Identity Permutation)

The permutation which maps each element of S onto itself is said to be the *identity permutation* and is denoted by I . Thus, if $S = \{a_1, a_2, \dots, a_n\}$, then

$$I = \begin{pmatrix} a_1 & a_2 & \cdots & a_n \\ a_1 & a_2 & \cdots & a_n \end{pmatrix}$$

- Let $f : S \rightarrow S$ and $g : S \rightarrow S$ be two permutations on S . Since $\text{range}.f = \text{dom}.g$, where $\text{range}.f$ and $\text{dom}.g$ denote the range of f and domain of g respectively, the composition is defined.
- Since f and g are both bijective, $g \circ f : S \rightarrow S$ is also bijective. Therefore, $g \circ f$ is a permutation on S . Similarly, $f \circ g$ is also a permutation on S .
- The products gf and fg are defined by the composite $g \circ f$ and $f \circ g$, respectively.

Multiplication of Permutations

• If

$$f = \begin{pmatrix} a_1 & a_2 & \cdots & a_n \\ f(a_1) & f(a_2) & \cdots & f(a_n) \end{pmatrix}$$

and

$$g = \begin{pmatrix} a_1 & a_2 & \cdots & a_n \\ g(a_1) & g(a_2) & \cdots & g(a_n) \end{pmatrix}$$

then

$$fg = \begin{pmatrix} a_1 & a_2 & \cdots & a_n \\ f[g(a_1)] & f[g(a_2)] & \cdots & f[g(a_n)] \end{pmatrix}$$

and

$$gf = \begin{pmatrix} a_1 & a_2 & \cdots & a_n \\ g[f(a_1)] & g[f(a_2)] & \cdots & g[f(a_n)] \end{pmatrix}$$

Inverse of a permutation

- The inverse of $p : S \rightarrow S$, where $S = \{a_1, a_2, \dots, a_n\}$ is

$$p^{-1} = \begin{pmatrix} p(a_1) & p(a_2) & \cdots & p(a_n) \\ a_1 & a_2 & \cdots & a_n \end{pmatrix}$$

where

$$p = \begin{pmatrix} a_1 & a_2 & \cdots & a_n \\ p(a_1) & p(a_2) & \cdots & p(a_n) \end{pmatrix}$$

- If

$$p = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 3 & 5 & 4 & 2 & 1 \end{pmatrix}$$

then

$$\begin{aligned} p^{-1} &= \begin{pmatrix} 3 & 5 & 4 & 2 & 1 \\ 1 & 2 & 3 & 4 & 5 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 5 & 4 & 1 & 3 & 2 \end{pmatrix} \end{aligned}$$

Definition (Cycle)

Let $S = \{a_1, a_2, \dots, a_n\}$. A permutation $f : S \rightarrow S$ is said to be a cycle of length r or an r -cycle, if there are r elements $a_{i_1}, a_{i_2}, \dots, a_{i_r}$ in S such that

$f(a_{i_1}) = a_{i_2}, f(a_{i_2}) = a_{i_3}, \dots, f(a_{i_{r-1}}) = a_{i_r}, f(a_{i_r}) = a_{i_1}$, and $f(a_j) = a_j$, $j \neq i_1, i_2, \dots, i_r$.

The cycle is denoted by $(a_{i_1} a_{i_2} \dots a_{i_r})$ or by $(a_{i_2} a_{i_3} \dots a_{i_r} a_{i_1})$ or any other form provided the elements appear in a fixed cyclic order.

Index Laws

- $f^m . f^n = f^{m+n}$
- $(f^m)^n = f^{mn}$

hold for integral values of m and n .

- By the law $f^m . g^m = (fg)^m$ does not hold, since $fg \neq gf$, in general.
- The identity permutation

$$I = \begin{pmatrix} a_1 & a_2 & \cdots & a_n \\ a_1 & a_2 & \cdots & a_n \end{pmatrix}$$

on a set $S = \{a_1, a_2, \dots, a_n\}$, is the product of n cycles (a_1) , (a_2) , \dots , (a_n) , each of length 1.

Definition (Transposition)

A 2-cycle is called a transposition.

Definition (Even Permutation)

If a permutation contains even number of transpositions, it is called an even permutation.

Definition (Odd Permutation)

If a permutation contains odd number of transpositions, it is called an odd permutation.

Problem. Examine whether the permutation

$$p = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 2 & 4 & 5 & 6 & 3 & 1 \end{pmatrix}$$

is odd or even.

Solution. Given p can be written as

$$p = \begin{pmatrix} 1 & 2 & 4 & 6 & 3 & 5 \\ 2 & 4 & 6 & 1 & 5 & 3 \end{pmatrix}$$

$$= (1\ 2\ 4\ 6) (3\ 5)$$

$$= (1\ 6) (1\ 4) (1\ 2) (3\ 5)$$

Since p has four transpositions, that is, even number of transpositions, therefore it is EVEN.

Problem. Prove that $(1\ 2\ 3\ \dots\ n) \circ (1\ i) = (1\ i+1\ i+2\ \dots\ n) \circ (2\ 3\ \dots\ i-1\ i)$.

Solution.

$$\begin{aligned}
 \text{LHS} &= (1\ 2\ 3\ \dots\ n) \circ (1\ i) \\
 &= \begin{pmatrix} 1 & 2 & 3 & \dots & i-1 & i & i+1 & \dots & n \\ 2 & 3 & 4 & \dots & i & i+1 & i+2 & \dots & 1 \end{pmatrix} \\
 &= \begin{pmatrix} 1 & 2 & 3 & \dots & i-1 & i & i+1 & \dots & n \\ i & 2 & 3 & \dots & i-1 & 1 & i+1 & \dots & n \end{pmatrix} \\
 &= \begin{pmatrix} 1 & 2 & 3 & \dots & i-1 & i & i+1 & \dots & n \\ i+1 & 3 & 4 & \dots & i & 2 & i+2 & \dots & 1 \end{pmatrix} \\
 &= \begin{pmatrix} 1 & i+1 & \dots & n & 2 & 3 & 4 & \dots & i-1 & i \\ i+1 & i+2 & \dots & 1 & 3 & 4 & 5 & \dots & i & 2 \end{pmatrix} \\
 &= (1\ i+1\ i+2\ \dots\ n) \circ (2\ 3\ 4\ \dots\ i-1\ i) \\
 &= \text{RHS.}
 \end{aligned}$$