

Program Verification: Reversing an Array

HE Mingxin, Max

CS104: `program07 @ yeah.net`

CS108: `mxhe1 @ yeah.net`

I2ML(H) Spring 2023 (CS104 | CS108)

Outline

Program Verification: Reversing an Array

Key Points

Introducing the array assignment rule

Summary

Key Points

Key points to learn:

Partial correctness for array assignments

- ▶ Prove that a Hoare triple is satisfied under partial correctness for a program containing array assignment statements.

The array assignment inference rule

Let A be an array of n integers.

First, write down the sequence of changes.

Resolve all of the changes when we prove the implied's.

$\langle Q[A\{e1 \leftarrow e2\}/A] \rangle$

$A[e1] = e2;$

$\langle Q \rangle$ array assignment

- ▶ A is the original array.
- ▶ $A\{e1 \leftarrow e2\}$ is the new array, which is identical to array A except that the $e1^{th}$ element is $e2$.

The array re-assignment notation

The array reassignment notation:

$$A\{e1 \leftarrow e2\}[i] = \begin{cases} e2, & \text{if } i = e1 \\ A[i], & \text{if } i \neq e1 \end{cases}$$

Note that $e1$ is an index whereas $e2$ is an array element.

We apply assignments from left to right.

Examples:

- ▶ $A\{1 \leftarrow 3\}[1] = 3$
- ▶ $A\{1 \leftarrow 3\}\{1 \leftarrow 4\}[1] = 4$

Reversing an array

Consider an array R of n integers, $R[1], R[2], \dots, R[n]$.

We want to reverse the order of its elements.

Our algorithm:

For each $1 \leq j \leq \lfloor n/2 \rfloor$,
we will swap $R[j]$ with $R[n + 1 - j]$.

Reversing an array

R is an array of n integers, $R[1], R[2], \dots, R[n]$. Prove that the following triple is satisfied under partial correctness.

$$\langle\langle \forall x ((1 \leq x \leq n) \rightarrow (R[x] = r_x)) \rangle\rangle$$
$$j = 1;$$
$$\mathbf{while} \ (2 * j \leq n) \ \{$$
$$t = R[j];$$
$$R[j] = R[n+1-j];$$
$$R[n+1-j] = t;$$
$$j = j + 1;$$
$$\}$$
$$\langle\langle \forall x ((1 \leq x \leq n) \rightarrow (R[x] = r_{n+1-x})) \rangle\rangle$$

Reversing an array

R is an array of n integers, $R[1], R[2], \dots, R[n]$. Prove that the following triple is satisfied under partial correctness.

Let $Inv(j)$ denote our invariant.

$$\begin{aligned} & \langle (\forall x ((1 \leq x \leq n) \rightarrow (R[x] = r_x))) \rangle \\ & j = 1; \\ & \mathbf{while} \ (2 * j \leq n) \ \{ \\ & \quad t = R[j]; \\ & \quad R[j] = R[n+1-j]; \\ & \quad R[n+1-j] = t; \\ & \quad j = j + 1; \\ & \} \\ & \langle (\forall x ((1 \leq x \leq n) \rightarrow (R[x] = r_{n+1-x}))) \rangle \end{aligned}$$

CQ 1 Reversing an array

CQ 1: Consider **the premise of implied (A)**.

Which of the following is an accurate description of the formula?

- (A) No swap has occurred.
- (B) Elements in $[1, j - 1]$ have been swapped, and elements in $[j, (n + 1)/2]$ have NOT been swapped.
- (C) Elements in $[1, j]$ have been swapped, and elements in $[j + 1, (n + 1)/2]$ have NOT been swapped.
- (D) All swaps have been completed.
- (E) None of the above

CQ 2 Reversing an array

CQ 2: Consider **the conclusion of implied (A)**.

Which of the following is an accurate description of the formula?

- (A) No swap has occurred.
- (B) Elements in $[1, j - 1]$ have been swapped, and elements in $[j, (n + 1)/2]$ have NOT been swapped.
- (C) Elements in $[1, j]$ have been swapped, and elements in $[j + 1, (n + 1)/2]$ have NOT been swapped.
- (D) All swaps have been completed.
- (E) None of the above

CQ 3 Reversing an array

CQ 3: Consider **the premise of implied (C)**.

Which of the following is an accurate description of the formula?

- (A) No swap has occurred.
- (B) Elements in $[1, j - 1]$ have been swapped, and elements in $[j, (n + 1)/2]$ have NOT been swapped.
- (C) Elements in $[1, j]$ have been swapped, and elements in $[j + 1, (n + 1)/2]$ have NOT been swapped.
- (D) All swaps have been completed.
- (E) None of the above

CQ 4 Reversing an array

CQ 4: Consider **the conclusion of implied (C)**.

Which of the following is an accurate description of the formula?

- (A) No swap has occurred.
- (B) Elements in $[1, j - 1]$ have been swapped, and elements in $[j, (n + 1)/2]$ have NOT been swapped.
- (C) Elements in $[1, j]$ have been swapped, and elements in $[j + 1, (n + 1)/2]$ have NOT been swapped.
- (D) All swaps have been completed.
- (E) None of the above

CQ 5 Reversing an array

CQ 5: Consider **the premise of implied (B)**.

Which of the following is an accurate description of the formula?

- (A) No swap has occurred.
- (B) Elements in $[1, j - 1]$ have been swapped, and elements in $[j, (n + 1)/2]$ have NOT been swapped.
- (C) Elements in $[1, j]$ have been swapped, and elements in $[j + 1, (n + 1)/2]$ have NOT been swapped.
- (D) All swaps have been completed.
- (E) None of the above

CQ 6 Reversing an array

CQ 6: Consider **the conclusion of implied (B)**.

Which of the following is an accurate description of the formula?

- (A) No swap has occurred.
- (B) Elements in $[1, j - 1]$ have been swapped, and elements in $[j, (n + 1)/2]$ have NOT been swapped.
- (C) Elements in $[1, j]$ have been swapped, and elements in $[j + 1, (n + 1)/2]$ have NOT been swapped.
- (D) All swaps have been completed.
- (E) None of the above

Summary

Key points learnt:

Partial correctness for array assignments

- ▶ Prove that a Hoare triple is satisfied under partial correctness for a program containing array assignment statements.