

Proof System for \mathcal{L}_0

Recall

Two aspects of a formal language.

- ▶ Syntax
 - ▶ formulas
 - ▶ connectives
- ▶ Semantics
 - ▶ truth value/truth assignment
 - ▶ truth table/truth function

A proof system for \mathcal{L}_0

Suppose that φ_1 , φ_2 and φ_3 are \mathcal{L}_0 -formulas. Then each of the following \mathcal{L}_0 -formulas is a **logical axiom**:

(Group I axioms)

- ▶ $(\varphi_1 \rightarrow (\varphi_2 \rightarrow \varphi_3)) \rightarrow ((\varphi_1 \rightarrow \varphi_2) \rightarrow (\varphi_1 \rightarrow \varphi_3))$
- ▶ $\varphi_1 \rightarrow \varphi_1$
- ▶ $\varphi_1 \rightarrow (\varphi_2 \rightarrow \varphi_1)$

(Group II axioms)

- ▶ $\varphi_1 \rightarrow (\neg\varphi_1 \rightarrow \varphi_2)$

(Group III axioms)

▶ $(\neg\varphi_1 \rightarrow \varphi_1) \rightarrow \varphi_1$

(Group IV axioms)

▶ $\neg\varphi_1 \rightarrow (\varphi_1 \rightarrow \varphi_2)$

▶ $\varphi_1 \rightarrow (\neg\varphi_2 \rightarrow \neg(\varphi_1 \rightarrow \varphi_2))$

Δ_0 denote the set of these four groups of logical axioms.

(Group III axioms)

▶ $(\neg\varphi_1 \rightarrow \varphi_1) \rightarrow \varphi_1$

(Group IV axioms)

▶ $\neg\varphi_1 \rightarrow (\varphi_1 \rightarrow \varphi_2)$

▶ $\varphi_1 \rightarrow (\neg\varphi_2 \rightarrow \neg(\varphi_1 \rightarrow \varphi_2))$

Δ_0 denote the set of these four groups of logical axioms.

Proposition 1

Every logical axioms above is a tautology.

Γ -proof

Suppose that $\Gamma \subseteq \mathcal{L}_0$.

Definition 2

Suppose that $s = \langle \varphi_i : i \leq n \rangle$ is a finite sequence of propositional formulas. s is a Γ -proof if for each $i \leq n$ at least one of the following happens:

- ▶ $\varphi_i \in \Gamma$;
- ▶ φ_i is a logical axiom;
- ▶ there exists $j_1, j_2 < i$ such that $\varphi_{j_2} = \varphi_{j_1} \rightarrow \varphi_i$. This rule is called **Modus Ponens**.

Γ -proof

Suppose that $\Gamma \subseteq \mathcal{L}_0$.

Definition 2

Suppose that $s = \langle \varphi_i : i \leq n \rangle$ is a finite sequence of propositional formulas. s is a Γ -proof if for each $i \leq n$ at least one of the following happens:

- ▶ $\varphi_i \in \Gamma$;
- ▶ φ_i is a logical axiom;
- ▶ there exists $j_1, j_2 < i$ such that $\varphi_{j_2} = \varphi_{j_1} \rightarrow \varphi_i$. This rule is called **Modus Ponens**.

Γ -proof

Suppose that $\Gamma \subseteq \mathcal{L}_0$.

Definition 2

Suppose that $s = \langle \varphi_i : i \leq n \rangle$ is a finite sequence of propositional formulas. s is a Γ -proof if for each $i \leq n$ at least one of the following happens:

- ▶ $\varphi_i \in \Gamma$;
- ▶ φ_i is a logical axiom;
- ▶ there exists $j_1, j_2 < i$ such that $\varphi_{j_2} = \varphi_{j_1} \rightarrow \varphi_i$. This rule is called **Modus Ponens**.

Definition 3

$\Gamma \vdash \varphi$ (Γ proves φ) iff there exists a finite sequence $s = \langle \varphi_i : i \leq n \rangle$ such that s is a Γ -proof and such that $\varphi_n = \varphi$.

Such sequence s is called a proof from Γ to φ , and φ is called a consequence of Γ .

When $\Gamma = \emptyset$, write $\vdash \varphi$.

Some properties of Γ -proofs

1. If s is a Γ -proof, and t is an initial segment of s , then t is also a Γ -proof.
2. If $s = \langle \varphi_i : i \leq n \rangle$ and $t = \langle \psi_i : i \leq m \rangle$ are two Γ -proofs, then so is

$$s + t = \langle \varphi_1, \dots, \varphi_n, \psi_1, \dots, \psi_m \rangle.$$

Definition 4

Suppose that $\Gamma \subseteq \mathcal{L}_0$.

1. Γ is **inconsistent** if for some formula φ , $\Gamma \vdash \varphi$ and $\Gamma \vdash \neg\varphi$.
2. Γ is **consistent** if Γ is not inconsistent.
3. Γ is **maximally consistent** if and only if for each formula ψ if $\Gamma \cup \{\psi\}$ is consistent then $\psi \in \Gamma$.

Definition 4

Suppose that $\Gamma \subseteq \mathcal{L}_0$.

1. Γ is **inconsistent** if for some formula φ , $\Gamma \vdash \varphi$ and $\Gamma \vdash \neg\varphi$.
2. Γ is **consistent** if Γ is not inconsistent.
3. Γ is **maximally consistent** if and only if for each formula ψ if $\Gamma \cup \{\psi\}$ is consistent then $\psi \in \Gamma$.

Definition 4

Suppose that $\Gamma \subseteq \mathcal{L}_0$.

1. Γ is **inconsistent** if for some formula φ , $\Gamma \vdash \varphi$ and $\Gamma \vdash \neg\varphi$.
2. Γ is **consistent** if Γ is not inconsistent.
3. Γ is **maximally consistent** if and only if for each formula ψ if $\Gamma \cup \{\psi\}$ is consistent then $\psi \in \Gamma$.

Compare with Γ being **satisfiable**. Our ultimate goal is to show that

Γ is consistent if and only if Γ is satisfiable.

We first show the ‘if’ direction.

Soundness

Theorem 5 (Soundness, version I)

If $\Gamma \subseteq \mathcal{L}_0$ is satisfiable. Then Γ is consistent.

Definition 6 (Logical implication)

Suppose $\Gamma \subseteq \mathcal{L}_0$ and $\varphi \in \mathcal{L}_0$. Then Γ **logically implies** φ , write $\Gamma \models \varphi$, if and only if for every truth assignment ν , $\nu \models \Gamma$ implies $\nu \models \varphi$.

Theorem 7 (Soundness, version II)

Suppose that $\Gamma \subseteq \mathcal{L}_0$, $\varphi \in \mathcal{L}_0$ and that $\Gamma \vdash \varphi$. Then $\Gamma \models \varphi$.

Theorem 7 (Soundness, version II)

Suppose that $\Gamma \subseteq \mathcal{L}_0$, $\varphi \in \mathcal{L}_0$ and that $\Gamma \vdash \varphi$. Then $\Gamma \models \varphi$.

Proof.

By induction on the length of Γ -proofs.



Some lemmas of this proof system

Lemma 8 (Inference)

Suppose that $\Gamma \subseteq \mathcal{L}_0$, $\varphi, \psi \in \mathcal{L}_0$. Suppose that $\Gamma \vdash \psi$ and $\Gamma \vdash (\psi \rightarrow \varphi)$. Then $\Gamma \vdash \varphi$.¹

Lemma 9 (Deduction)

Suppose that $\Gamma \subseteq \mathcal{L}_0$, $\varphi, \psi \in \mathcal{L}_0$ and $\Gamma \cup \{\varphi\} \vdash \psi$. Then $\Gamma \vdash (\varphi \rightarrow \psi)$.²

¹No logical axioms required.

²Group I axioms are needed.

Lemma 10

Suppose that $\Gamma \subseteq \mathcal{L}_0$ and that Γ is inconsistent. Suppose that $\psi \in \mathcal{L}_0$. Then $\Gamma \vdash \psi$.³

Lemma 11

Suppose that $\Gamma \subseteq \mathcal{L}_0$ and that Γ is consistent. Suppose that $\varphi \in \mathcal{L}_0$. Then at least one of $\Gamma \cup \{\varphi\}$ or $\Gamma \cup \{\neg\varphi\}$ is consistent, possibly both.⁴

³Uses the Deduction lemma and Group II axioms.

⁴Needs Group III axioms.

Corollary 12

Suppose that $\Gamma \subseteq \mathcal{L}_0$ and that Γ is maximally consistent. Suppose that $\varphi \in \mathcal{L}_0$. Then

- 1. Either $\varphi \in \Gamma$ or $(\neg\varphi) \in \Gamma$.*
- 2. If $\Gamma \vdash \varphi$ then $\varphi \in \Gamma$.*

Lemma 13

Suppose that $\Gamma \subseteq \mathcal{L}_0$ and that Γ is maximally consistent. Suppose that $\varphi_1, \varphi_2 \in \mathcal{L}_0$. Then $(\varphi_1 \rightarrow \varphi_2) \in \Gamma$ iff either $\varphi_1 \notin \Gamma$ or $\varphi_2 \in \Gamma$.⁵

⁵Uses Group IV axioms.

Lemma 14

*Suppose that $\Gamma \subseteq \mathcal{L}_0$ and that Γ is maximally consistent.
Then Γ is satisfiable.*

Lemma 14

Suppose that $\Gamma \subseteq \mathcal{L}_0$ and that Γ is maximally consistent. Then Γ is satisfiable.

Lemma 15

Suppose that $\Gamma \subseteq \mathcal{L}_0$ and that Γ is consistent. Then there exists a set $\Gamma^ \subset \mathcal{L}_0$ such that $\Gamma \subseteq \Gamma^*$ and such that Γ^* is maximally consistent.*

Completeness Theorem

Theorem 16 (Completeness, version I)

Suppose that $\Gamma \subseteq \mathcal{L}_0$ and that Γ is consistent. Then Γ is satisfiable.

Completeness Theorem

Theorem 16 (Completeness, version I)

Suppose that $\Gamma \subseteq \mathcal{L}_0$ and that Γ is consistent. Then Γ is satisfiable.

This follows from Lemma ?? and then Lemma ??.

Completeness Theorem

Theorem 16 (Completeness, version I)

Suppose that $\Gamma \subseteq \mathcal{L}_0$ and that Γ is consistent. Then Γ is satisfiable.

This follows from Lemma ?? and then Lemma ??.

Theorem 17 (Completeness, version II)

Suppose that $\Gamma \subseteq \mathcal{L}_0$, $\varphi \in \mathcal{L}_0$ and that $\Gamma \models \varphi$. Then $\Gamma \vdash \varphi$.

Exercise 1 (Slaman & Woodin)

Exercise 1.4.1 (1)–(3)

Exercise (思考题)

Show that

1. $\vdash \neg\neg\alpha \rightarrow \alpha$
2. $\vdash (\alpha \rightarrow \neg\beta) \rightarrow (\beta \rightarrow \neg\alpha)$
3. $\vdash \alpha \rightarrow \neg\neg\alpha$
4. $\vdash (\alpha \rightarrow \beta) \leftrightarrow (\neg\beta \rightarrow \neg\alpha)$
5. If $\Gamma \cup \{\alpha\} \vdash \beta$ and $\Gamma \cup \{\neg\alpha\} \vdash \beta$, then $\Gamma \vdash \beta$.

Proof of Inference.

Let s be a Γ -proof of φ , t be a Γ -proof of $(\varphi \rightarrow \psi)$, Then by MP, $s + t + \langle \psi \rangle$ is a Γ -proof of ψ . □

Proof of Inference.

Let s be a Γ -proof of φ , t be a Γ -proof of $(\varphi \rightarrow \psi)$, Then by MP, $s + t + \langle \psi \rangle$ is a Γ -proof of ψ . □

Proof of Deduction.

From a $(\Gamma \cup \{\varphi\})$ -proof of ψ , $\langle \psi_1, \dots, \psi_n \rangle$, produce a Γ -proof of $(\varphi \rightarrow \psi)$. Prove by induction on $i \leq n$ that $\Gamma \vdash (\varphi \rightarrow \psi_i)$. Use Inference.

Proof of Inference.

Let s be a Γ -proof of φ , t be a Γ -proof of $(\varphi \rightarrow \psi)$, Then by MP, $s + t + \langle \psi \rangle$ is a Γ -proof of ψ . □

Proof of Deduction.

From a $(\Gamma \cup \{\varphi\})$ -proof of ψ , $\langle \psi_1, \dots, \psi_n \rangle$, produce a Γ -proof of $(\varphi \rightarrow \psi)$. Prove by induction on $i \leq n$ that $\Gamma \vdash (\varphi \rightarrow \psi_i)$. Use Inference.

CASE 1. $\psi_i \in \Gamma \cup \Delta$.

CASE 2. $\psi_i = \varphi$.

CASE 3. ψ_i is obtained from MP.



Proof of Inference.

Let s be a Γ -proof of φ , t be a Γ -proof of $(\varphi \rightarrow \psi)$, Then by MP, $s + t + \langle \psi \rangle$ is a Γ -proof of ψ . □

Proof of Deduction.

From a $(\Gamma \cup \{\varphi\})$ -proof of ψ , $\langle \psi_1, \dots, \psi_n \rangle$, produce a Γ -proof of $(\varphi \rightarrow \psi)$. Prove by induction on $i \leq n$ that $\Gamma \vdash (\varphi \rightarrow \psi_i)$. Use Inference.

CASE 1. $\psi_i \in \Gamma \cup \Delta$.

Need $\psi_i \rightarrow (\varphi \rightarrow \psi_i)$.

CASE 2. $\psi_i = \varphi$.

CASE 3. ψ_i is obtained from MP.



Proof of Inference.

Let s be a Γ -proof of φ , t be a Γ -proof of $(\varphi \rightarrow \psi)$, Then by MP, $s + t + \langle \psi \rangle$ is a Γ -proof of ψ . □

Proof of Deduction.

From a $(\Gamma \cup \{\varphi\})$ -proof of ψ , $\langle \psi_1, \dots, \psi_n \rangle$, produce a Γ -proof of $(\varphi \rightarrow \psi)$. Prove by induction on $i \leq n$ that $\Gamma \vdash (\varphi \rightarrow \psi_i)$. Use Inference.

CASE 1. $\psi_i \in \Gamma \cup \Delta$.

Need $\psi_i \rightarrow (\varphi \rightarrow \psi_i)$.

CASE 2. $\psi_i = \varphi$.

Need $\varphi \rightarrow \varphi$.

CASE 3. ψ_i is obtained from MP.

□

Proof of Inference.

Let s be a Γ -proof of φ , t be a Γ -proof of $(\varphi \rightarrow \psi)$, Then by MP, $s + t + \langle \psi \rangle$ is a Γ -proof of ψ . \square

Proof of Deduction.

From a $(\Gamma \cup \{\varphi\})$ -proof of ψ , $\langle \psi_1, \dots, \psi_n \rangle$, produce a Γ -proof of $(\varphi \rightarrow \psi)$. Prove by induction on $i \leq n$ that $\Gamma \vdash (\varphi \rightarrow \psi_i)$. Use Inference.

CASE 1. $\psi_i \in \Gamma \cup \Delta$.

Need $\psi_i \rightarrow (\varphi \rightarrow \psi_i)$.

CASE 2. $\psi_i = \varphi$.

Need $\varphi \rightarrow \varphi$.

CASE 3. ψ_i is obtained from MP.

Need

$(\varphi \rightarrow (\psi_{j_1} \rightarrow \psi_i)) \rightarrow [(\varphi \rightarrow \psi_{j_1}) \rightarrow (\varphi \rightarrow \psi_i)]$.

\square

Proof of Lemma ??.

Need $\varphi \rightarrow (\neg\varphi \rightarrow \psi)$. Use Inference.



Proof of Lemma ??.

Need $\varphi \rightarrow (\neg\varphi \rightarrow \psi)$. Use Inference. □

Proof of Lemma ??.

Suppose that Γ is consistent, both $\Gamma \cup \{\varphi\}$ and $\Gamma \cup \{\neg\varphi\}$ are inconsistent. Prove by contradiction.

Proof of Lemma ??.

Need $\varphi \rightarrow (\neg\varphi \rightarrow \psi)$. Use Inference. □

Proof of Lemma ??.

Suppose that Γ is consistent, both $\Gamma \cup \{\varphi\}$ and $\Gamma \cup \{\neg\varphi\}$ are inconsistent. Prove by contradiction.

First show $\Gamma \vdash \varphi$. Since $\Gamma \cup \{\neg\varphi\}$ is inconsistent, by Lemma ?? and Deduction, $\Gamma \vdash \neg\varphi \rightarrow \varphi$. Use $(\neg\varphi \rightarrow \varphi) \rightarrow \varphi \in \Delta$.

Proof of Lemma ??.

Need $\varphi \rightarrow (\neg\varphi \rightarrow \psi)$. Use Inference. □

Proof of Lemma ??.

Suppose that Γ is consistent, both $\Gamma \cup \{\varphi\}$ and $\Gamma \cup \{\neg\varphi\}$ are inconsistent. Prove by contradiction.

First show $\Gamma \vdash \varphi$. Since $\Gamma \cup \{\neg\varphi\}$ is inconsistent, by Lemma ?? and Deduction, $\Gamma \vdash \neg\varphi \rightarrow \varphi$. Use $(\neg\varphi \rightarrow \varphi) \rightarrow \varphi \in \Delta$.

As $\Gamma \cup \{\varphi\}$ is inconsistent, by Lemma ?? and Deduction, $\Gamma \vdash \varphi \rightarrow \neg\varphi$. We've shown $\Gamma \vdash \varphi$. By Inference, $\Gamma \vdash \neg\varphi$.

Proof of Lemma ??.

Need $\varphi \rightarrow (\neg\varphi \rightarrow \psi)$. Use Inference. □

Proof of Lemma ??.

Suppose that Γ is consistent, both $\Gamma \cup \{\varphi\}$ and $\Gamma \cup \{\neg\varphi\}$ are inconsistent. Prove by contradiction.

First show $\Gamma \vdash \varphi$. Since $\Gamma \cup \{\neg\varphi\}$ is inconsistent, by Lemma ?? and Deduction, $\Gamma \vdash \neg\varphi \rightarrow \varphi$. Use $(\neg\varphi \rightarrow \varphi) \rightarrow \varphi \in \Delta$.

As $\Gamma \cup \{\varphi\}$ is inconsistent, by Lemma ?? and Deduction, $\Gamma \vdash \varphi \rightarrow \neg\varphi$. We've shown $\Gamma \vdash \varphi$. By Inference, $\Gamma \vdash \neg\varphi$.

Therefore Γ is inconsistent. Contradiction! □

Proof of Corollary ??.

1. follows from maximality.

Proof of Corollary ??.

1. follows from maximality.
2. follows from 1.



Proof of Corollary ??.

1. follows from maximality.
2. follows from 1.



Proof of Lemma ??.

“ \Leftarrow ”. Suppose $\varphi_1 \notin \Gamma$. Use Corollary ?? and a logical axiom, $(\neg\varphi_1) \rightarrow (\varphi_1 \rightarrow \varphi_2)$, and Inference to get $(\varphi_1 \rightarrow \varphi_2) \in \Gamma$.

Proof of Corollary ??.

1. follows from maximality.
2. follows from 1.



Proof of Lemma ??.

“ \Leftarrow ”. Suppose $\varphi_1 \notin \Gamma$. Use Corollary ?? and a logical axiom, $(\neg\varphi_1) \rightarrow (\varphi_1 \rightarrow \varphi_2)$, and Inference to get $(\varphi_1 \rightarrow \varphi_2) \in \Gamma$.

Suppose $\varphi_2 \in \Gamma$. Need a logical axiom $\varphi_2 \rightarrow (\varphi_1 \rightarrow \varphi_2)$.

Proof of Corollary ??.

1. follows from maximality.
2. follows from 1.



Proof of Lemma ??.

“ \Leftarrow ”. Suppose $\varphi_1 \notin \Gamma$. Use Corollary ?? and a logical axiom, $(\neg\varphi_1) \rightarrow (\varphi_1 \rightarrow \varphi_2)$, and Inference to get $(\varphi_1 \rightarrow \varphi_2) \in \Gamma$.

Suppose $\varphi_2 \in \Gamma$. Need a logical axiom $\varphi_2 \rightarrow (\varphi_1 \rightarrow \varphi_2)$.

“ \Rightarrow ”. Suppose $\varphi_1 \in \Gamma$ and $\varphi_2 \notin \Gamma$. By Corollary ??, $\neg\varphi_2 \in \Gamma$. Need $\varphi_1 \rightarrow ((\neg\varphi_2) \rightarrow \neg(\varphi_1 \rightarrow \varphi_2))$. Therefore $(\neg(\varphi_1 \rightarrow \varphi_2)) \in \Gamma$. So it can't be that $(\varphi_1 \rightarrow \varphi_2) \in \Gamma$.



Proof of Lemma ??.

Need to give a truth assignment that works for Γ . Set $\nu(A_i) = T$ iff $A_i \in \Gamma$.



Proof of Lemma ??.

Need to give a truth assignment that works for Γ . Set $\nu(A_i) = T$ iff $A_i \in \Gamma$. □

Proof of Lemma ??.

Extend Γ to either $\Gamma \cup \{\varphi\}$ or $\Gamma \cup \{\neg\varphi\}$, which ever is consistent, for every $\varphi \in \mathcal{L}_0$. □