

# Variational Quantum Nuclear-Electronic Dynamics: Proton Transfer on Quantum Computers

Dilhan Manawadu, Edoardo Altamura

The Hartree Centre, STFC, Sci-Tech Daresbury, Warrington WA4 4AD, United Kingdom

## Quantum effects of proton transfer

- Proton transfer reactions are ubiquitous in many chemical and biological processes.
- Accurate description of proton transfer requires a quantum description of the kinetic nuclei that captures quantum tunnelling and zero-point energy.
- Several theoretical approaches to accurately model proton transfer like Ring Polymer Molecular Dynamics [1] and Multi-configuration time-dependent Hartree [2] have been developed, however they exhibit prohibitive scaling with the system size.

## Nuclear-Electronic Orbital (NEO) Theory

- NEO approach solves the mixed nuclear-electronic time-independent Schrödinger equation by constructing a mixed nuclear-electronic wavefunction [3].
- NEO framework has been complemented with existing electronic structure methods at different levels of theory like NEO Hartree-Fock (NEOHF) and NEO Complete Active Space Configuration Interaction (NEOCASCI) [4].
- These, however, still suffer from the scalability limitations of the underlying electronic structure methods.

## NEO approach for modelling proton transfer

- NEO framework has been extended for implementation on quantum hardware [5].
- Kovyrshin *et al.* utilised NEO framework with the Unitary Coupled Cluster (UCC) ansatz to calculate the energy barrier of proton transfer in malonaldehyde [6].
- This work demonstrated that NEO with UCC singlets and doublets could reach energy barrier predicted by NEOCASCI to within  $10^{-4}$  Hartree.
- Electronic-only CASCI calculation overestimates the barrier of proton transfer, which emphasises the importance of electron-nuclear correlation




Figure 1: Malonaldehyde

Method	$\Delta E/\text{Ha}$
CASCI	0.011962
NEOCASCI	0.005011
NEOUCCSD	0.004924
NEOUCCSDT	0.005007
NEOUCCSDTQ	0.005009

Table 1: Barrier for proton transfer in malonaldehyde calculated using different flavours of NEO. Values reproduced from Ref. [6].

## Quantum resource reduction using ADAPT-VQE

- NEO framework with UCC ansatz offers a way of modelling proton transfer in polynomial time using the Quantum Phase Estimation (QPE) [7] algorithm.
- However, these circuits are too deep to be meaningfully realised on current hardware for chemically interesting systems.
- Nykänen *et al.* extended the hardware efficient Adaptive Derivative-Assembled Pseudo-Trotter Ansatz Variational Quantum Eigensolver (ADAPT-VQE) for NEO framework. This method, titled Frozen Natural orbital (FNO)-NEO-ADAPT-VQE enables high fidelity circuit construction tailored for currently available quantum hardware [8].
- By using a FNO basis, they were able to reduce the qubit requirement, while the adaptive ansatz building led to shallower circuits.

## Adaptive approximate matrix product state preparation

- Adaptive Approximate Circuit Compilation (ADAPT-AQC), an adaptive algorithm based on approximate circuit compiling to construct high fidelity shallow circuits that approximates a given quantum circuits was recently proposed by Jadenberg *et al.* [9].
- In conjunction with ADAPT-VQE, this provides a toolbox of implementing quantum circuits to capture proton dynamics of small molecules on currently available quantum hardware.

## Approximate quantum compilation of proton-transfer dynamic for quantum processors

- In this work, we demonstrate a quantum computing pipeline based on NEO framework to model proton transfer of malonaldehyde.
- We suggest strategies to construct quantum circuits to tailor for current and future quantum hardware.
- We perform proton transfer barrier estimation simulations using realistic hardware noise models.

## Theory

- The dynamics of the system is described by the Time-Dependent Schrödinger Equation (TDSE)

$$i\hbar \frac{\partial}{\partial t} |\Psi(t)\rangle = [\hat{T}_p + \hat{V}(t)] |\Psi(t)\rangle, \quad (1)$$

where  $\hat{T}_p$  is the kinetic energy operator for the proton, and  $\hat{V}(t)$  is a time-dependent potential given by

$$\hat{V}(t) = \hat{V}_p(t) + \hat{V}_{ep}(t) + \hat{H}_e(t) \quad (2)$$

where  $\hat{V}_p(t)$  describes the interaction of the proton with the classical nuclear scaffold,  $\hat{V}_{ep}(t)$  is the proton-electron interaction and  $\hat{H}_e(t)$  is the electronic Hamiltonian contribution.

- We can write the NEOCI wavefunction with nuclear and electronic components  $|\Phi_v^n\rangle$  and  $|\Phi_\mu^e\rangle$  respectively as

$$|\Psi[\vec{C}(t)]\rangle = \sum_{\mu\nu} C_{\mu\nu}(t) |\Phi_\mu^e\rangle |\Phi_\nu^n\rangle. \quad (3)$$

- This leads to an equation of motion for the CI coefficients  $\vec{C}(t)$

$$\vec{H}(t)\vec{C}(t) = i\frac{\partial}{\partial t}\vec{C}(t), \quad (4)$$

where the matrix elements of  $\vec{H}(t)$  are defined by

$$H_{\kappa\lambda,\mu\nu}(t) = \langle \Phi_\kappa^e | \langle \Phi_\lambda^n | \hat{T}_p + \hat{V}(t) | \Phi_\mu^e \rangle | \Phi_\nu^n \rangle. \quad (5)$$

- Previously, Kovyrshin *et al.* explored the dynamics of this model through Suzuki-Trotter decomposition [10]. This approach is too computationally demanding for current hardware.
- To make our approach viable with current and near-future hardware, we explore the adiabatic regime of proton transfer by slowly varying the Hamiltonian. At each time step, the energy of the instantaneous proton-electron Hamiltonian is obtained by variationally minimising

$$E(t) = \min_{\vec{C}(t)} \frac{\langle \Psi[\vec{C}(t)] | \hat{T}_p + \hat{V}(t) | \Psi[\vec{C}(t)] \rangle}{\langle \Psi[\vec{C}(t)] | \Psi[\vec{C}(t)] \rangle}. \quad (6)$$

## Pipeline for proton transfer simulation




Figure 2: Starting with a variationally optimized FNO-NEO-ADAPT-VQE ansatz, we generate two sets of ADAPT-VQE circuits using different convergence thresholds. These are further truncated via ADAPT-AQC to optimize circuit depth. Deeper circuits are simulated without noise to emulate early fault tolerance, while shallower ones are simulated on the noisy IBM Fez model with ZNE.

## Results

Soon to be published!

## Acknowledgements

This work was supported by the Hartree National Centre for Digital Innovation, a UK Government-funded collaboration between STFC and IBM.

## References

- [1] Habershon, S.; Manolopoulos, D. E.; Markland, T. E.; Miller III, T. F. *Annual Review of Physical Chemistry*, Vol 62 **2013**, 64, 387–413, Publisher: Annual Reviews.
- [2] Beck, M. H.; Jäckle, A.; Worth, G. A.; Meyer, H.-D. **2000**, 324, 1–105, Publisher: Elsevier.
- [3] Webb, S. P.; Iordanov, T.; Hammes-Schiffer, S. *Journal of Chemical Physics* **2002**, 117, 4106–4118, Publisher: American Institute of Physics.
- [4] Pavošević, F.; Culpitt, T.; Hammes-Schiffer, S. *Chemical Reviews* **2020**, 120, 4222–4253, Publisher: ACS Publications.
- [5] Pavošević, F.; Hammes-Schiffer, S. *Journal of Chemical Theory and Computation* **2021**, 17, 3252–3258.
- [6] Kovyrshin, A.; Skogh, M.; Broo, A.; Mensa, S.; Sahin, E.; Crain, J.; Tavernelli, I. *Journal of Chemical Physics* **2023**, 158, 214119, Publisher: AIP Publishing.
- [7] Aspuru-Guzik, A.; Dutoi, A. D.; Love, P. J.; Head-Gordon, M. *Science* **2005**, 309, 1704–1707.
- [8] Nykänen, A.; Miller, A.; Talarico, W.; Knecht, S.; Kovyrshin, A.; Skogh, M.; Tornberg, L.; Broo, A.; Mensa, S.; Symons, B. C. B.; Sahin, E.; Crain, J.; Tavernelli, I.; Pavošević, F. *Journal of Chemical Theory and Computation* **2023**, 19, 9269–9277.
- [9] Jaderberg, B.; Pennington, G.; Marshall, K. V.; Anderson, L. W.; Agarwal, A.; Lindoy, L. P.; Rungger, I.; Mensa, S.; Crain, J. Variational preparation of normal matrix product states on quantum computers. 2025; <http://arxiv.org/abs/2503.09683>, arXiv:2503.09683 [quant-ph].
- [10] Kovyrshin, A.; Skogh, M.; Tornberg, L.; Broo, A.; Mensa, S.; Sahin, E.; Symons, B. C. B.; Crain, J.; Tavernelli, I. *Journal of Physical Chemistry Letters* **2023**, 14, 7065–7072, Publisher: ACS Publications.