

Day-61, Jan-30, 2025 (Mogh 17, 2081 B.S.)

Introduction to Covariance

X : age of a child

Y_1 : height of the child (in)

Age (X)	Height (Y_1)
6	50
7	52
8	55
9	57
10	60
11	62
12	64
13	65
14	67
15	68

Y_2 : grades in a test

Age (X)	Grades (Y_2)
6	5
7	7
8	8
9	3
10	1
11	1
12	6
13	10
14	2


Y_3 : number of naps per day

Age (X)	Naps per day (Y_3)
6	8
7	7
8	6
9	5
10	4
11	3
12	2
13	1
14	1
15	0


Introduction to Covariance

X : age of a child


Y_1 : height of the child (in)



Y_2 : grades in a test



Y_3 : number of naps per day



To help us better
visualize what's going on

DeepLearning.AI


DeepLearning.AI

and the third one is like this.


Covariance means variance of two variables together

Correlation means having relationship with each other variables.


Mean?



$$\mu_x = 10.5$$



$$\mu_x = 10.5$$




$$\mu_x = 10.5$$




Now we can also look
at the variances.


Horizontal (X) Variance



$$Var(X) = 9.17$$



$$Var(X) = 9.17$$




$$Var(X) = 9.17$$




which are 9.17 because it's
the same data-set for ages.


Vertical (Y) Variance



$$Var(Y) = 39.56$$




$$Var(Y) = 9.78$$




$$Var(Y) = 7.57$$

which are 39.56, 9.78 and 7.57.


Still no Way To Compare Them



Covariance > 0



Covariance ≈ 0




Covariance < 0

the covariance of the
second plot is almost zero,


So, what is the step?

First Step: Center Them



DeepLearning.AI


First Step: Center Them



DeepLearning.AI


this really nice plots where

Second Step: Notice Trend



and the lack of
trends in the middle.

Second Step: Notice Trend




move to Right Up

the coordinates move up,
and the one in the middle,

DeepLearning.AI


Second Step: Notice Trend



there seems to be no rule
and on the one on the right,

DeepLearning.AI



Second Step: Notice Trend



if you move to the right,
the coordinates move down.

DeepLearning.AI




Positives and Negatives



The quantities always have the same sign.

DeepLearning.AI




Positives and Negatives



the one in the middle, nothing seems to happen.

DeepLearning.AI




Positives and Negatives



Now let's look at the product of coordinates,

DeepLearning.AI




Positives and Negatives



Now, what if I were to add

DeepLearning.AI


Covariance




This sum is called
the co-variance.

DeepLearning.AI


Covariance



$$Cov(X, Y) > 0$$



$$Cov(X, Y) \approx 0$$



$$Cov(X, Y) < 0$$

like age and naps per day.

DeepLearning.AI

1

Covariance

$$Cov(X, Y) = \sum xy$$

Almost...

$$Cov(X, Y) = \frac{\sum (x_i - \mu_x)(y_i - \mu_y)}{n}$$

take the average of
all these products.

DeepLearning.AI

2

Covariance Formula

Age vs Height

Covariance > 0

$$\mu_x = 10.5 \quad \mu_y = 60$$

$$Cov(X, Y) = \frac{\sum (x_i - \mu_x)(y_i - \mu_y)}{n}$$

$$Cov(X, Y) = \frac{170}{10} = 17 > 0$$

that when age grows,
height grows.

$$\sum = 170$$

Age (x)	Height (y)	$(x_i - \mu_x)$	$(y_i - \mu_y)$	$(x_i - \mu_x)(y_i - \mu_y)$
6	50	-4.5	-10	45
7	52	-3.5	-8	28
8	55	-2.5	-5	12.5
9	57	-1.5	-3	4.5
10	60	-0.5	0	0
11	62	0.5	2	1
12	64	1.5	4	6
13	65	2.5	5	12.5
14	67	3.5	7	24.5
15	68	4.5	8	36

DeepLearning.AI

Covariance Formula

Age vs Height

Covariance > 0

$$\mu_x = 10.5 \quad \mu_y = 60$$

$$Cov(X, Y) = \frac{\sum (x_i - \mu_x)(y_i - \mu_y)}{n}$$

$$Cov(X, Y) = \frac{170}{10} = 17 > 0$$

that when age grows,
height grows.

(1)

Age (x)	Height (y)	$(x_i - \mu_x)$	$(y_i - \mu_y)$	$(x_i - \mu_x)(y_i - \mu_y)$
6	50	-4.5	-10	45
7	52	-3.5	-8	28
8	55	-2.5	-5	12.5
9	57	-1.5	-3	4.5
10	60	-0.5	0	0
11	62	0.5	2	1
12	64	1.5	4	6
13	65	2.5	5	12.5
14	67	3.5	7	24.5
15	68	4.5	8	36

$\sum = 170$

DeepLearning.AI

Covariance Formula

Age vs Grades

Covariance ≈ 0

$$\mu_x = 10.5 \quad \mu_y = 5$$

$$Cov(X, Y) = \frac{\sum (x_i - \mu_x)(y_i - \mu_y)}{n}$$

$$Cov(X, Y) = \frac{1}{10} = 0.1 \approx 0$$

or at least very
little influence for

Age (x)	Grades (y)	$(x_i - \mu_x)$	$(y_i - \mu_y)$	$(x_i - \mu_x)(y_i - \mu_y)$
6	5	-4.5	0	0
7	7	-3.5	2	-7
8	8	-2.5	3	-7.5
9	3	-1.5	-2	3
10	1	-0.5	-4	2
11	1	0.5	-4	-2
12	6	1.5	1	1.5
13	10	2.5	5	12.5
14	2	3.5	-3	-10.5
15	7	4.5	2	9

$\sum = 1$

DeepLearning.AI

-ve

and .

No

Covariance

Covariance Formula

(2)

Age (x)	Naps (y)	$(x_i - \mu_x)$	$(y_i - \mu_y)$	$(x_i - \mu_x)(y_i - \mu_y)$
6	8	-4.5	4.3	-19.35
7	7	-3.5	3.3	-11.55
8	6	-2.5	2.3	-5.75
9	5	-1.5	1.3	-1.95
10	4	-0.5	0.3	-0.15
11	3	0.5	-0.7	-0.35
12	2	1.5	-1.7	-2.55
13	1	2.5	-2.7	-6.75
14	1	3.5	-2.7	-9.45
15	0	4.5	-3.7	-16.65

$\sum = -74.5$

That is negative
because as you know,

Comparing Correlations


(3)



Age vs Height

Covariance > 0



$$Cov(x, y) = 17$$



Age vs Grades

Covariance ≈ 0

$$Cov(x, y) = 0.1$$



Age vs Naps per Day

Covariance < 0

$$Cov(x, y) = -7.45$$

be negatively correlated
have a covariance of -7.45,


Covariance of a Probability Distribution: Motivation

X and Y are playing 3 games to either win or lose a dollar

GAME 1

a: Both players win \$1 each


b: Both players lose \$1 each



GAME 2


c: X wins \$1 and Y loses \$1

d: X loses \$1 and Y win \$1



GAME 3

a, b, c or d




So four possibilities, basically a, b, c and d all with probability 1/4.

DeepLearning.AI

Covariance of a Probability Distribution: Motivation


GAME 1



$$\mathbb{E}[X_1] = \frac{1}{2}(1) + \frac{1}{2}(-1) = 0$$

$$\mathbb{E}[Y_1] = \frac{1}{2}(1) + \frac{1}{2}(-1) = 0$$

GAME 2




$$\mathbb{P}(d) = 1/2$$

$$\mathbb{P}(c) = 1/2$$

They win a dollar or lose a dollar with probability 1/2.


GAME 3



DeepLearning.AI

Covariance of a Probability Distribution: Motivation

GAME 1



$$\mathbb{E}[X_1] = 0$$


$$\mathbb{E}[Y_1] = 0$$

$$\mathbb{E}[X_2] = \frac{1}{2}(1) + \frac{1}{2}(-1) = 0$$

$$\mathbb{E}[Y_2] = \frac{1}{2}(1) + \frac{1}{2}(-1) = 0$$


So again, it's 0.

GAME 3



Covariance of a Probability Distribution: Motivation


GAME 1



$$\mathbb{E}[X_1] = 0$$

$$\mathbb{E}[Y_1] = 0$$


GAME 2



$$\mathbb{E}[X_2] = 0$$

$$\mathbb{E}[Y_2] = 0$$

GAME 3




$$\mathbb{E}[X_3] = 2\left(\frac{1}{4}(1)\right) + 2\left(\frac{1}{4}(-1)\right) = 0$$

$$\mathbb{E}[Y_3] = 2\left(\frac{1}{4}(1)\right) + 2\left(\frac{1}{4}(-1)\right) = 0$$

So these games are basically the same if you only think of one player at a time.

DeepLearning.AI

Covariance of a Probability Distribution: Motivation




$$\mathbb{E}[X_1] = 0 \quad \mathbb{E}[Y_1] = 0$$

$$Var(X_1) = \mathbb{E}[X_1^2] - \mathbb{E}[X_1]^2 = \frac{1}{2}(-1)^2 + \frac{1}{2}(1)^2 - 0^2 = 1$$


So all we care is the expected value
of X_1 squared and that's going to be 1.

DeepLearning.AI



$$\mathbb{E}[X_2] = 0 \quad \mathbb{E}[Y_2] = 0$$

①



$$\mathbb{E}[X_3] = 0 \quad \mathbb{E}[Y_3] = 0$$

$$Var(X_1) = \mathbb{E}[X_1^2] - \mathbb{E}[X_1]^2 = \frac{1}{2}(-1)^2 + \frac{1}{2}(1)^2 - 0^2 = 1$$


So all we care is the expected value
of X_1 squared and that's going to be 1.

DeepLearning.AI

Where does difference lie?

Covariance of a Probability Distribution: Motivation

GAME 1




$$\mathbb{E}[X_1] = 0 \quad \mathbb{E}[Y_1] = 0$$

$$Var(X_1) = 1$$

$$Var(Y_1) = 1$$

GAME 2




$$\mathbb{E}[X_2] = 0 \quad \mathbb{E}[Y_2] = 0$$

$$Var(X_2) = 1$$

$$Var(Y_2) = 1$$

GAME 3



$$\mathbb{E}[X_3] = 0 \quad \mathbb{E}[Y_3] = 0$$

$$Var(X_3) = 1$$

$$Var(Y_3) = 1$$


DeepLearning.AI

How similar are these 3 games for player X and player Y?

But obviously they are different games.

Covariance of a Probability Distribution: Motivation

GAME 1




$$\mathbb{E}[X_1] = 0 \quad \mathbb{E}[Y_1] = 0$$

$$Var(X_1) = 1$$

$$Var(Y_1) = 1$$

DeepLearning.AI

GAME 2



$$\mathbb{E}[X_2] = 0 \quad \mathbb{E}[Y_2] = 0$$


$$Var(X_2) = 1$$

$$Var(Y_2) = 1$$

The difference lies in that you have to
look at both players at the same time,

if you
look one player at a time

GAME 3



$$\mathbb{E}[X_3] = 0 \quad \mathbb{E}[Y_3] = 0$$

$$Var(X_3) = 1$$


$$Var(Y_3) = 1$$

$$\mathbb{E}[X_2] = 0 \quad \mathbb{E}[Y_2] = 0$$

$$Var(X_3) = 1$$

$$Var(Y_3) = 1$$

GAME 2




$$\mathbb{E}[X_2] = 0 \quad \mathbb{E}[Y_2] = 0$$

$$Var(X_2) = 1$$

$$Var(Y_2) = 1$$

GAME 3




$$\mathbb{E}[X_3] = 0 \quad \mathbb{E}[Y_3] = 0$$

$$Var(X_3) = 1$$

$$Var(Y_3) = 1$$

Covariance of a Probability Distribution: Motivation

GAME 1



$$\mathbb{E}[X_1] = 0 \quad \mathbb{E}[Y_1] = 0$$

$$Var(X_1) = 1$$

$$Var(Y_1) = 1$$

DeepLearning.AI

Covariance of X and Y : $Cov(X, Y)$

How similar are these 3 games for player X and player Y?

$$\mathbb{E}[X_1] = 0 \quad \mathbb{E}[Y_1] = 0$$

$$Var(X_1) = 1 \quad Var(Y_1) = 1$$

So we need to look at the covariance.

To know the difference!

Covariance of a Probability Distribution: Motivation

Covariance of X and Y : $\text{Cov}(X, Y)$

$$\text{Cov}(X, Y) = \frac{\sum (x_i - \mu_x)(y_i - \mu_y)}{n}$$


we're going to calculate it for
the three games.

DeepLearning.AI

1

Covariance of a Probability Distribution: Motivation

GAME 1



$$\text{Cov}(X, Y) = \frac{\sum (x_i - \mu_x)(y_i - \mu_y)}{n}$$


x	y	$(x_i - \mu_x)$	$(y_i - \mu_y)$	$(x_i - \mu_x)(y_i - \mu_y)$
1	1	1	1	1
-1	-1	-1	-1	-1

$$\text{Cov}(X, Y) = \frac{\sum (x_i - \mu_x)(y_i - \mu_y)}{n} = \frac{2}{2} = 1$$

So the covariance of the first game is 1.

Covariance of a Probability Distribution: Motivation

GAME 1



$$\mu_x = \mathbb{E}[X_1] = 0$$

$$\mu_y = \mathbb{E}[Y_1] = 0$$

$$\text{Cov}(X, Y) = \frac{\sum (x_i - \mu_x)(y_i - \mu_y)}{n}$$

x	y	$(x_i - \mu_x)$	$(y_i - \mu_y)$	$(x_i - \mu_x)(y_i - \mu_y)$
1	1	1	1	1
-1	-1	-1	-1	-1

$$\text{Cov}(X, Y) = \frac{\sum (x_i - \mu_x)(y_i - \mu_y)}{n} = \frac{2}{2} = 1$$

the more player one wins,
the more player two wins.

Either both 'L'
Either both 'S'.

So, the $\text{Cov}(X, Y)$ is
+ve but the

Game 1 shows +ve


Covariance =

DeepLearning.AI

2

1

Covariance of a Probability Distribution: Motivation

GAME 2

$$\mu_x = \mathbb{E}[X_2] = 0$$

$$\mu_y = \mathbb{E}[Y_2] = 0$$

$$Cov(X, Y) = \frac{\sum (x_i - \mu_x)(y_i - \mu_y)}{n}$$

x	y	$(x_i - \mu_x)$	$(y_i - \mu_y)$	$(x_i - \mu_x)(y_i - \mu_y)$
1	-1	1	-1	-1
-1	1	-1	1	-1

$$Cov(X, Y) = \frac{\sum (x_i - \mu_x)(y_i - \mu_y)}{n} = \frac{-2}{2} = -1$$


So this one has a covariance of -1,
which is reflected in the fact that

DeepLearning.AI

Game 2.

2

Covariance of a Probability Distribution: Motivation

GAME 2

$$\mu_x = \mathbb{E}[X_2] = 0$$

$$\mu_y = \mathbb{E}[Y_2] = 0$$

$$Cov(X, Y) = \frac{\sum (x_i - \mu_x)(y_i - \mu_y)}{n}$$

x	y	$(x_i - \mu_x)$	$(y_i - \mu_y)$	$(x_i - \mu_x)(y_i - \mu_y)$
1	-1	1	-1	-1
-1	1	-1	1	-1


$$Cov(X, Y) = \frac{\sum (x_i - \mu_x)(y_i - \mu_y)}{n} = \frac{-2}{2} = -1$$

And that means that either one is happy and the other one is sad, or

DeepLearning.AI

③ No pattern All states

Covariance of a Probability Distribution: Motivation

GAME 3

$$\mu_x = \mathbb{E}[X_3] = 0$$

$$\mu_y = \mathbb{E}[Y_3] = 0$$

$$Cov(X, Y) = \frac{\sum (x_i - \mu_x)(y_i - \mu_y)}{n}$$

x	y	$(x_i - \mu_x)$	$(y_i - \mu_y)$	$(x_i - \mu_x)(y_i - \mu_y)$
1	1	1	1	1
1	-1	1	-1	-1
-1	1	-1	1	-1
-1	-1	-1	-1	1


$$Cov(X, Y) = \frac{\sum (x_i - \mu_x)(y_i - \mu_y)}{n} = \frac{0}{4} = 0$$

So this covariance is 0.

DeepLearning.AI

100DaysOfMaths_@dilli_hangrae

Covariance of a Probability Distribution: Motivation

GAME 3

$$\mu_x = \mathbb{E}[X_3] = 0$$

$$\mu_y = \mathbb{E}[Y_3] = 0$$

$$Cov(X, Y) = \frac{\sum (x_i - \mu_x)(y_i - \mu_y)}{n}$$

x	y	$(x_i - \mu_x)$	$(y_i - \mu_y)$	$(x_i - \mu_x)(y_i - \mu_y)$
1	1	1	1	1
1	-1	1	-1	-1
-1	1	-1	1	-1
-1	-1	-1	-1	1


$$Cov(X, Y) = \frac{\sum (x_i - \mu_x)(y_i - \mu_y)}{n} = \frac{0}{4} = 0$$

And that shows the fact that there's not really a pattern among these.

DeepLearning.AI


Covariance of a Probability Distribution: Motivation

GAME 1




$$\text{Cov}(X, Y) = 1$$

GAME 2



$$\text{Cov}(X, Y) = -1$$

GAME 3



$$\text{Cov}(X, Y) = 0$$

In the second one a covariance of -1,
and in the third one a covariance of 0.

DeepLearning.AI

So, the final Covariance of a Probability Distribution are five, 0 and -ve.

So, let's introduce 3rd Game with three conditions of shifts for unequal probabilities:


Covariance of a Probability Distribution: Motivation

GAME 4

a: Both players win \$1 each

b: Both players lose \$1 each

c: Neither players wins nor lose anything



Step 1

Both players lose a dollar and nothing happens.

DeepLearning.AI


Covariance of a Probability Distribution: Motivation

GAME 4

a: Both players win \$1 each $P(a) = 1/2$

b: Both players lose \$1 each $P(b) = 1/3$

c: Neither players wins nor lose anything $P(c) = 1/6$



Unequal Probabilities

Step 2

So now if we look at only one player, the X player,

DeepLearning.AI


Covariance of a Probability Distribution: Motivation

GAME 4

a: Both players win \$1 each $P(a) = 1/2$

b: Both players lose \$1 each $P(b) = 1/3$

c: Neither players wins nor lose anything $P(c) = 1/6$



$$\mathbb{E}[X_4] = \frac{1}{6}$$

$$Var(X_4) = \sum_{i=1}^N (x_i - \mu_x)^2 \cdot P(x_i)$$

$$= \frac{1}{2}\left(1 - \frac{1}{6}\right)^2 + \frac{1}{6}\left(0 - \frac{1}{6}\right)^2 + \frac{1}{3}\left(-1 - \frac{1}{6}\right)^2$$

$$= 0.806$$

then multiply with the probabilities and

we get 0.806.

Unequal Probabilities

$$\text{So, } Var(X_4) \Rightarrow 0.806$$

$$Var(Y_4) = 0.806$$

$$\text{So, what's } Cov(X_4, Y_4) = ?$$

$$\text{use } Cov(X_4, Y_4) = P_{XY} (x_i, y_i)(x_i - \mu_x)(y_i - \mu_y)$$

DeepLearning.AI

Covariance of Probability Distributions

$$Cov(X, Y) = \frac{\sum (x_i - \mu_x)(y_i - \mu_y)}{n} = \frac{1}{n} \sum (x_i - \mu_x)(y_i - \mu_y)$$

$$Cov(X, Y) = \sum p_{XY}(x_i, y_i)(x_i - \mu_x)(y_i - \mu_y)$$

equal probabilities

unequal probabilities

DeepLearning.AI

So this is what happens in general.

$$\text{Ex-xx. } Cov(X, Y) \neq E(XY) - E(X)E(Y)$$


$$Cov(X, Y) \Rightarrow 18.014 - 4.903 \times 5.280 \\ \Rightarrow -7.878$$

-ve Covariance

Slope
Downward

Next is Covariance Matrix !

Covariance of a Probability Distribution: Motivation



Unequal Probabilities

$$Cov(X, Y) = \sum p_{XY}(x_i, y_i)(x_i - \mu_x)(y_i - \mu_y) = E[XY] - E[X]E[Y]$$

$$= \frac{1}{2}\left(1 - \frac{1}{6}\right)^2 + \frac{1}{6}\left(0 - \frac{1}{6}\right)^2 + \frac{1}{3}\left(-1 - \frac{1}{6}\right)^2$$

$$Cov(X, Y) = 0.806$$

$$\mu_x = \mu_y = E[X_4] = E[Y_4] = 1/6$$

$$Var(X_4) = Var(Y_4) = 0.806$$

DeepLearning.AI

the covariance XY is 0.806.


Covariance?

$$E(X) = 4.903$$

$$E(Y) = 5.280$$

$$E(XY) = 18.014$$


Ex-xx




Recall that we had two marginal distributions, X and Y and

DeepLearning.AI


Covariance Matrix



$$\begin{aligned}Var(X) &= Var(Y) = 1 \\Cov(X, Y) &= 1\end{aligned}$$



$$\begin{aligned}Var(X) &= Var(Y) = 1 \\Cov(X, Y) &= -1\end{aligned}$$




$$\begin{aligned}Var(X) &= Var(Y) = 1 \\Cov(X, Y) &= 0\end{aligned}$$

the same expectations and covariance of 1 minus 1 and zero.

DeepLearning.AI

Covariance Matrix




$$\begin{aligned}Var(X) &= 8.526 \\Var(Y) &= 10.163 \\Cov(X, Y) &= -7.878\end{aligned}$$

because the higher the waiting time, the lower the ratings.

DeepLearning.AI

Covariance Matrix




So we would have ten pairs over here and that's going to be the covariance matrix.

DeepLearning.AI

Covariance Matrix

1



$$\text{Var}(X) = 8.526$$

$$\text{Var}(Y) = 10.163$$

$$\text{Cov}(X, Y) = -7.878$$


	X	Y
X	$\text{Var}(X)$	$\text{Cov}(X, Y)$
Y	$\text{Cov}(X, Y)$	$\text{Var}(Y)$

the covariances on the antidiagonal.

DeepLearning.AI

Covariance Matrix

2



	X	Y
X	$\text{Var}(X)$	$\text{Cov}(X, Y)$
Y	$\text{Cov}(X, Y)$	$\text{Var}(Y)$

$$\begin{bmatrix} \text{Var}(X) & \text{Cov}(X, Y) \\ \text{Cov}(X, Y) & \text{Var}(Y) \end{bmatrix}$$

$$\begin{bmatrix} 8.526 & -7.878 \\ -7.878 & 10.163 \end{bmatrix}$$


Covariance Matrix

Covariance matrix is very, very useful in statistics and also in machine learning.

DeepLearning.AI

Covariance Matrix

coursera.org - To exit full screen, press esc



Age vs Height

$$\text{Var}(X) = 9.17$$

$$\text{Var}(Y) = 39.56$$

$$\text{Cov}(X, Y) = 17$$

$$\begin{bmatrix} \text{Var}(X) & \text{Cov}(X, Y) \\ \text{Cov}(X, Y) & \text{Var}(Y) \end{bmatrix}$$

$$\begin{bmatrix} 9.17 & 17 \\ 17 & 39.56 \end{bmatrix}$$

Matrix with Diagonal Var


and antidiagonal Cov are Covariance Matrix

$$\boxed{\begin{bmatrix} \text{Var}(X) & \text{Cov}(X, Y) & \text{Var}(Y) \\ \text{Cov}(X, Y) & \text{Var}(Y) \end{bmatrix}}$$

DeepLearning.AI


100DaysOfMaths_@dilli_hangrae

Covariance Matrix



$$\begin{aligned}Var(X) &= 1 \\Var(Y) &= 1 \\Cov(X, Y) &= 1\end{aligned}$$

$$\begin{bmatrix}Var(X) & Cov(X, Y) \\Cov(X, Y) & Var(Y)\end{bmatrix}$$



$$\begin{aligned}Var(X) &= 1 \\Var(Y) &= 1 \\Cov(X, Y) &= -1\end{aligned}$$

They are these two for games 1 and 2.

DeepLearning.AI

1

Covariance of a Joint Continuous Distribution

Dataset with 3 Variables:

X	Y	Z	Var(X)	Cov(X, Y)
			Var(Y)	Cov(X, Z)
			Var(Z)	Cov(Y, Z)

X	Y	Z	Var(X)	Cov(X, Y)	Cov(X, Z)
Y			Cov(X, Y)	Var(Y)	Cov(Y, Z)
Z			Cov(X, Z)	Cov(Y, Z)	Var(Z)

In the coordinates X set we have
the covariance of X set, etc.

DeepLearning.AI

2

Covariance of a Joint Continuous Distribution

Dataset with 5 Variables:

V	W	X	Y	Z	Var(V)	Cov(V, W)
					Var(W)	Cov(V, X)
					Var(X)	Cov(V, Y)
					Var(Y)	Cov(V, Z)
					Var(Z)	Cov(W, X)
					⋮	
						Cov(Y, Z)

V	W	X	Y	Z	Var(V)	Cov(V, W)	Cov(V, X)	Cov(V, Y)	Cov(V, Z)
W					Cov(V, W)	Var(W)	Cov(W, X)	Cov(W, Y)	Cov(W, Z)
X					Cov(V, X)	Cov(W, X)	Var(X)	Cov(X, Y)	Cov(X, Z)
Y					Cov(V, Y)	Cov(W, Y)	Cov(X, Y)	Var(Y)	Cov(Y, Z)
Z					Cov(V, Z)	Cov(W, Z)	Cov(X, Z)	Cov(Y, Z)	Var(Z)

covariance of the row variable and
the column variable.

DeepLearning.AI

3

Covariance of a Joint Continuous Distribution

$$\sum =$$


Covariance Matrix

V	W	X	Y	Z	Var(V)	Cov(V, W)	Cov(V, X)	Cov(V, Y)	Cov(V, Z)
W					Cov(V, W)	Var(W)	Cov(W, X)	Cov(W, Y)	Cov(W, Z)
X					Cov(V, X)	Cov(W, X)	Var(X)	Cov(X, Y)	Cov(X, Z)
Y					Cov(V, Y)	Cov(W, Y)	Cov(X, Y)	Var(Y)	Cov(Y, Z)
Z					Cov(V, Z)	Cov(W, Z)	Cov(X, Z)	Cov(Y, Z)	Var(Z)

This matrix is going to be called sigma,
the covariance matrix.

DeepLearning.AI

Correlation Coefficient




$$\begin{aligned}Var(X) &= 9.17 \\Var(Y) &= 7.57\end{aligned}$$

$$Cov(X, Y) = -7.45$$

When this is a positive value it tells us that


DeepLearning.AI



$$\begin{aligned}Var(X) &= 9.17 \\Var(Y) &= 39.56\end{aligned}$$

$$Cov(X, Y) = 17$$

Correlation Coefficient




$$\begin{aligned}Var(X) &= 9.17 \\Var(Y) &= 7.57\end{aligned}$$

$$Cov(X, Y) = -7.45$$

grow however because of 17 is bigger than 7.45 forgetting about


DeepLearning.AI



$$\begin{aligned}Var(X) &= 9.17 \\Var(Y) &= 39.56\end{aligned}$$

$$Cov(X, Y) = 17$$

Correlation Coefficient




$$\begin{aligned}Var(X) &= 9.17 \\Var(Y) &= 9.78 \\Cov(X, Y) &= 0.1\end{aligned}$$

$$\begin{aligned}\text{Correlation Coefficient} &= \frac{Cov(X, Y)}{\sqrt{Var(X)} \cdot \sqrt{Var(Y)}} \\&= \frac{0.1}{\sqrt{9.17} \cdot \sqrt{9.78}}\end{aligned}$$

$$\approx 0.01$$

DeepLearning.AI

Correlation Coefficient



$$\begin{aligned}Var(X) &= 8.526 \\Var(Y) &= 10.163 \\Cov(X, Y) &= -7.878\end{aligned}$$

$$\begin{aligned}\text{Correlation Coefficient} &= \frac{Cov(X, Y)}{\sqrt{Var(X)} \cdot \sqrt{Var(Y)}} \\&= \frac{-7.878}{\sqrt{8.562} \cdot \sqrt{10.163}}\end{aligned}$$

$$\approx -0.845$$

DeepLearning.AI

Correlation Coefficient

$$\text{Correlation Coefficient} = \frac{\text{Cov}(X, Y)}{\sigma_x \cdot \sigma_y} = \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X)} \cdot \sqrt{\text{Var}(Y)}}$$



DeepLearning.AI


Multivariate Gaussian Distribution

For a single variable, X

$$f_X(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{1}{2} \frac{(x-\mu)^2}{\sigma^2}}$$

Parameters:

- μ : center of the bell
- σ : spread of the bell



Now here's a question. What if you had more than one variable?

DeepLearning.AI

Multivariate Gaussian Distribution: an Example


Two variables

H : Height of an adult in inches

$$H \sim \mathcal{N}(\mu_H, \sigma_H)$$

W : Weight of an adult in pounds


$$W \sim \mathcal{N}(\mu_W, \sigma_W)$$



then you can look at the margins
and notice that they are both normal

DeepLearning.AI

Multivariate Gaussian Distribution: an Example




If W, H were independent

$$f_{HW}(h, w) = f_H(h)f_W(w)$$

And rearranging the terms a little bit,
you would get this expression over here.

DeepLearning.AI

Multivariate Gaussian Distribution: an Example




If W, H were independent

$$\begin{aligned} f_{HW}(h, w) &= f_H(h)f_W(w) \\ &= \frac{1}{\sqrt{2\pi}\sigma_H} e^{-\frac{1}{2}\frac{(h-\mu_H)^2}{\sigma_H^2}} \cdot \frac{1}{\sqrt{2\pi}\sigma_W} e^{-\frac{1}{2}\frac{(w-\mu_W)^2}{\sigma_W^2}} \\ &= \frac{1}{2\pi\sigma_H\sigma_W} e^{-\frac{1}{2}\left(\frac{(h-\mu_H)^2}{\sigma_H^2} + \frac{(w-\mu_W)^2}{\sigma_W^2}\right)} \end{aligned}$$


DeepLearning.AI

Multivariate Gaussian Distribution: an Example





DeepLearning.AI

is completely symmetric in a dependence case you see that distribution is elongated



Multivariate Gaussian Distribution: an Example



curves are high up and the purple curves are down.

DeepLearning.AI

Multivariate Gaussian Distribution: an Example

If W, H were independent

$$\begin{aligned} f_{HW}(h, w) &= f_H(h)f_W(w) \\ &= \frac{1}{2\pi\sigma_H\sigma_W} e^{-\frac{1}{2}\left(\frac{(h-\mu_H)^2}{\sigma_H^2} + \frac{(w-\mu_W)^2}{\sigma_W^2}\right)} \\ &\quad \xrightarrow{\left\| \begin{bmatrix} \frac{h-\mu_H}{\sigma_H} \\ \frac{w-\mu_W}{\sigma_W} \end{bmatrix} \right\|_2^2} = \left[\begin{bmatrix} h-\mu_H \\ w-\mu_W \end{bmatrix} \right] \begin{bmatrix} \frac{1}{\sigma_H^2} & 0 \\ 0 & \frac{1}{\sigma_W^2} \end{bmatrix} \left[\begin{bmatrix} h \\ w \end{bmatrix} - \begin{bmatrix} \mu_H \\ \mu_W \end{bmatrix} \right] \\ &= \left(\begin{bmatrix} h \\ w \end{bmatrix} - \begin{bmatrix} \mu_H \\ \mu_W \end{bmatrix} \right)^T \begin{bmatrix} \sigma_H^2 & 0 \\ 0 & \sigma_W^2 \end{bmatrix}^{-1} \left(\begin{bmatrix} h \\ w \end{bmatrix} - \begin{bmatrix} \mu_H \\ \mu_W \end{bmatrix} \right) \end{aligned}$$

diagonal matrix with the variances in each element

Multivariate Gaussian Distribution: an Example

If W, H were independent

$$\begin{aligned} f_{HW}(h, w) &= f_H(h)f_W(w) \\ &= \frac{1}{2\pi\sigma_H\sigma_W} e^{-\frac{1}{2}\left(\frac{(h-\mu_H)^2}{\sigma_H^2} + \frac{(w-\mu_W)^2}{\sigma_W^2}\right)} \end{aligned}$$
$$= ([h \ w] - [\mu_H \ \mu_W]) \begin{bmatrix} \frac{1}{\sigma_H^2} & 0 \\ 0 & \frac{1}{\sigma_W^2} \end{bmatrix} ([h \ w] - [\mu_H \ \mu_W])$$

Covariance matrix!
(Σ)

$$= ([h \ w] - [\mu_H \ \mu_W])^T \begin{bmatrix} \sigma_H^2 & 0 \\ 0 & \sigma_W^2 \end{bmatrix}^{-1} ([h \ w] - [\mu_H \ \mu_W])$$

Notice that since the variables are independent this has to be the diagonal.

DeepLearning.AI

Multivariate Gaussian Distribution: an Example

If W, H were independent

$$\begin{aligned} f_{HW}(h, w) &= f_H(h)f_W(w) \\ &= \frac{1}{2\pi\sigma_H\sigma_W} e^{-\frac{1}{2}\left(\frac{(h-\mu_H)^2}{\sigma_H^2} + \frac{(w-\mu_W)^2}{\sigma_W^2}\right)} \end{aligned}$$

Covariance matrix!
(Σ)

$$= ([h \ w] - [\mu_H \ \mu_W])^T \begin{bmatrix} \sigma_H^2 & 0 \\ 0 & \sigma_W^2 \end{bmatrix}^{-1} ([h \ w] - [\mu_H \ \mu_W])$$

the determinant of the covariance matrix.

DeepLearning.AI

Multivariate Gaussian Distribution: an Example

If W, H were independent

$$\begin{aligned} f_{HW}(h, w) &= f_H(h)f_W(w) \\ &= \frac{1}{2\pi\sigma_H\sigma_W} e^{-\frac{1}{2}\left(\frac{(h-\mu_H)^2}{\sigma_H^2} + \frac{(w-\mu_W)^2}{\sigma_W^2}\right)} \\ &= \frac{1}{2\pi\sigma_H\sigma_W} \exp\left(-\frac{1}{2}([h \ w] - [\mu_H \ \mu_W])^T \begin{bmatrix} \sigma_H^2 & 0 \\ 0 & \sigma_W^2 \end{bmatrix}^{-1} ([h \ w] - [\mu_H \ \mu_W])\right) \end{aligned}$$

Remember that this is valid only if w and h were independent.

DeepLearning.AI

Multivariate Gaussian Distribution: an Example

Dependent case:

$$\begin{aligned} f_{HW}(h, w) &= f_H(h)f_W(w) \\ &= \frac{1}{2\pi\sigma_H\sigma_W} e^{-\frac{1}{2}\left(\frac{(h-\mu_H)^2}{\sigma_H^2} + \frac{(w-\mu_W)^2}{\sigma_W^2}\right)} \\ &= \frac{1}{2\pi\det\Sigma^{1/2}} \exp\left(-\frac{1}{2}([h \ w] - \underline{\mu})^T \Sigma^{-1} ([h \ w] - \underline{\mu})\right) \end{aligned}$$

Well, of course, it's no longer true that the joint

DeepLearning.AI

Multivariate Gaussian Distribution: an Example

Dependent case:

$$f_{HW}(h, w) = f_H(h)f_W(w)$$

$$= \frac{1}{2\pi\sigma_H\sigma_W} e^{-\frac{1}{2}\left(\frac{(h-\mu_H)^2}{\sigma_H^2} + \frac{(w-\mu_W)^2}{\sigma_W^2}\right)}$$

$$= \frac{1}{2\pi\det\Sigma^{1/2}} \exp\left(-\frac{1}{2}\left(\begin{bmatrix} h \\ w \end{bmatrix} - \boldsymbol{\mu}\right)^T \Sigma^{-1} \left(\begin{bmatrix} h \\ w \end{bmatrix} - \boldsymbol{\mu}\right)\right)$$

and now the off-diagonal amounts

correspond to the covariances between variables.

1

$$\Sigma = \begin{bmatrix} \sigma_H^2 & Cov(H, W) \\ Cov(H, W) & \sigma_W^2 \end{bmatrix}$$

2

Multivariate Gaussian Distribution: an Example

Dependent case:

$$f_{HW}(h, w) = f_H(h)f_W(w)$$

$$= \frac{1}{2\pi\sigma_H\sigma_W} e^{-\frac{1}{2}\left(\frac{(h-\mu_H)^2}{\sigma_H^2} + \frac{(w-\mu_W)^2}{\sigma_W^2}\right)}$$

$$= \frac{1}{2\pi\det\Sigma^{1/2}} \exp\left(-\frac{1}{2}\left(\begin{bmatrix} h \\ w \end{bmatrix} - \boldsymbol{\mu}\right)^T \Sigma^{-1} \left(\begin{bmatrix} h \\ w \end{bmatrix} - \boldsymbol{\mu}\right)\right)$$

and now the off-diagonal amounts

correspond to the covariances between variables.

DeepLearning.AI

DeepLearning.AI

Multivariate Gaussian Distribution: General Definition

3

$$f_X(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{1}{2}\frac{(x-\mu)^2}{\sigma^2}}$$

$$f_X(x_1, x_2, \dots, x_n) = \frac{1}{(2\pi)^{n/2} |\Sigma|^{\frac{1}{2}}}$$

covariance matrix
spread of the bell

The bigger the

determinant the bigger the spread.

4

Multivariate Gaussian Distribution: General Definition


$$f_X(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{1}{2}\frac{(x-\mu)^2}{\sigma^2}}$$

$$f_X(x_1, x_2, \dots, x_n) = \frac{1}{(2\pi)^{n/2} |\Sigma|^{\frac{1}{2}}} e^{-\frac{1}{2}(\boldsymbol{x}-\boldsymbol{\mu})^T \Sigma^{-1} (\boldsymbol{x}-\boldsymbol{\mu})}$$

covariance matrix
spread of the bell

Mean vector
 $\boldsymbol{\mu} = [\mu_1, \mu_2, \dots, \mu_n]^T$

Multivariate Gaussian Distribution: General Definition



DeepLearning.AI

Gaussian Distribution
Multivariate:

M and # (Σ) are vectors.

$$f_X(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{1}{2} \frac{(x-\mu)^2}{\sigma^2}}$$

so $|\Sigma|$ determinant of the covariance matrix

$$\text{Whose } x = [x_1, x_2, \dots, x_n]^T$$

$$\mu = [\mu_1, \mu_2, \dots, \mu_n]^T$$

SOURCE: Coursene Probability & Statistics for Data Science
& Machine Learning!