

Quantum Mechanics Primer for Nuclear Astrophysics

Note for NS 3035

Dehiwalage Don Dilruwan
Colombo

“A ship in harbor is safe, but that is not what ships are built for” — John A. Shedd

*In the first three classes we discussed the foundations of a simple stellar model. We began with non rotating uniform stars and developed several important concepts such as the free fall timescale, hydrostatic equilibrium, and the virial theorem (I encourage you to practice these derivations on your own). Using kinetic theory, we realized that most ordinary stars exist in a **non relativistic** regime where the thermal motion of particles dominates the internal physics.¹ In Classes 3 and 4, we stepped back to look at the bigger astrophysical picture. Our prime question was how stars are formed ? We discussed the evolution from the interstellar medium (ISM) to giant molecular clouds (GMCs), the formation of open clusters and the importance of the Jeans criterion in initiating gravitational collapse. We did not want to remain at a surface level discussion by simply looking at pictures and memorizing a few fancy names. Those kinds of courses are for toddlers!. As science students, our goal is to develop deeper physical intuition even when the concepts initially feel uncomfortable.*

From this week onward, we will pause our astronomy discussions and move toward the more familiar territory of nuclear science. I understand that the first few classes may have felt somewhat overwhelming, but I believe you will become much more comfortable from this point onward. However, even if some of the coming weeks feel like Greek to you, I invite you to reflect on the quote placed at the beginning of this document, The best learning often begins just beyond our comfort zone. It means that, truly understanding this subject is rarely comfortable at the beginning. Derivations may feel messy, and sometimes the mathematics may appear intimidating. But that discomfort is often a sign that real learning is taking place. So as we move deeper into nuclear astrophysics hang on tight and prepare for rough seas ahead. We are leaving the calm harbor and sailing into open waters !!

¹We also briefly mentioned the ultra relativistic regime where electron degeneracy pressure becomes important

Contents

Quantum Grammar	4
1 The Wave Function and Probability Density	5
2 The Schrödinger Equation	7
2.1 From Classical Energy to Quantum Mechanics	7
2.2 The Time Dependent Schrödinger Equation	7
2.3 What Does the Wave Function Physically Mean?	8
2.4 Time Independent Schrödinger Equation	8
2.4.1 Why Quantization Happens	9
2.5 Why the Schrödinger Equation Matters for Nuclear Astrophysics	9
2.5.1 Quantum Tunneling Inside the Sun	9
3 Spherical Coordinates and Angular Momentum	11
3.1 Why Spherical Coordinates Matter in Quantum Mechanics	11
3.2 Spherical Symmetry and the Schrödinger Equation	12
3.3 Radial Motion and Angular Motion	12
3.4 Angular Momentum Quantization	12
3.4.1 What is an Operator?	13
3.4.2 Angular Momentum Operators	13
3.5 The Centrifugal Barrier	14
3.5.1 Importance of the Centrifugal Barrier in Nuclear Astrophysics	14
4 Applications of Quantum Mechanics in Nuclear Astrophysics	15
4.1 The Nuclear Shell Model	16
4.1.1 Spin Orbit Coupling and Magic Numbers	17
4.2 Cross Sections and Reaction Probabilities	17
4.2.1 Transition Amplitudes and Reaction Probabilities	17
4.2.2 Cross Sections in Nuclear Astrophysics	18
4.2.3 Reaction Rates in Stars	18
4.3 Inverse Reactions and Detailed Balance	18
4.3.1 Quantum Mechanical Interpretation	19
4.3.2 Detailed Balance	19
4.3.3 Inverse Reactions in Stellar Environments	19
4.3.4 Reaction Equilibrium and Nuclear Statistical Equilibrium	20
4.4 Resonant and Non Resonant Reactions	20
4.4.1 Non Resonant Reactions	20
4.4.2 Resonant Reactions	21
4.4.3 Breit Wigner Resonance Formula	21
4.4.4 Comparison Between Resonant and Non Resonant Reactions	21
4.4.5 Importance in Nuclear Astrophysics	22

Quantum Grammar

In nuclear astrophysics we study nuclear reactions inside stars. The stellar plasma in ordinary star exists in non relativistic regime where nuclear reactions are driven by the thermal motion of particles. Inside stellar cores nuclei move continuously because of the extremely high temperatures. As these nuclei move, they frequently collide with one another. Although we developed kinetic theory of gases and the virial theorem using classical treatment, classical physics alone cannot explain how these reactions actually occur.

In nuclear scale, positively charged nuclei strongly repel each other through the coulomb force. Classically, the thermal energies inside stars are often too small to overcome this Coulomb barrier. In fact, if stars relied purely on classical physics, thermonuclear fusion inside the Sun would be impossible. This is where quantum mechanics becomes essential in nuclear astrophysics. At nuclear scales, particles no longer behave purely as localized classical objects. Instead, they also exhibit wave like behavior described by the de Broglie wavelength,

$$\lambda = \frac{h}{p}$$

where h is Planck's constant and p is the particle momentum. Here is the mathematical and conceptual breakdown of the core quantum foundations you need for the success in Nuclear Astrophysics (NS 3035)

Chapter 1

The Wave Function and Probability Density

One of the most fundamental and crude ways to understand quantum physics is to compare it with classical physics. In the classical world a particle possesses a precise position and momentum at any given time. In the quantum world, a particle such as a proton or neutron does not have an exactly defined position and momentum simultaneously. *Instead, the particle is completely described by a wave function* denoted by the Greek letter $\Psi(\vec{r}, t)$, where $\vec{r}$ represents the position vector and t represents time. The wave function itself isn't directly observable but its absolute square represents a probability density. The probability P of finding a particle within a specific volume V is given by

$$P = \int_V |\Psi(\vec{r}, t)|^2 dV$$

where

$$|\Psi(\vec{r}, t)|^2$$

is called the probability density. This probabilistic interpretation is one of the key differences between classical and quantum physics. Instead of predicting an exact trajectory for a particle, quantum mechanics predicts the probability of observing the particle at different positions. This wave-like nature of matter becomes extremely important at nuclear scales and plays a central role in phenomena such as quantum tunneling, which allows nuclear fusion to occur inside stars.

Why it matters for Nuclear Astrophysics ?? The probability density becomes especially important because nuclear reactions inside stars depend strongly on the likelihood of particles being sufficiently close to one another. Since nuclei behave as quantum mechanical waves their wave functions can partially penetrate the Coulomb barrier even when their classical kinetic energies are too small to overcome it. The probability density near or inside the barrier determines the likelihood of tunneling and therefore the probability of fusion occurring.

As a result, quantities such as reaction **cross sections**, **tunneling probabilities**, and **thermonuclear reaction rates** are all fundamentally connected to the behavior of the wave function and its probability density. Without this quantum mechanical description, you are blind when you explain how stars like the Sun sustain nuclear fusion and continuously generate energy over billions of years.

Interactive Python Experiment : Gaussian Wave Packet and Probability Density

To better understand the connection between the quantum wave function, probability density, and stellar fusion processes, an interactive Python simulation was developed using Google Colab. The experiment visualizes the Gaussian wave packet,

$$\psi(x) = e^{-\frac{(x-x_0)^2}{2\sigma^2}} e^{ikx}$$

and its probability density

$$|\psi(x)|^2$$

allowing students to directly observe how the spatial spread, localization, and wave oscillations change as the parameters x_0 , σ , and k are modified.




Figure 1.1: Gaussian wave packet showing the real part, imaginary part, and probability density $|\psi(x)|^2$. The probability density determines the likelihood of finding the particle at a given position and forms the quantum foundation for tunneling and thermonuclear fusion inside stars.

This simulation provides an intuitive computational introduction to:

- Quantum wave functions
- Probability density
- de Broglie wave behavior
- Localization and uncertainty
- Quantum tunneling
- Nuclear fusion probability inside stars

Interactive Experiment Link: <https://colab.research.google.com/drive/1w150mJ31JNgn0Xe6wpYLF9w7NPH-mHe5?usp=sharing>

Chapter 2

The Schrödinger Equation

One of the biggest questions in quantum mechanics is the following: *If particles behave like waves, how do these waves evolve with time?* In classical physics, the motion of an object is determined using Newton's laws. If we know the force acting on a particle, we can calculate its trajectory, velocity, and future position. However, at atomic and nuclear scales this picture begins to fail. Electrons, protons, neutrons and entire nuclei exhibit wave like behavior. Therefore instead of tracking precise trajectories, quantum mechanics describes particles using a wave function

$$\Psi(\vec{r}, t)$$

which contains the complete probabilistic description of the particle. The equation that determines how this wave function evolves is called the **Schrödinger Equation**. In many ways, the Schrödinger Equation plays a role in quantum mechanics similar to Newton's second law in classical mechanics. It tells us how a quantum system changes with time and how particles behave under different physical conditions

2.1 From Classical Energy to Quantum Mechanics

To understand where the Schrödinger Equation comes from, we begin with a familiar idea from classical mechanics. For a particle with mass m , the total energy is

$$E = \frac{p^2}{2m} + V$$

where

- E is the total energy
- p is the momentum
- V is the potential energy

This equation simply states that the total energy is the sum of kinetic energy and potential energy. Quantum mechanics introduces a new idea: particles behave like waves. According to de Broglie, a particle with momentum p possesses a wavelength

$$\lambda = \frac{h}{p}$$

This means that matter itself has wave-like properties. As a result, quantities such as momentum and energy become connected to wave behavior. The Schrödinger Equation was constructed to describe how these matter waves evolve and interact.

2.2 The Time Dependent Schrödinger Equation

The most general form of the Schrödinger Equation is the Time Dependent Schrödinger Equation (TDSE)

$$i\hbar \frac{\partial \Psi(\vec{r}, t)}{\partial t} = -\frac{\hbar^2}{2m} \nabla^2 \Psi(\vec{r}, t) + V(\vec{r}, t) \Psi(\vec{r}, t)$$

At first glance this equation may look intimidating, but each term has a very clear physical meaning.

Left Hand Side: Time Evolution

$$i\hbar \frac{\partial \Psi}{\partial t}$$

This term describes how the wave function changes with time. It tells us how the quantum state evolves as the system moves forward in time.

First Term on the Right: Kinetic Energy

$$-\frac{\hbar^2}{2m} \nabla^2 \Psi$$

This term is related to the kinetic energy of the particle. The operator

$$\nabla^2$$

called the Laplacian measures how strongly the wave function bends or curves in space. Large curvature corresponds to rapidly varying waves, which are associated with larger momentum and kinetic energy.

Second Term on the Right : Potential Energy

$$V(\vec{r}, t) \Psi$$

This term represents the influence of external forces or interactions acting on the particle. For example

- electromagnetic forces,
- nuclear forces,
- gravitational potentials,
- Coulomb barriers between nuclei.

Together, these terms determine how the quantum wave evolves in space and time.

2.3 What Does the Wave Function Physically Mean?

The wave function itself is not directly observable. Instead, measurable quantities come from its probability density

$$|\Psi(\vec{r}, t)|^2$$

which gives the probability of finding the particle at a particular position. The total probability must always equal one

$$\int |\Psi(\vec{r}, t)|^2 dV = 1$$

because the particle must exist somewhere in space. This condition is known as **normalization**.

2.4 Time Independent Schrödinger Equation

In many physical systems, the potential energy does not change with time. For example

- electrons bound inside atoms,
- nucleons inside nuclei,
- particles trapped in stable potentials.

In such cases, the wave function can be separated into spatial and temporal parts. This leads to the Time Independent Schrödinger Equation (TISE)

$$-\frac{\hbar^2}{2m} \nabla^2 \psi(\vec{r}) + V(\vec{r}) \psi(\vec{r}) = E \psi(\vec{r})$$

This equation determines the allowed quantum states and energies of the system. Unlike classical systems, quantum systems do not possess arbitrary energies. Instead, only specific discrete energy values are allowed. This phenomenon is known as **energy quantization**.

2.4.1 Why Quantization Happens

One intuitive way to understand quantization is to compare quantum waves with standing waves on a string. Only certain wave patterns can fit on a stretched string without destroying themselves through destructive interference. Similarly, only certain wave functions can exist stably inside atoms or nuclei. As a result, only specific energy states are physically allowed.

This idea forms the foundation of

- atomic structure
- nuclear shell structure
- nuclear excited states
- resonance reactions in stars

2.5 Why the Schrödinger Equation Matters for Nuclear Astrophysics


The Schrödinger Equation is one of the most important equations in nuclear astrophysics because stellar environments are fundamentally quantum mechanical. Many important astrophysical processes depend directly on wave behavior and probability densities, including:

- Quantum tunneling
- Nuclear fusion
- Resonance reactions
- Neutron capture
- Nuclear scattering
- Beta decay
- Nuclear shell structure

Without quantum mechanics, most fusion reactions inside stars would be impossible

2.5.1 Quantum Tunneling Inside the Sun

One of the most remarkable predictions of the Schrödinger Equation is **quantum tunneling**. The dominant fusion process inside the Sun begins with the proton-proton reaction



where two protons combine to form deuterium. However, protons are both positively charged. As they approach each other they experience strong electromagnetic repulsion known as the Coulomb barrier. The Coulomb potential is approximately

$$V(r) = \frac{1}{4\pi\epsilon_0} \frac{e^2}{r}$$

Inside the solar core, the kinetic energy of protons is usually much smaller than this barrier energy.

$$E < V(r)$$

Classically, fusion should therefore be impossible. However, the Schrödinger Equation predicts that the wave function does not abruptly stop at the barrier. Instead, the wave penetrates partially into the classically forbidden region. Inside the barrier, the solution behaves approximately like

$$\psi(r) \propto e^{-\kappa r}$$

This means the wave function decreases exponentially but never becomes exactly zero. Because the probability density is

$$|\psi(r)|^2$$

there remains a finite probability that the protons can appear on the opposite side of the barrier. Once the protons become sufficiently close, the strong nuclear force dominates and fusion can occur. Therefore stars shine because quantum mechanics allows particles to tunnel through barriers that classical physics cannot overcome. This quantum tunneling mechanism is one of the fundamental reasons why thermonuclear fusion inside stars is possible.

Chapter 3

Spherical Coordinates and Angular Momentum

Many important systems in nuclear astrophysics exhibit approximate spherical symmetry (We briefly discussed this idea in the first class), for example

- atomic nuclei are roughly spherical
- stars are approximately spherical
- many nuclear interaction potentials depend mainly on distance from the center.

Because of this symmetry, it is often more natural to describe quantum systems using spherical coordinates rather than Cartesian coordinates. In ordinary Cartesian coordinates, the position of a particle is written as

$$(x, y, z)$$

However for systems involving circular or spherical motion, Cartesian coordinates can make the mathematics unnecessarily complicated. Instead, we use spherical coordinates

$$(r, \theta, \phi)$$

where

- r is the radial distance from the origin
- θ is the polar angle
- ϕ is the azimuthal angle

Physically, spherical coordinates separate motion into two independent parts

1. Motion toward or away from the center
2. Motion around the center

This turns out to be extremely useful in quantum mechanics because many nuclear systems naturally behave this way

3.1 Why Spherical Coordinates Matter in Quantum Mechanics

In quantum mechanics, particles are described using wave functions. For a particle moving in three dimensions, the wave function depends on position

$$\psi(x, y, z)$$

When spherical symmetry is present, it becomes more natural to write the wave function as

$$\psi(r, \theta, \phi)$$

This allows the wave function to separate naturally into

- a radial part
- an angular part

Mathematically, we write

$$\psi(r, \theta, \phi) = R(r) Y(\theta, \phi)$$

where

- $R(r)$ describes how the wave behaves with distance
- $Y(\theta, \phi)$ describes how the wave behaves around the sphere

This idea is extremely important because it transforms one complicated three dimensional problem into two simpler problems.

3.2 Spherical Symmetry and the Schrödinger Equation

Many nuclear potentials depend only on distance from the center

$$V(\vec{r}) \rightarrow V(r)$$

This means the potential does not care about direction. It only depends on how far the particle is from the center. Because of this symmetry, the Schrödinger Equation becomes separable in spherical coordinates. The Time Independent Schrödinger Equation becomes

$$-\frac{\hbar^2}{2m} \nabla^2 \psi(r, \theta, \phi) + V(r) \psi(r, \theta, \phi) = E \psi(r, \theta, \phi)$$

The Laplacian operator in spherical coordinates is

$$\nabla^2 = \frac{1}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial}{\partial r} \right) + \frac{1}{r^2 \sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial}{\partial \theta} \right) + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2}{\partial \phi^2}$$

At first this expression may look complicated, but something important should be noticed

- the first term depends only on radius r
- the remaining terms depend only on angles (θ, ϕ)

This strongly suggests that the wave function itself can also be separated into radial and angular components. This procedure is called **separation of variables**

3.3 Radial Motion and Angular Motion

After separation, quantum motion inside a spherical system can be understood as a combination of

- radial motion
- angular motion

Radial Motion The radial part describes how the particle moves toward or away from the center. For example

- approaching the nucleus
- moving away from the nucleus
- tunneling through barriers

Angular Motion The angular part describes rotational or orbital behavior around the center. This introduces the concept of angular momentum.

3.4 Angular Momentum Quantization

In classical mechanics, angular momentum can take any value. However, quantum mechanics predicts something very different. At atomic and nuclear scales, angular momentum is **quantized**. This means particles are only allowed to possess specific discrete angular momentum values rather than arbitrary continuous values. Before introducing the equations we first need to understand an important idea in quantum mechanics called an **operator**

3.4.1 What is an Operator?

In classical physics, physical quantities such as position, momentum, and energy are represented simply by numbers. In quantum mechanics, physical observables are represented using mathematical objects called **operators**. An operator acts on a wave function and extracts physical information about the system. For example:

- the energy operator measures energy
- the momentum operator measures momentum
- the angular momentum operator measures angular momentum

Operators are usually written with a small hat symbol

^

placed above the quantity. For example,

$$\hat{L}$$

represents the angular momentum operator. Physically, this means that angular momentum is no longer treated as an ordinary classical number. Instead, it becomes a quantum mechanical observable acting on the wave function.

3.4.2 Angular Momentum Operators

Solving the angular part of the Schrödinger Equation introduces the angular momentum operators

$$\hat{L}^2 Y_{lm} = l(l+1)\hbar^2 Y_{lm}$$

and

$$\hat{L}_z Y_{lm} = m\hbar Y_{lm}$$

where

- l is the orbital angular momentum quantum number
- m is the magnetic quantum number

The allowed values are

$$l = 0, 1, 2, 3, \dots$$

and

$$-l \leq m \leq l$$

Physically, the quantum number l determines the orbital shape of the wave function.

$$l = 0 \rightarrow s\text{-wave}$$

$$l = 1 \rightarrow p\text{-wave}$$

$$l = 2 \rightarrow d\text{-wave}$$

$$l = 3 \rightarrow f\text{-wave}$$

These orbital patterns become extremely important in atomic structure, nuclear structure, and reaction probabilities.

3.5 The Centrifugal Barrier

After separating the Schrödinger Equation, the radial equation contains an additional term called the **centrifugal barrier**. The radial equation becomes

$$\left[-\frac{\hbar^2}{2m} \frac{d^2}{dr^2} + V_{\text{eff}}(r) \right] u(r) = Eu(r)$$

where

$$u(r) = rR(r)$$

and the effective potential is

$$V_{\text{eff}}(r) = V(r) + \frac{l(l+1)\hbar^2}{2mr^2}$$

The second term

$$\frac{l(l+1)\hbar^2}{2mr^2}$$

is the centrifugal barrier.

Physically, this term acts like an additional repulsive barrier for particles carrying angular momentum. A useful classical analogy is circular motion. In classical physics, an object rotating around a center resists falling directly inward because of rotational motion. Similarly in quantum mechanics, particles with larger angular momentum experience a stronger effective barrier near the center. Therefore:

- particles with $l = 0$ can approach the center more easily
- particles with larger l values are increasingly suppressed near small radii

3.5.1 Importance of the Centrifugal Barrier in Nuclear Astrophysics

The centrifugal barrier is a major concept in stellar nuclear reactions. Inside stars, nuclei already struggle against the Coulomb barrier caused by electromagnetic repulsion. If the reacting particles also carry angular momentum they must overcome an additional centrifugal barrier. Therefore fusion reactions involving larger angular momentum become much less probable. This means:

- s -wave reactions ($l = 0$) occur most easily
- p -wave and d -wave reactions are more strongly suppressed

This angular momentum dependence strongly affects

- stellar fusion rates
- neutron capture probabilities
- resonance reactions
- nucleosynthesis pathways

Chapter 4

Applications of Quantum Mechanics in Nuclear Astrophysics

Now that we have introduced the mathematical foundations of quantum mechanics, such as the Schrödinger equation, spherical coordinates, angular momentum, and quantized wave functions, we can begin applying these ideas to real nuclear astrophysics systems. If you already have a background in basic of quantum mechanics, you may not need to study Chapters 1, 2, and 3 in detail. However, I strongly encourage you to read this chapter carefully in order to develop a better physical understanding of how quantum mechanics is applied within nuclear astrophysics.

Quantum States and Bra Ket Notation

In quantum mechanics, the complete state of a physical system is represented using a mathematical object called a **state vector**. These states are commonly written using **Dirac bra ket notation**. A quantum state is written as

$$|\psi\rangle$$

and is called a **ket**. The corresponding conjugate state is written as

$$\langle\psi|$$

and is called a **bra**. Together, bras and kets provide a compact way to describe quantum mechanical probabilities and transitions. For nuclear reactions we often define

- the initial state $|i\rangle$
- the final state $|f\rangle$

For example, in a reaction

$$p + p \rightarrow d + e^+ + \nu_e$$

the two incoming protons correspond to the initial quantum state $|i\rangle$, while the deuterium and emitted particles correspond to the final state $|f\rangle$.

The Hamiltonian Operator

In quantum mechanics, physical observables are represented using operators. One of the most important operators is the **Hamiltonian operator**

$$\hat{H}$$

which represents the total energy of the system. The Hamiltonian contains

- kinetic energy terms,
- potential energy terms,
- interaction terms responsible for nuclear reactions.

The Schrödinger Equation can be written compactly using the Hamiltonian operator as

$$\hat{H}|\psi\rangle = E|\psi\rangle$$

which states that the Hamiltonian acting on a quantum state produces the energy of that state. Physically, the Hamiltonian determines how quantum systems evolve and interact.

4.1 The Nuclear Shell Model

One of the most important applications of quantum mechanics in nuclear astrophysics is the nuclear shell model. The model assumes that nucleons occupy discrete quantum states inside the nucleus, similar to electrons in atoms. The average nuclear potential is first approximated using the three-dimensional harmonic oscillator potential $V(r) = \frac{1}{2}m\omega^2 r^2$, where m is the nucleon mass and ω is the oscillator frequency. Substituting this potential into the Schrödinger equation $-\frac{\hbar^2}{2m}\nabla^2\psi + V(r)\psi = E\psi$ produces quantized energy levels

$$E_N = \left(N + \frac{3}{2}\right) \hbar\omega$$

where the principal oscillator quantum number is $N = 2(n-1) + l$. Here $n = 1, 2, 3, \dots$ is the radial quantum number and $l = 0, 1, 2, \dots$ is the orbital angular momentum quantum number. Each shell contains several magnetic substates. For a given l , the number of substates is $2l + 1$ since $-l \leq m \leq l$. Including nucleon spin introduces an additional factor of 2, giving the total number of nucleons in a shell $2(2l + 1)$

First Shell: $N = 0$

For $N = 0$, the only allowed value is $l = 0$ corresponding to the s -state. The number of nucleons becomes

$$2(2l + 1) = 2(1) = 2$$

Thus the first shell closes at

$$2$$

Second Shell: $N = 1$

For $N = 1$, the allowed value is $l = 1$ corresponding to the p -state. The number of states is

$$2(2(1) + 1) = 6$$

Adding to the previous shell gives

$$2 + 6 = 8$$

Thus the second shell closes at

$$8$$

Third Shell: $N = 2$

For $N = 2$, the allowed angular momentum values are $l = 0, 2$, corresponding to one s -state and one d -state. The number of states becomes

$$2(1) + 2(5) = 12$$

Adding to previous shells gives

$$8 + 12 = 20$$

Thus the third shell closes at

$$20$$

Therefore, the harmonic oscillator shell model predicts the primitive magic numbers

$$2, 8, 20$$

which successfully explain many lighter nuclei. Nuclei with filled shells possess enhanced stability similar to noble gases in atomic physics.

4.1.1 Spin Orbit Coupling and Magic Numbers

Although the harmonic oscillator model explains lighter nuclei, it fails to reproduce heavier magic numbers such as

$$28, 50, 82, 126$$

The missing ingredient is spin-orbit coupling. Nucleons possess both orbital angular momentum $\vec{L}$ and intrinsic spin angular momentum $\vec{S}$. Their interaction behaves approximately as

$$\hat{H}_{ls} \propto \vec{L} \cdot \vec{S}$$

The total angular momentum becomes $\vec{J} = \vec{L} + \vec{S}$. For nucleons with $s = \frac{1}{2}$, the allowed total angular momentum values are

$$j = l + \frac{1}{2}$$

and

$$j = l - \frac{1}{2}$$

These states possess slightly different energies, splitting the shell structure and shifting some levels downward. This produces the experimentally observed magic numbers

$$2, 8, 20, 28, 50, 82, 126$$

These magic nuclei are extremely important in nuclear astrophysics because their enhanced stability strongly influences neutron capture processes, stellar nucleosynthesis, beta decay pathways, supernova element formation, and r-process abundance peaks.

4.2 Cross Sections and Reaction Probabilities

One of the central goals of nuclear astrophysics is to determine the probability that a nuclear reaction will occur inside stars. In classical physics, particles are often imagined as tiny spheres that collide directly with one another. However, quantum mechanics provides a very different picture. In quantum mechanics, nuclei behave as wave packets rather than perfectly localized particles. Nuclear reactions therefore occur through interactions between quantum mechanical wave functions. The quantity used to measure the probability of a nuclear reaction is called the *cross section*, denoted by σ . Physically, the cross section represents an effective interaction area for the reaction. A larger cross section corresponds to a higher probability that the reaction occurs. However unlike classical collision theory (kinetic theory) quantum mechanical cross sections are determined by wave function overlap, tunneling probabilities, angular momentum barriers and transition amplitudes between quantum states.

4.2.1 Transition Amplitudes and Reaction Probabilities

In nuclear reactions, the probability that the system transitions from an initial state $|i\rangle$ to a final state $|f\rangle$ depends on the interaction Hamiltonian. The transition amplitude is written as

$$\langle f | \hat{H} | i \rangle$$

This quantity measures how strongly the initial and final wave functions overlap through the interaction. The reaction probability is proportional to the square of this amplitude

$$P_{i \rightarrow f} \propto |\langle f | \hat{H} | i \rangle|^2$$

This equation is extremely important in nuclear astrophysics because it shows that reactions are fundamentally quantum mechanical transitions between allowed states. If the overlap between the wave functions is large, the reaction probability becomes large. If the overlap is weak, the reaction becomes suppressed.

4.2.2 Cross Sections in Nuclear Astrophysics

The nuclear cross section is directly related to the transition probability. Quantum mechanically, the cross section depends on

- wave function overlap
- tunneling probability
- resonance enhancement
- angular momentum selection rules
- conservation laws
- available phase space

For stellar fusion reactions, tunneling through the Coulomb barrier plays a dominant role. Even if nuclei possess insufficient classical energy to overcome the barrier, their wave functions can partially penetrate the forbidden region, giving a finite reaction probability. This is why fusion reactions inside stars are possible despite the strong electromagnetic repulsion between nuclei.

4.2.3 Reaction Rates in Stars

Inside stars, nuclei possess a distribution of velocities described approximately by the Maxwell Boltzmann distribution. The reaction rate per unit volume between two nuclear species is

$$R = n_1 n_2 \langle \sigma v \rangle$$


where

- n_1 and n_2 are the number densities,
- σ is the reaction cross section,
- v is the relative velocity,
- $\langle \sigma v \rangle$ is the thermally averaged reaction rate.


This quantity is one of the most important parameters in stellar evolution calculations because it determines how rapidly nuclear reactions proceed inside stars. Therefore, the evolution of stars, nucleosynthesis pathways, and elemental abundances throughout the Universe are ultimately controlled by quantum mechanical transition probabilities and nuclear cross sections.

4.3 Inverse Reactions and Detailed Balance

In nuclear astrophysics, many nuclear reactions can proceed in both forward and reverse directions. If a reaction of the form



is possible, then the inverse reaction



may also occur under suitable physical conditions. Classically, one might imagine these reactions simply as ordinary collisions occurring in opposite directions. However, quantum mechanics provides a much deeper interpretation. Since nuclei behave as quantum mechanical waves, nuclear reactions are fundamentally described as transitions between quantum states. The probability that a reaction occurs depends on the overlap between the initial and final wave functions, as well as conservation laws such as energy, angular momentum, parity, and spin.

4.3.1 Quantum Mechanical Interpretation

In quantum mechanics, a nuclear reaction is treated as a transition between two quantum states

$$|i\rangle \rightarrow |f\rangle$$

where $|i\rangle$ represents the initial state and $|f\rangle$ represents the final state. The probability amplitude for this transition is determined by the matrix element

$$\langle f | \hat{H} | i \rangle$$


where $\hat{H}$ is the interaction Hamiltonian responsible for the reaction. The reaction probability is proportional to the square of this amplitude,

$$P_{i \rightarrow f} \propto |\langle f | \hat{H} | i \rangle|^2$$

which means that nuclear reactions are fundamentally controlled by wave function overlap and quantum transition probabilities. If the wave functions overlap strongly, the reaction probability becomes large. If the overlap is weak, the reaction becomes strongly suppressed. This is one of the reasons why tunneling probabilities, angular momentum barriers, and resonance structures are so important in nuclear astrophysics.

4.3.2 Detailed Balance

One of the most important consequences of quantum mechanics is that forward and inverse reactions are not independent. They are related through the principle of *detailed balance*. At thermal equilibrium, the forward and reverse reaction rates must balance



This occurs because microscopic quantum processes are reversible. If quantum mechanics allows a transition from $|i\rangle \rightarrow |f\rangle$, then the reverse transition $|f\rangle \rightarrow |i\rangle$ is also allowed. The ratio between forward and reverse reaction rates depends on

- phase space availability
- statistical degeneracy
- thermal population
- reaction Q -value
- spin multiplicities

For reactions in thermal equilibrium, detailed balance relates the cross sections approximately through

$$\frac{\sigma_{f \rightarrow i}}{\sigma_{i \rightarrow f}} \propto \frac{g_i p_i^2}{g_f p_f^2}$$


where

- $g = 2J + 1$ is the spin degeneracy factor
- p is the particle momentum
- J is the total angular momentum quantum number


This equation shows that forward and inverse reactions are fundamentally connected through quantum statistics and conservation laws

4.3.3 Inverse Reactions in Stellar Environments

In stars, inverse reactions become extremely important at high temperatures because photons and particles possess enough thermal energy to break nuclei apart. For example, radiative capture reactions such as



can proceed in reverse through **photodisintegration**



Similarly, alpha capture reactions may be balanced by alpha emission reactions at sufficiently high temperatures. Quantum mechanically, the same compound nuclear states that enhance forward reactions may also enhance inverse reactions. Therefore, resonance structures strongly affect both directions simultaneously. The competition between forward fusion and inverse photodisintegration strongly influences stellar nucleosynthesis pathways. In extremely hot astrophysical environments such as

- supernova explosions
- neutron star mergers
- X-ray bursts
- late stage stellar burning

inverse reactions can become comparable to or even dominate forward reactions.

4.3.4 Reaction Equilibrium and Nuclear Statistical Equilibrium

When forward and inverse reactions occur rapidly compared to stellar evolution timescales, the system may reach *nuclear statistical equilibrium (NSE)*. In this NSE regime, nuclear abundances are determined not by individual reactions alone, but by the balance between all forward and reverse quantum transitions. Under these conditions, nuclei continuously form and break apart while maintaining an overall statistical equilibrium distribution. This equilibrium depends strongly on

- temperature
- density
- binding energy
- nuclear partition functions
- quantum degeneracy

Therefore inverse reactions are not simply *reverse collisions*. They are fundamentally connected to forward reactions through quantum mechanical transition probabilities, resonance structures, conservation laws and thermal equilibrium. Without these quantum mechanical effects, it would be impossible to accurately describe stellar nucleosynthesis and the chemical evolution of the Stars and Universe.

4.4 Resonant and Non Resonant Reactions

In nuclear astrophysics, nuclear reactions do not always occur in the same way. Some reactions proceed smoothly over a wide range of energies, while others become enormously enhanced at very specific energies. Quantum mechanics explains this behavior naturally through the wave nature of nuclei and the existence of discrete nuclear energy levels. Nuclear reactions are generally classified into two categories

1. Non resonant reactions
2. Resonant reactions

4.4.1 Non Resonant Reactions

A *non resonant reaction* occurs when the interacting nuclei fuse directly without forming a long lived intermediate excited state. In this case, the reaction probability changes relatively smoothly with energy. Quantum mechanically, the incoming nuclear wave function tunnels through the Coulomb barrier and directly overlaps with the final nuclear state. The reaction therefore depends mainly on

- tunneling probability
- wave function overlap
- Coulomb barrier penetration
- angular momentum barriers

The cross section for non resonant reactions usually changes gradually with energy because there is no special energy level strongly enhancing the reaction. A useful physical picture is to imagine two nuclei approaching each other and immediately reacting once they tunnel sufficiently close together. The reaction does not temporarily “pause” inside an excited compound state. Many proton proton fusion reactions inside stars behave approximately as non resonant tunneling reactions.

4.4.2 Resonant Reactions

A *resonant reaction* occurs when the interacting nuclei temporarily form an excited compound nucleus with an energy very close to one of the allowed quantum energy levels of the system. Quantum mechanically, the incoming nuclear wave function matches a quasi stable standing wave configuration inside the nucleus. This dramatically increases the probability of interaction. The process can be written schematically as



where

- C^* is an excited compound nucleus
- the intermediate state exists temporarily before decaying

Resonance occurs when the incoming energy approximately matches an allowed excited energy level

$$E \approx E_r$$

where E_r is the resonance energy. At this energy, the nuclear wave functions interfere constructively and produce a large enhancement in the reaction probability. This is similar to resonance in classical systems such as

- pushing a swing at the correct frequency
- acoustic resonance in musical instruments
- standing waves on a string

In nuclear physics, resonance appears because nuclei possess discrete quantized excited states. If the incoming particles supply exactly the correct energy, the compound nucleus can temporarily occupy one of these states

4.4.3 Breit Wigner Resonance Formula

Near a resonance energy, the reaction cross section approximately follows the Breit Wigner formula

$$\sigma(E) \propto \frac{\Gamma^2/4}{(E - E_r)^2 + \Gamma^2/4}$$

where

- E_r is the resonance energy
- Γ is the resonance width

The resonance width Γ is related to the lifetime of the excited state through the uncertainty principle

$$\Gamma \sim \frac{\hbar}{\tau}$$

where τ is the lifetime of the resonance. Therefore

- narrow resonances correspond to longer lived states
- broad resonances correspond to short lived states

4.4.4 Comparison Between Resonant and Non Resonant Reactions

Non Resonant Reactions goes

- direct tunneling process
- smooth energy dependence
- no intermediate quasi stable state
- reaction probability mainly controlled by barrier penetration

Resonant Reactions goes

- temporary compound nucleus formation,
- strong enhancement at specific energies,
- associated with quantized excited states,
- reaction probability dominated by resonance matching.

4.4.5 Importance in Nuclear Astrophysics

Resonant and non resonant reactions play central roles in stellar nucleosynthesis. Many stellar fusion reactions occur extremely slowly because of Coulomb barrier suppression. However, even a narrow resonance can dramatically increase the reaction rate. Some of the most important reactions in astrophysics are resonance dominated, including reactions involved in

- carbon production
- oxygen production
- neon burning
- silicon burning
- heavy element nucleosynthesis

A famous example is the Hoyle state in carbon-12. Without this resonance, the triple alpha process inside stars would be far less efficient, and the cosmic abundance of carbon would be dramatically smaller.