
ACR Classification (ACR+ vs ACR-)
Model: set_mil_mt_no_sab [BACC 0.623 +/- 0.034] | Cluster attribution Δ affinity (mean +/- s.d., 5 splits)
HE [top 14 clusters]
<- ACR- | ACR+ ->
Alveolar with hemorrhage and inflammation
Alveolar with empty spaces
Alveolar with empty spaces
Alveolar
Alveolar with hemorrhage and inflammation
Alveolar with hemorrhage and inflammation
Bronchial
Alveolar with empty spaces
Alveolar with hemorrhage and inflammation
Alveolar with hemorrhage and inflammation
Alveolar with hemorrhage and inflammation
Alveolar with empty spaces
Alveolar
Alveolar
Enriched in ACR+
Enriched in ACR-
0
2
4
6
8
10
12
Delta cluster affinity
(high-risk minus low-risk)
BAL [top 14 clusters]
<- ACR- | ACR+ ->
Proliferating macrophages
TRAM-7
Perivascular macrophages
CD4 T naive cells
CD4 T cells-3
MoAM-3
MoAM-4
TRAM-5
TRAM-8
CD4 T cells-2
TRAM-3
TRAM-6
TRAM-4
CD8 T cells-2
n=5 splits
0
2
4
6
8
10
12
Delta cluster affinity
(high-risk minus low-risk)
CT [top 14 clusters]
<- ACR- | ACR+ ->
13
27
1
11
5
9
17
18
28
22
0
7
8
6
n=5 splits
0
2
4
6
8
10
12
Delta cluster affinity
(high-risk minus low-risk)