Northbridge General Hospital (fictional)

100 Example Avenue, Northbridge - fictional address, telephone 000 0000

Endocrinology Outpatient Consultation

Department of Endocrinology & Diabetes

PATIENT **Rajesh Menon**	UHID **MG25001**	AGE / SEX **62 years / Male**
REPORT DATE **15 Sep 2025**	REFERENCE NO. **CON-80889773**	DEPARTMENT **Department of Endocrinology & Diabetes**

Reason for Visit

Routine four-monthly review of type 2 diabetes mellitus (T2DM) with chronic kidney disease (CKD). Referred by Dr. A. Kapoor, General Practice.

History of Presenting Illness

Mr. Menon is a 62-year-old man with T2DM diagnosed in 2014 and hypertension (HTN) since 2011. He reports good adherence to treatment and occasional fatigue after meals. He denies polyuria, visual disturbance, chest pain or foot symptoms.

Pertinent negatives: No history of diabetic foot ulcer or retinopathy. No history of myocardial infarction, stroke or peripheral vascular disease.

Past Medical History

- T2DM since 2014
- Hypertension since 2011
- CKD first noted in 2023
- Dyslipidaemia

Social History

- Retired bank clerk, lives with his wife
- Never smoked; no alcohol
- Vegetarian diet, walks 20 minutes daily

Examination

BLOOD PRESSURE **148/88 mmHg**	HEART RATE **78 /min, regular**	WEIGHT **84 kg**
BMI **29.1 kg/m2**	SPO2 **98% on room air**	FOOT EXAM **Monofilament sensation intact**

Investigations

Results from 12 Sep 2025 (outside laboratory), reviewed at this visit.

Test	Result	Reference	Comment
HbA1c	8.6 %	4.0 - 5.6 %	Above the 7.5 % target
Fasting plasma glucose	164 mg/dL	70 - 99 mg/dL	High
Serum creatinine	1.6 mg/dL	0.7 - 1.3 mg/dL	High
eGFR (CKD-EPI)	46 mL/min/1.73 m2	>= 60	CKD stage G3a
Urine albumin/creatinine ratio	64 mg/g	< 30 mg/g	Moderately increased

Current Medications

Medicine	Brand	Dose	Frequency
Metformin	Glucophage	1000 mg	Twice daily (BD) after meals
Glimepiride	Amaryl	2 mg	Once daily (OD) before breakfast

SYNTHETIC DATA — FOR DEMONSTRATION ONLY

Medicine	Brand	Dose	Frequency
Amlodipine	Norvasc	5 mg	Once daily (OD)
Telmisartan	Micardis	40 mg	Once daily (OD)
Atorvastatin	Lipitor	20 mg	At night

Allergies

No known drug allergies (NKDA).

Impression

T2DM with suboptimal glycaemic control (HbA1c 8.6 %) and CKD stage G3a with moderately increased albuminuria. Blood pressure above target.

Plan

- Continue current medications; review metformin dose if eGFR falls below 45.
- Repeat HbA1c, renal function and urine albumin/creatinine ratio in 4 months.
- Dietitian referral; foot-care education reinforced.
- Annual retinal screening to be booked with ophthalmology.

Electronically signed: **Dr. Helen Marsh, Consultant Endocrinologist** on 15 Sep 2025

*** End of report ***

SYNTHETIC DATA — FOR DEMONSTRATION ONLY