

## Secondary Metabolites from *Cycas lacrimans*

Vincent Antonio S. Ng<sup>1</sup>, Esperanza Maribel Agoo<sup>2</sup>, Chien-Chang Shen<sup>3</sup>, Consolacion Y. Ragasa<sup>1,4\*</sup>

<sup>1</sup>Chemistry Department, De La Salle University, 2401 Taft Avenue, 1004 Manila, Philippines

<sup>2</sup>Biology Department, De La Salle University, 2401 Taft Avenue, 1004 Manila, Philippines

<sup>3</sup>National Research Institute of Chinese Medicine, Ministry of Health and Welfare, 155-1, Li-Nong St., Sec. 2, Taipei, Taiwan

<sup>4</sup>Chemistry Department, De La Salle University Science & Technology Complex Leandro V. Locsin Campus, Biñan City, Laguna, Philippines

Available Online: 31<sup>st</sup> August, 2015

### ABSTRACT

Chemical investigation of the sarcotesta of *Cycas lacrimans*, a plant endemic to the Philippines afforded  $\beta$ -sitosteryl-3 $\beta$ -glucopyranoside-6'-O-palmitate (1), a mixture of alkyl 4-hydroxy-*trans*-cinnamate esters (2a) and alkyl 4-hydroxy-*cis*-cinnamate esters (2b),  $\beta$ -sitosterol (3), unsaturated triacylglycerols (4), and methyl fatty acid esters (5). The structures of 1-5 were identified by comparison of their NMR data with those reported in the literature.

**Keywords:** *Cycas lacrimans*, Cycadaceae,  $\beta$ -sitosteryl-3 $\beta$ -glucopyranoside-6'-O-palmitate, alkyl 4-hydroxy-*trans*-cinnamate esters, alkyl 4-hydroxy-*cis*-cinnamate esters,  $\beta$ -sitosterol, triacylglycerols, methyl fatty acid esters

### INTRODUCTION

*Cycas*, the only currently known genus of the Family Cycadaceae, are considered as fossil plants though they may have evolved only about 12 million years ago<sup>1</sup>. Their long existence and persistence through time have sparked special interest in their biology and evolution. The cycads resemble palms in morphology thus are commonly called sago palm. They bear naked seeds and are dioecious (male and female as separate individuals). These are widely distributed in the Tropics, with species found in Asia, Africa, Southeast Asia, Pacific, and Australia<sup>2</sup>. They also grow on volcanic, limestone, ultramafic, sandy, or even water-logged soils in grassland and forest habitats<sup>3</sup>.

In the Philippines, there are eleven (11) cycad species namely, *C. aenigma* K. D. Hill & Lindstrom, *C. curranii* (J. Schust.) K. D. Hill, *C. edentata* de Laub., *C. lacrimans* Lindstrom & K. D. Hill, *C. nitida* K. D. Hill & Lindstrom, *C. riuminiana* Porte ex Regel, *C. saxatilis* K. D. Hill & Lindstrom, *C. sancti-lasallei* Agoo & Madulid, *C. wadei* Merr., *C. vespertilio* Lindstrom & K. D. Hill, and *C. zambalensis* Madulid & Agoo<sup>3-5</sup>. All species, except for *C. edentata*, are endemic to the country<sup>4</sup>. *C. revoluta*, a widely cultivated species, is an introduced species from Japan and Taiwan.

The demand of *Cycas* species for domestic and international horticultural trade, grassland and forest fires of anthropogenic origin, and conversion of their natural habitats to settlements and other land uses have threatened to varying degrees the wild populations of the genus<sup>6</sup>. Some of these species which have been assessed and evaluated in the 2010 IUCN Red List of Threatened Species are *C. curranii*<sup>7</sup>, *C. wadei*<sup>8</sup> and *C. zambalensis* as

Critically Endangered (CR)<sup>7</sup>, *C. riuminiana* as Endangered (E)<sup>7</sup>, and *C. saxatilis* Vulnerable (V)<sup>9</sup>.

This study is part of our research on the chemical constituents of *Cycas* species that are endemic to the Philippines. We earlier reported the isolation of isopimarane-19-ol (I) from the megasporophyll lamina; 9 $\alpha$ H-isopimarane-7,15-diene (II) and triacylglycerols (III) from the bark; III, oleic acid (IV), and 1,2-dioleoylglycerol (V) from the leaflets; III,  $\beta$ -sitosterol (VI), and stigmastanol (VII) from the petiole and rachis; VI from the roots; and III and VI from the endotesta and sclerotesta of *C. lacrimans*<sup>10</sup>. In another study, we reported the isolation of III, VI, VII, and squalene (VIII) from the sarcotesta; III, VI, VII, and phytol fatty acid esters (IX) from the endotesta; III, VI, VII, and  $\beta$ -sitosteryl fatty acid esters (X) from the sclerotesta; and III and X from the bark of *C. sancti-lasallei*<sup>1</sup>. Another *Cycas* species, *C. vespertilio* yielded III, a mixture of VI and VII, pinosresinol (XI), sesamin (XII), and paulownin (XIII) from the cone base; III, VI, VII, XI, XIII, and lariciresinol (XIV) from the cataphylls; VI from the megasporophyll lamina; VI and a mixture of *trans*-4-hydroxycinnamate fatty acid esters (XV) and *cis*-4-hydroxycinnamate fatty acid esters (XVI) from the unripe sarcotesta; and III and VI from the ripe sarcotesta<sup>12</sup>. We recently reported the isolation of  $\beta$ -sitosteryl-3 $\beta$ -glucopyranoside-6'-O-palmitate (XVII) from the sarcotesta; X, unsaturated fatty acid methyl esters (XVIII), VI and VII from the endotesta; chlorophyll a (XIX) from the leaflets; and triacylglycerols (III) from the male cone of *C. edentata*<sup>13</sup>. We report herein the isolation of  $\beta$ -sitosteryl-3 $\beta$ -glucopyranoside-6'-O-palmitate (1), a mixture of alkyl 4-

\*Author for Correspondence




Figure 1: Chemical structures of  $\beta$ -sitosterol-3- $\beta$ -glucopyranoside-6'-*O*-palmitate (**1**), alkyl 4-hydroxy-*trans*-cinnamate esters (**2a**), alkyl 4-hydroxy-*cis*-cinnamate esters (**2b**),  $\beta$ -sitosterol (**3**), unsaturated triacylglycerols (**4**), and methyl fatty acid esters (**5**) from *C. lacrimans*.

hydroxy-*trans*-cinnamate esters (**2a**) and alkyl 4-hydroxy-*cis*-cinnamate esters (**2b**),  $\beta$ -sitosterol (**3**), unsaturated triacylglycerols (**4**), and methyl fatty acid esters from the sarcotesta of *Cycas lacrimans*. The structures of **1-5** were elucidated by comparison of their  $^1\text{H}$  and/or  $^{13}\text{C}$  NMR data with those reported in the literature. To the best of our knowledge this is the first report on the isolation of **1-2b** and **5** from *C. lacrimans*.

## MATERIALS AND METHODS

### General Experimental Procedure

NMR spectra were recorded on a Varian VNMRs spectrometer in  $\text{CDCl}_3$  at 600 MHz for  $^1\text{H}$  NMR and 150 MHz for  $^{13}\text{C}$  NMR spectra. Column chromatography was performed with silica gel 60 (70-230 mesh). Thin layer chromatography was performed with plastic backed plates coated with silica gel F<sub>254</sub> and the plates were visualized by spraying with vanillin/ $\text{H}_2\text{SO}_4$  solution followed by warming.

### Sample Collection

*Cycas lacrimans* sarcotesta were collected in 2013. Voucher specimens were collected and authenticated by

one of the authors (EMGA) and deposited in the De La Salle University- Manila Herbarium (DLSUH3113).

#### General Isolation Procedure

A glass column 18 inches in height and 1.0 inch internal diameter was used for the fractionation of the crude extracts. Ten milliliter fractions were collected. Fractions with spots of the same  $R_f$  values were combined and rechromatographed in appropriate solvent systems until TLC pure isolates were obtained. A glass column 12 inches in height and 0.5 inch internal diameter was used for the rechromatography. Five milliliter fractions were collected. Final purifications were conducted using Pasteur pipettes as columns. One milliliter fractions were collected.

#### Isolation of the Chemical Constituents of the Sarcotesta

The freeze-dried sarcotesta of *C. lacrimans* (105 g) were ground in a blender, soaked in  $\text{CH}_2\text{Cl}_2$  for 3 days and then filtered. The filtrate was concentrated under vacuum to afford a crude extract (0.4 g) which was chromatographed using increasing proportions of acetone in  $\text{CH}_2\text{Cl}_2$  at 10% increment. The 20% acetone in  $\text{CH}_2\text{Cl}_2$  fraction was rechromatographed (2  $\times$ ) using 10% EtOAc in petroleum ether to yield **5** (4 mg). The 40% acetone in  $\text{CH}_2\text{Cl}_2$  fraction was rechromatographed using 10% EtOAc in petroleum ether. The less polar fractions were combined and rechromatographed (2  $\times$ ) using 7.5% EtOAc in petroleum ether to afford **4** (5 mg). The more polar fractions were combined and rechromatographed (3  $\times$ ) using  $\text{CH}_3\text{CN}:\text{Et}_2\text{O}:\text{CH}_2\text{Cl}_2$  (0.5:0.5:9, v/v) to yield a mixture of **2a** and **2b** (3 mg). The 60% acetone in  $\text{CH}_2\text{Cl}_2$  fraction was rechromatographed (3  $\times$ ) using  $\text{CH}_3\text{CN}:\text{Et}_2\text{O}:\text{CH}_2\text{Cl}_2$  (2.5:2.5:5, v/v) to yield **1** (5 mg).

In the second isolation, the freeze-dried sarcotesta of *C. lacrimans* (140 g) were ground in a blender, soaked in  $\text{CH}_2\text{Cl}_2$  for 3 days and then filtered. The filtrate was concentrated under vacuum to afford a crude extract (1.2 g) which was chromatographed using increasing proportions of acetone in  $\text{CH}_2\text{Cl}_2$  at 10% increment. The 30% acetone in  $\text{CH}_2\text{Cl}_2$  fraction was rechromatographed (3  $\times$ ) in 7.5% EtOAc in petroleum ether to afford **4** (12 mg). The 40% acetone in  $\text{CH}_2\text{Cl}_2$  fraction was rechromatographed (4  $\times$ ) in 15% EtOAc in petroleum ether to afford **3** (15 mg) after washing with petroleum ether.

**$\beta$ -Sitosteryl-3 $\beta$ -glucopyranoside-6'-O-palmitate (1):**  $^1\text{H}$  NMR (600 MHz,  $\text{CDCl}_3$ ):  $\delta$  3.55 (m, H-3), 5.34 (m, H-6), 0.66 (s,  $\text{CH}_3$ -18), 0.99 (s,  $\text{CH}_3$ -19), 0.92 (3H, d,  $J = 6.0$  Hz,  $\text{CH}_3$ -21), 0.82 (3H, d,  $J = 7.2$  Hz,  $\text{CH}_3$ -26), 0.85 (d,  $J = 7.2$  Hz,  $\text{CH}_3$ -27), 0.86 (t,  $J = 7.2$  Hz,  $\text{CH}_3$ -29), 4.37 (d,  $J = 7.8$  Hz, H-1'), 3.36 (dd,  $J = 7.5, 9.0$ , H-2'), 3.56 (dd,  $J = 9.0, 10.0$ , H-3'), 3.36 (dd,  $J = 9.0, 10.0$  Hz, H-4'), 3.43 (m, H-5'), 4.23 (dd,  $J = 12.0, 2.0$  Hz, H-6a'), 4.49 (dd,  $J = 5.0, 12.0$  Hz, H-6b'), 2.34 (t,  $J = 7.5$  Hz, H-2'');  $^{13}\text{C}$  NMR (150 MHz,  $\text{CDCl}_3$ ):  $\delta$  37.24 (C-1), 29.71 (C-2), 79.53 (C-3), 38.88 (C-4), 140.26 (C-5), 122.19 (C-6), 31.93, 31.85 (C-7, C-8), 50.15 (C-9), 36.72 (C-10), 21.06 (C-11), 39.74 (C-12), 42.31 (C-13), 56.74 (C-14), 24.28 (C-15), 28.24 (C-16), 56.05 (C-17), 11.84 (C-18), 19.34 (C-19), 36.13 (C-20), 18.77 (C-21), 33.93 (C-22), 26.05 (C-23), 45.81 (C-24), 29.13 (C-25), 19.01 (C-26), 19.81 (C-27), 23.05 (C-28), 11.97 (C-29), 101.18 (C-1'), 73.58 (C-2'), 75.87 (C-

3'), 69.98 (C-4'), 73.98 (C-5'), 75.87 (C-6'), 174.80 (C-1''), 34.21 (C-2''), 24.94 (C-3''), 29.31 (C-4''), 29.52 (C-5''), 29.71 (C-6''), 29.71 (C-7''-C-12''), 29.36 (C-13''), 31.93 (C-14''), 22.69 (C-15''), 14.12 (C-16'').

**Trans-4-hydroxycinnamate fatty acid esters (2a):**  $^1\text{H}$  NMR (600 MHz,  $\text{CDCl}_3$ ):  $\delta$  6.83 (d,  $J = 9.0$  Hz, H-3, H-5), 7.43 (d,  $J = 9.0$  Hz, H-2, H-6), 7.61 (d,  $J = 15.6$  Hz, H-7), 6.28 (d,  $J = 15.6$  Hz, H-8), 4.16 (t,  $J = 6.6$  Hz, H-1'), 1.67 (2H, quintet,  $J = 6.6$  Hz, H-2'), 1.23–1.38 (H-3'-H-n'), 0.88 (t,  $J = 6.6$  Hz, Me);  $^{13}\text{C}$  NMR (150 MHz,  $\text{CDCl}_3$ ):  $\delta$  127.41 (C-1), 115.81 (C-3, C-5), 129.90 (C-2, C-6), 157.47 (C-4), 144.10 (C-7), 115.87 (C-8), 167.44 (C-9), 64.46/64.61 (C-1'), 28.24 (C-2'), 129.90 (=CH), 27.79 (allylic  $\text{CH}_2$ '), 22.69, 26.04, 29.06–29.70 ( $\text{CH}_2$ )<sub>n</sub>, 31.92–31.65 ( $\text{CH}_2$ '), 14.12 ( $\text{CH}_3$ ' terminal).

**Cis-4-hydroxycinnamate fatty acid esters (2b):**  $^1\text{H}$  NMR (600 MHz,  $\text{CDCl}_3$ ):  $\delta$  0.88 (3H, t,  $J = 6.6$  Hz, Me), 1.23–1.38 (m), 1.67 (2H, quintet,  $J = 6.6$  Hz, H-2'), 4.09 (2H, t,  $J = 6.6$  Hz, H-1'), 5.83 (1H, d,  $J = 12.6$  Hz, H-8), 6.78 (2H, d,  $J = 9.0$  Hz, H-3, H-5), 6.82 (1H, d,  $J = 12.6$  Hz, H-7), 7.59 (2H, d,  $J = 9.0$  Hz, H-2, H-6);  $^{13}\text{C}$ -NMR (150 MHz,  $\text{CDCl}_3$ ):  $\delta$  127.11 (C-1), 114.90 (C-3, C-5), 132.33 (C-2, C-6), 156.48 (C-4), 143.02 (C-7), 117.42 (C-8), 166.59 (C-9), 64.46/64.61 (C-1'), 22.69, 26.04, 28.24 (C-2'), 129.90 (=CH), 29.06–29.70 ( $\text{CH}_2$ )<sub>n</sub>, 31.92–31.65 ( $\text{CH}_2$ '), 14.12 ( $\text{CH}_3$ ' terminal).

**$\beta$ -Sitosterol (3):**  $^1\text{H}$  NMR (600 MHz,  $\text{CDCl}_3$ ):  $\delta$  3.51 (m, H-3), 2.28, 2.24 (H<sub>2</sub>-4), 5.33 (dd,  $J = 5.4, 2.4$  Hz, H-6), 0.66 (s,  $\text{CH}_3$ -18), 0.99 (s,  $\text{CH}_3$ -19), 0.91 (d,  $J = 6.6$  Hz,  $\text{CH}_3$ -21), 0.80 (d,  $J = 6.6$  Hz,  $\text{CH}_3$ -26), 0.82 (d,  $J = 7.2$  Hz,  $\text{CH}_3$ -27), 0.86 (t,  $J = 7.2$  Hz,  $\text{CH}_3$ -29).  $^{13}\text{C}$  NMR (150 MHz,  $\text{CDCl}_3$ ):  $\delta$  37.24 (C-1), 31.66 (C-2), 71.81 (C-3), 42.30 (C-4), 140.75 (C-5), 121.72 (C-6), 31.90, 31.91 (C-7, C-8), 50.12 (C-9), 36.14 (C-10), 21.07 (C-11), 39.76 (C-12), 42.30 (C-13), 56.76 (C-14), 24.30 (C-15), 28.24 (C-16), 56.04 (C-17), 11.85 (C-18), 19.39 (C-19), 36.50 (C-20), 19.02 (C-21), 33.93 (C-22), 29.14 (C-23), 45.82 (C-24), 26.06 (C-25), 18.77 (C-26), 19.81 (C-27), 23.06 (C-28), 11.97 (C-29).

**Triacylglycerols (4):**  $^1\text{H}$  NMR (600 MHz,  $\text{CDCl}_3$ ):  $\delta$  4.28 (2H, dd,  $J = 4.2, 12.0$  Hz, glyceryl  $\text{CH}_2\text{O}$ ), 4.12 (2H, dd,  $J = 6.0, 12.0$  Hz, glyceryl  $\text{CH}_2\text{O}$ ), 5.32 (1H, m, glyceryl CHO), 2.30 (6H, t,  $J = 7.2$  Hz,  $\alpha$ - $\text{CH}_2$ ), 5.33 (m, olefinic H), 2.75 (double allylic  $\text{CH}_2$ ), 1.98–2.05 (allylic,  $\text{CH}_2$ ), 1.23–1.35 ( $\text{CH}_2$ ), 0.87 (t,  $J = 6.6$  Hz,  $\text{CH}_3$ );  $^{13}\text{C}$  NMR (150 MHz,  $\text{CDCl}_3$ ):  $\delta$  62.09 (glyceryl  $\text{CH}_2$ ), 68.87 (glyceryl CH), 173.29, 173.25 (C=O  $\alpha$ ), 172.84 (C=O  $\beta$ ), 34.03, 34.05, 34.19 (C-2), 24.84, 24.86 (C-3), 29.05, 29.08, 29.12 (C-4), 29.17, 29.20, 29.27 (C-5), 29.48 (C-6), 22.57, 22.68 (C-8), 130.23, 130.01, 129.70 (C-9), 127.74, 128.29, 129.68 (C-10), 25.61, 27.17, 27.20, 27.22, 29.32, 29.35, 29.36, 29.52, 29.62, 29.66, 29.70, 29.76 ( $\text{CH}_2$ ), 31.52, 31.90, 31.92 ( $\text{CH}_2$ ), 14.07, 14.11, 14.27 (terminal  $\text{CH}_3$ ).

**Fatty acid methyl esters (5):**  $^1\text{H}$  NMR (600 MHz,  $\text{CDCl}_3$ ):  $\delta$  0.86 (t,  $J = 6.6$  Hz), 0.96 (t,  $J = 7.2$  Hz), 1.23–1.35 (m), 1.53–1.61 (m), 2.02–2.06 (m), 2.28 (t,  $J = 7.5$  Hz), 2.78 (t,  $J = 5.7$  Hz), 3.65 (s), 5.30–5.37 (m);  $^{13}\text{C}$  NMR (150 MHz,  $\text{CDCl}_3$ ):  $\delta$  14.07, 14.12, 14.27, 20.54, 22.57, 22.69, 24.94, 24.96, 25.52, 25.60, 25.62, 27.15, 27.19, 27.19, 29.09, 29.11, 29.15, 29.25, 29.31, 29.34, 29.35, 29.45, 29.52,

29.58, 29.66, 29.70, 29.76, 31.52, 31.90, 31.92, 34.10, 34.12, 51.44, 127.10, 127.72, 127.90, 128.03, 128.24, 128.27, 129.75, 130.00, 130.05, 130.21, 130.27, 131.96, 174.32.

## RESULTS AND DISCUSSION

Silica gel chromatography of the dichloromethane extract of the sarcotesta of *C. lacrimans* afforded  $\beta$ -sitosteryl-3 $\beta$ -glucopyranoside-6'-O-palmitate (**1**)<sup>14,15</sup>, a mixture of alkyl 4-hydroxy-*trans*-cinnamate ester (**2a**)<sup>12</sup> and alkyl 4-hydroxy-*cis*-cinnamate ester (**2b**)<sup>12,16</sup>,  $\beta$ -sitosterol (**3**)<sup>10,17</sup>, unsaturated triacylglycerols (**4**)<sup>10</sup>, and methyl fatty acid esters (**5**)<sup>18</sup>. The structures of **1-5** were identified by comparison of their <sup>1</sup>H and <sup>13</sup>C NMR data with literature data.

$\beta$ -sitosterol (**3**) isolated from the sarcotesta was also present in the petiole and rachis, roots, endotesta and sclerotesta of *C. lacrimans*<sup>0</sup>, while triacylglycerols (**4**) obtained from the sarcotesta were also found in the bark, leaflets, petiole and rachis, roots, endotesta and sclerotesta of *C. lacrimans*<sup>0</sup>. Results of this study indicate that *C. lacrimans* shares similar chemical characteristics with other members of the family Cycadaceae: *C. edentata* which contained  $\beta$ -sitosteryl-3 $\beta$ -glucopyranoside-6'-O-palmitate (**1**) and methyl fatty acid esters (**5**)<sup>13</sup> and *C. vespertilio* which yielded alkyl 4-hydroxy-*trans*-cinnamate esters (**2a**) and alkyl 4-hydroxy-*cis*-cinnamate esters (**2b**)<sup>12</sup>.

## ACKNOWLEDGEMENT

A research grant from the Commission on Higher Education – Philippine Higher Education Research Network (CHED-PHERNet) of the Philippines is gratefully acknowledged.

## REFERENCES

1. Nagalingum NS, Marshal CR, Quental TB, Tai HS, Little DP, Matthews S. Recent synchronous radiation of a living fossil. *Science* 2011; 334:796–799.
2. Donaldson JS. Cycads. Status Survey and Conservation Action Plan. IUCN Gland, Switzerland and Cambridge, U.K.; 2003.
3. Madulid DA, Agoo EMG. 2009. Taxonomy and conservation of Philippine Cycads. *Blumea* 2009; 54:99–102.
4. Lindstrom AJ, Hill KD, Stanberg LC. The genus *Cycas* (Cycadaceae) in the Philippines. *Telopea* 2008; 12:119–145.
5. Agoo EMG, Madulid DA. *Cycas sancti-lasallei* (Cycadaceae), a new species from the Philippines. *Blumea* 2012; 57:131–133.
6. IUCN Red List of Threatened Species. Version 2010.4. <www.iucnredlist.org>. Downloaded on 09 February 2011.
7. Agoo EMG, Madulid DA, Linis VC, Sambale E. In: IUCN 2010. IUCN Red List of Threatened Species. Version 2013.2. <www.iucnredlist.org>. Downloaded on 16 December 2013.
8. Hill KD. 2010. *Cycas wadei*. In: IUCN 2013. IUCN Red List of Threatened Species. Version 2013.2. <www.iucnredlist.org>. Downloaded on 26 December 2013.
9. Bosenberg, JD. 2010. In: IUCN 2013. IUCN Red List of Threatened Species. Version 2013.2. <www.iucnredlist.org>. Downloaded on 16 December 2013.
10. Ragasa CY, Ng VAS, Agoo EM, Shen C-C. Chemical constituents of *Cycas lacrimans*. *Int. J. Pharmacog. Phytochem. Res.* 2015; 7(3):616–620.
11. Ng VAS, Agoo EM, Shen C-C, Ragasa CY. Chemical constituents of *Cycas sancti-lasallei*. *J. Appl. Pharm. Sci.* 2015; 5(Suppl. 1):12–17.
12. Ragasa CY, Ng VAS, Agoo EM, Shen C-C. Chemical constituents of *Cycas vespertilio*. *Braz. J. Pharmacog.* 2015; doi:10.1016/j.bjp.2015.06.002.
13. Ng VAS, Agoo EM, Shen C-C, Ragasa CY. Secondary metabolites from *Cycas edentata*. *J. Pharm. Sci. Res.* (in-press).
14. Tsai P-W, de Castro-Cruz K, Shen C-C, Ragasa CY. Chemical constituents of *Ficus odorata*. *Pharm. Chem. J.* 2012; 46(4):225–227.
15. Nguyen AT, Malonne H, Duez P, Vanhaelen M, Fontaine J, Vanhaelen-Fastre R. Cytotoxic constituents from *Plumbago zeylanica*. *Fitoter.* 2004; 75:500 – 504.
16. Nishimura K, Takenaka Y, Kishi M, Tanahashi T, Yoshida H, Okuda C, Mizushima Y. Synthesis and DNA polymerase  $\alpha$  and  $\beta$  inhibitory activity of alkyl p-coumarates and related compounds. *Chem. Pharm. Bull.* 2009; 57:476–480.
17. Ragasa CY, Ng VAS, De Los Reyes MM, Mandia EH, Oyong GG, Shen C-C. Chemical constituents and cytotoxicity of the leaves of *Dysoxylum gaudichaudianum* (A. Juss.) Miq. *Der Pharma Chemica* 2014; 6(5):182–187.
18. Basumatary S, Deka DC. Identification of fatty acid methyl esters in biodiesel from *Pithecellobium monadelphum* seed oil. *Der Chemica Sinica.* 2012; 3(6): 1384–1393.