

Determining the composition and structure of hot Jupiter atmospheres using high-resolution spectroscopy

Diogo Godinho Gomes – Student No. 11407018

Anton Pannekoek Institute for Astronomy (API), University of Amsterdam, Science Park 904, 1098 XH Amsterdam
e-mail: seccr-astro-science@uva.nl

February 19, 2018

ABSTRACT

Context. ...

Aims. ...

Methods. ...

Results. ...

Conclusions. ...

Key words. ... – ... – ...

1. Observations

The targets of the observations were the host star HD 209458 ($B = 8.21 \pm 0.02$ mag (Høg et al. 2000), $K = 6.308 \pm 0.026$ mag (Cutri et al. 2003)) and its companion, HD 209458 b, a Hot Jupiter in a close transiting orbit ($M = 0.714 \pm 0.017 M_J$, $a = 0.04747 \pm 0.00055$ au, $i = 86.590 \pm 0.046$ deg (Southworth 2010)), for approximately 5 hours on the night of July 8th, 2014. The instrument used was the CRYogenic high-resolution InfraRed Echelle Spectrograph (CRIRES), mounted at the Nasmyth focus A of the 8.2-m UT1/Antu at the ESO's Very Large Telescope (VLT), Cerro Paranal, Chile. This infrared detector contains four Aladdin III InSb detectors, each of them with size 1024×512 pixels, and gaps of around 280 pixels between each pair of detectors. These observations (concerning the determination of the structure and composition of HD 209458 b's atmosphere) are part of ESO's program 186.C-0289, which intends to study the brightest transiting and non-transiting hot-Jupiter systems visible from Cerro Paranal.

During the observations, 33 spectra were collected on the L_[1] .. Band – the atmospheric transmission window in the mid-infrared around $3.5 \mu\text{m}$ – each one of them being the result of two different sets (A and B) of exposures of 8×30.0 s each, with a delay of about 1 min between exposures. Although setting a fixed position for the telescope relatively to the slit increases stability and decreases the overheads, in order to achieve an easier sky-background subtraction, these observations were done in the nodding mode: the telescope was nodded along the slit by 10 arcsec between each set of exposures in a ABBA pattern. As a consequence, the final extracted data consist of an AB or a BA pair per spectrum.

The range of wavelengths covered with the observations was $3.459 - 3.543 \mu\text{m}$, with a resolution of $R \sim 100000$ which corresponds to approximately 3 km s^{-1} per resolution element (or 1.5 km s^{-1} per pixel) on the exoplanet's radial velocity – this high resolution results from using a 0.2 arcsec slit together with the Multi Application Curvature Adaptive Optics (MACAO)

system (Arsenault et al. 2003). HD 209458 b was continuously observed shortly after a secondary eclipse, when a significant portion of the day side of the planet was visible, during the orbital phases ranging between 0.530 and 0.591. For these observed orbital phases and a radial velocity semi-amplitude (approximately the same as the orbital velocity, as the system has an inclination of almost 90 deg) of $K_p \approx 140 \text{ km s}^{-1}$ (Snellen et al. 2010), the expected variation in the exoplanet's radial velocity throughout the spectra is about $\Delta RV_p \approx 50 \text{ km s}^{-1}$ (or 32 pixels on the CRIRES detectors).

During the observations, the seeing worsened considerably towards the final spectra: it reached its minimum on the eighth spectrum, 0.67 arcsec, and its maximum on the very last one, 1.50 arcsec.

Acknowledgements. ...

References