

# Lecture 3: Equivariant Convolutional Neural Networks on Homogeneous Spaces

Geometry and Topology in Machine Learning Seminar

June 16th, 2025

# Recall: Representations and Bundles

Lecture 3:  
Equivariant  
Convolutional  
Neural  
Networks on  
Homogeneous  
Spaces

CNNs with  
respect to  
Groups

Equivariant  
Neural  
Networks

Example:  
3D Steerable  
CNNs

A glimpse of  
implementations

- ① Let  $G$  be a **group** and  $V$  be a **vector space**.
  - A  **$G$ -representation** is a group action of  $G$  on  $V$  where the actions are all linear maps.
  - Equivalently, this is the data of a map  $\rho : G \rightarrow \text{GL}(V)$  that respects the group structure on both sides<sup>1</sup>.
- ② A **principal  $G$ -bundle** is a fiber bundle  $\phi : X \rightarrow B$  where the fibers are  **$G$ -torsors**, i.e. spaces where  $G$  acts **freely** and **transitively** on.
- ③ Let  $\rho : G \rightarrow \text{GL}(V)$  be an  **$G$ -representation**. The **associated vector bundle** of  $\phi : X \rightarrow B$  with respect to  $\rho$  is

$$\phi' : E := X \times_{\rho} V := X \times V / \sim \rightarrow B$$

where  $(x \cdot g, y) \sim (x, \rho(g) \cdot y)$ . Each fiber over  $b \in B$  is a copy of  $V$ , equipped with the action of  $G$  from  $\rho$ .  $E$  is the **vector bundle associated to  $\phi$  via  $\rho$** .

---

<sup>1</sup>i.e. a group homomorphism  $\rho(gh) = \rho(g)\rho(h)$

# Recall: Sections and Mackey Functions

- ④ Given an associated vector bundle  $\phi' : E \rightarrow B$ , a **section**<sup>2</sup> is a map  $s : B \rightarrow E$  such that  $\phi' \circ s = id_B$ .
- ⑤ Let  $H \leq G$  and  $V$  be an  $H$ -representation  $\rho$ , a **Mackey function** is a map

$$f : G \rightarrow V, f(gh) = \rho(h^{-1})f(g), \quad \forall g \in G, h \in H.$$

In the last lecture, we focused on the case where the base space  $B$  is a **homogeneous space**, meaning we can write  $B = G/H$ .

- The quotient map  $G \rightarrow G/H$  is a **principal H-bundle**.
- Given an associated vector bundle  $E \rightarrow G/H$  from an  $H$ -representation  $\rho$ , there is an isomorphism between

$$\Gamma(E) = \{\text{sections of } E\} \text{ and } \mathcal{I}_G = \{\text{Mackey functions for } (H, \rho)\}$$

---

<sup>2</sup>Also known as a **field**, or a **stack of features**

# Recall: Induced Representations

Lecture 3:  
Equivariant  
Convolutional  
Neural  
Networks on  
Homogeneous  
Spaces

CNNs with  
respect to  
Groups

Equivariant  
Neural  
Networks

Example:  
3D Steerable  
CNNs

A glimpse of  
implementations

- ⑥ **Induced representations** describe how features transform in a globally consistent vector bundle with  $G$ -actions.

## Definition

Let  $H \leq G$  and  $\rho : H \rightarrow \text{GL}(V)$  be an  $H$ -representation. The **induced representation**  $\pi = \text{Ind}_H^G \rho$  is the data:

- ① The vector space of **Mackey functions**  $\mathcal{I}_G$
- ② An action of  $G$  on  $\mathcal{I}_G$  by

$$g \cdot f(k) := f(g^{-1}k)$$

Note in the case of  $G \rightarrow G/H$ ,  $\mathcal{I}_G$  is isomorphic to  $\Gamma(E)$ , so the induced representation describes how  $\Gamma(E)$  transforms!

Lecture 3:  
Equivariant  
Convolutional  
Neural  
Networks on  
Homogeneous  
Spaces

CNNs with  
respect to  
Groups

Equivariant  
Neural  
Networks

Example:  
3D Steerable  
CNNs

A glimpse of  
implementations

- 1 CNNs with respect to Groups
- 2 Equivariant Neural Networks
- 3 Example: 3D Steerable CNNs
- 4 A glimpse of implementations

## Let us talk more about ML in this lecture!

In the last lecture, we have seen the analogy of **feature spaces** and an example of **spherical CNNs**. Today, we will focus on **layers between feature spaces** in our set-up. Specifically, we should address the following questions:

- 1 Doing **convolutions** seems to require an integral - how do we integrate on groups?
- 2 Even if we can integrate on nice groups, what does **convolution** mean in this context?
- 3 How do we design **convolutional neural networks** from this?
- 4 How does **convolution** relate to **equivariance**?

For example, let us look at the group  $(\mathbb{R}^1, +)$  (the setting for 1D CNNs) where the operation is addition. Integration over  $\mathbb{R}^1$  is fundamentally based on:

$$\int_{[a,b]} 1dx = \text{length}([a, b]).$$

**Observation:** The **length** on  $\mathbb{R}^1$  satisfies the following:

- 1 For any interval  $[a, b]$ , if we shift the entire interval by  $t \in \mathbb{R}$ , the **length does not change**.
- 2 The length of any non-empty open interval is not zero.
- 3 The length of any closed and bounded interval is finite.

If we move to the group  $(\mathbb{R}^n, +)$  and **length** to **n-dimensional volume**, analogous statements to above would hold.

# Going More Generally

Lecture 3:  
Equivariant  
Convolutional  
Neural  
Networks on  
Homogeneous  
Spaces

CNNs with  
respect to  
Groups

Equivariant  
Neural  
Networks

Example:  
3D Steerable  
CNNs

A glimpse of  
implementations

- 1 For any interval  $[a, b]$ , if we shift the entire interval by  $t \in \mathbb{R}$ , the **length does not change**.
- 2 The length of any non-empty open interval is not zero.
- 3 The length of any closed and bounded interval is finite.

The general properties they should correspond to a group  $G$  are

- 1 **Length**  $\rightarrow$  a  $\mathbb{R}_{\geq 0}$ -valued function  $\mu$  on “nice” subsets of  $G$  (specifically a **measure**).
- 2  $t + [a, b]$  is an example of a **left coset** of  $[a, b]$  with respect to  $t \in \mathbb{R}$ . Thus, we can more generally consider:

$$gS = \{gs \in G \mid s \in S\}, g \in G, S \subset G \text{ and } \mu(gS) = \mu(S).$$

- 3 **Open intervals**  $\rightarrow$  **Open sets**.
- 4 **Closed and bounded intervals**  $\rightarrow$  **Compact sets**.



# Integration on Groups

Lecture 3:  
Equivariant  
Convolutional  
Neural  
Networks on  
Homogeneous  
Spaces

CNNs with  
respect to  
Groups

Equivariant  
Neural  
Networks

Example:  
3D Steerable  
CNNs

A glimpse of  
implementations

Let  $G$  be a **nice group**<sup>3</sup>, there is a way to define **integration** on  $G$  as follows:

## Theorem

$G$  admits a **left Haar measure**, which is a **measure**  $\mu : F(G)^4 \subset \{\text{Subsets of } G\} \rightarrow \mathbb{R}$  such that:

- ①  $\mu$  is **left-translation invariant**, ie.  $\mu(gS) = \mu(S)$  for all  $g \in G$  and  $S \in F(G)$ . Here
- ②  $\mu(U) > 0$  for all non-empty open subset of  $G$ .
- ③  $\mu(K) < \infty$  for all compact subsets of  $G$ .

Note that **left Haar measure** is **unique up to scaling**.

---

<sup>3</sup>locally compact and Hausdorff, this is satisfied for all examples of groups we have seen so far

<sup>4</sup>This is the Borel  $\sigma$ -algebra, but you can just think of this as **reasonable subsets**.

# Integration on Groups (Continued)

Lecture 3:  
Equivariant  
Convolutional  
Neural  
Networks on  
Homogeneous  
Spaces

CNNs with  
respect to  
Groups

Equivariant  
Neural  
Networks

Example:  
3D Steerable  
CNNs

A glimpse of  
implementations

There is a similar way to define a **right Haar measure** with **right cosets**. We say  $G$  is **unimodular** if its left and right measures agree after scaling.

## Definition

For unimodular nice groups  $G$ , we call  $\mu$  **the Haar measure** of  $G$ .

- 1 If  $G$  is **compact** (ex.  $\mathrm{SO}(n)$ ), then it is **unimodular**.
- 2 If  $G$  is **abelian**, then it is also **unimodular**.
- 3 (For Algebraic Enthusiasts): If more generally the **abelianization of  $G$**  is finite, then it is also **unimodular**.
- 4 (For Lie theory enthusiasts): For a Lie group  $G$  such that  $\det(\mathrm{Ad}_g) = 1$  for all  $g \in G$ ,  $G$  is **unimodular**.

# Examples of Integration on Groups

Given a **Haar measure**  $\mu$  on  $G$ , we can use it to define an **integration**  $\int_G \bullet d\mu$  on  $G$  such that

$$\int_G 1_S d\mu := \mu(S).$$

- ① If  $G$  is  $(\mathbb{R}^n, +)$ , this is just the usual (Lebesgue) **integration** from calculus.
- ② If  $G$  is  $S^1$  (group operation being rotation), for a function  $f : S^1 \rightarrow \mathbb{R}$ , this is the same as viewing  $f$  as a function  $f' : [0, 1] \rightarrow \mathbb{R}$  with  $f(0) = f(1)$  and

$$\int_{S^1} f d\mu = \int_0^1 f' dx.$$

- ③ If  $G$  is a **finite group** (or more generally a **discrete group**<sup>5</sup>), then the **integration** is the same as **discrete summation**.

---

<sup>5</sup>Think  $SL_2(\mathbb{Z})$  for example

In 1D functions, for two functions  $f, g : \mathbb{R} \rightarrow \mathbb{R}$ , convolution is given by

$$f * g(t) = \int_{\mathbb{R}} f(\tau)g(t - \tau)d\tau = \int_{\mathbb{R}} f(\tau)g((- \tau) + t)d\tau.$$

Another way to see this however is that the term  $(-\tau) + t$  is really an example of

$$g^{-1}g' \in G \text{ for some elements } g, g' \in G.$$

# Convolution on Groups

Lecture 3:  
Equivariant  
Convolutional  
Neural  
Networks on  
Homogeneous  
Spaces

CNNs with  
respect to  
Groups

Equivariant  
Neural  
Networks

Example:  
3D Steerable  
CNNs

A glimpse of  
implementations

More generally, let  $V_1, V_2$  be vector spaces (ie. codomain of features in CNNs).

## Definition

Given a function  $f : G \rightarrow V_1$  and  $\kappa : G \rightarrow \text{Hom}(V_1, V_2)$ , the **convolution** of  $f$  and  $\kappa$  is the function

$$f \star \kappa : G \rightarrow V_2, (f \star \kappa)(g) := \int_G \kappa(g^{-1}g')f(g')d\mu(g')$$

Here the integration is taken with respect to the **variable**  $g'$ .

Here  $\kappa$  is often called the (one-argument)  **$G$ -convolution kernel**.

Let  $G$  be a group.

## Definition

A  **$G$ -Convolutional Neural Network Layer** (**G-CNN Layer**) is a layer  $L_i : \mathcal{F}_i \rightarrow \mathcal{F}_{i+1}$  in a neural network where

- 1  $\mathcal{F}_i$  is the **feature space** of  $G$  with respect to some  $(H_i \leq G, \rho_i : H_i \rightarrow \mathrm{GL}(V_i))$ . In other words,  $\mathcal{F}_i$  is the space of **sections**  $\Gamma(E_i)$  of the associated vector bundle  $E_i = G \times_{\rho_i} V_i \rightarrow G/H_i$ .  $\mathcal{F}_{i+1}$  is defined similarly.
- 2 The map  $L_i : \Gamma(E_i) \rightarrow \Gamma(E_{i+1})$  is a **convolution** with respect to some  $\kappa_i$  where  $\kappa_i$  is to be **optimized/learned**.

Here we recall that  $\Gamma(E_i)$  (resp.  $\Gamma(E_{i+1})$ ) is isomorphic to the collection of **Mackey functions**  $f : G \rightarrow V_i$  (resp.  $f : G \rightarrow V_{i+1}$ ), so the **convolution** here makes sense.

Lecture 3:  
Equivariant  
Convolutional  
Neural  
Networks on  
Homogeneous  
Spaces

CNNs with  
respect to  
Groups

Equivariant  
Neural  
Networks

Example:  
3D Steerable  
CNNs

A glimpse of  
implementations

- 1 CNNs with respect to Groups
- 2 Equivariant Neural Networks
- 3 Example: 3D Steerable CNNs
- 4 A glimpse of implementations

Let  $G$  be a group.

## Definition

A  **$G$ -Equivariant Neural Network Layer** is a layer

$L_i : \mathcal{F}_i \rightarrow \mathcal{F}_{i+1}$  in a neural network where

- 1  $\mathcal{F}_i$  is the **feature space** of  $G$  with respect to some  $(H_i \leq G, \rho_i : H_i \rightarrow \text{GL}(V_i))$ . In other words,  $\mathcal{F}_i$  is the space of **sections**  $\Gamma(E_i)$  of the associated vector bundle  $E_i = G \times_{\rho_i} V_i \rightarrow G/H_i$ .  $\mathcal{F}_{i+1}$  is defined similarly.
- 2 The map  $L_i : \Gamma(E_i) \rightarrow \Gamma(E_{i+1})$  is a **linear map** that is **equivariant** with respect to the **induced representation structure**  $(\pi_i \text{ and } \pi_{i+1})$  on both sides.



Recall the following **slogan** from Lecture 1:

“Translation equivariant linear maps are convolutions.”

Since we have defined convolutions to a broader generality now, the **new slogan** we would hope is that:

“ $G$ -equivariant linear maps are  $G$ -convolutions.”

# Convolution is All You Need

Lecture 3:  
Equivariant  
Convolutional  
Neural  
Networks on  
Homogeneous  
Spaces

CNNs with  
respect to  
Groups

Equivariant  
Neural  
Networks

Example:  
3D Steerable  
CNNs

A glimpse of  
implementations

## Theorem (Convolution is All You Need [Cohen et al., 2019])

Under reasonable assumptions<sup>6</sup>, a  $G$ -equivariant neural network layer is a  $G$ -convolutional neural network layer.

In other words, under reasonable hypothesis, a (bounded) equivariant linear map  $\phi$  can be expressed as some convolution with respect to  $G$ .

---

<sup>6</sup>which will be clarified in the proceeding slides

# Why Should We Expect This? (Proof Sketch)

Let us consider a (bounded) **equivariant linear map**

$$\phi : \Gamma(E_i) \cong \mathcal{I}_G^i \rightarrow \Gamma(E_{i+1}) \cong \mathcal{I}_G^{i+1}.$$

(Here  $\mathcal{I}_G^i$  refers to the respective **Mackey function**).

In many nice cases, it is possible to **represent  $\phi$  as an integral transform**. In the sense that, for any  $f : G \rightarrow V_i \in \mathcal{I}_G^i$ , there exists a **two-argument kernel**  $\kappa^+ : G \times G \rightarrow \text{Hom}(V_i, V_{i+1})$  such that

$$(\phi f)(y) = \int_G \kappa^+(y, x) f(x) d\mu_G(x)^7 \quad (\diamond)$$

If  $V_i$  and  $V_{i+1}$  are both **finite dimensional**, we can fix a choice of basis to make this a **matrix-valued kernel**.

---

<sup>7</sup>The right hand side is called an **integral transform**.

# Why Should We Expect This? (Proof Sketch)

Lecture 3:  
Equivariant  
Convolutional  
Neural  
Networks on  
Homogeneous  
Spaces

CNNs with  
respect to  
Groups

Equivariant  
Neural  
Networks

Example:  
3D Steerable  
CNNs

A glimpse of  
implementations

Let  $\pi_i, \pi_{i+1}$  be the **induced representations** for  $\mathcal{I}_G^i$  and  $\mathcal{I}_G^{i+1}$  respectively. Since  $\phi$  is **equivariant** w.r.t the induced actions on the functions, this means that for all  $u, x \in G$  and  $f \in \mathcal{I}_G^i$

$$[\phi(\pi_i(u) \cdot f)](x) = [\pi_{i+1}(u) \cdot (\phi(f))](x) \quad (\spadesuit).$$

Recall the action of an arbitrary induced representation  $\pi$  on  $f : G \rightarrow V$  is

$$g \cdot f(x) := f(g^{-1}x), \forall g, x \in G.$$

Expanding both sides of  $(\spadesuit)$  out using the **integral transforms**, we have that

$$\int_G \kappa^+(y, x) f(u^{-1}x) d\mu_G(x) = \int_G \kappa^+(u^{-1}y, x) f(x) d\mu_G(x). \quad (\square)$$

# Why Does ( $\spadesuit$ ) Expand to ( $\square$ )?

Starting with the equivariant condition

$$[\phi(\pi_i(u) \cdot f)](x) = [\pi_{i+1}(u) \cdot (\phi(f))](x) \quad (\spadesuit),$$

we have that

$$\begin{aligned} \text{LHS} &= [\phi(\pi_i(u) \cdot f)](y) \\ &= \int_G \kappa^+(y, x) [\pi_i(u) \cdot f](x) d\mu_G(x) \quad \text{by } (\diamond). \\ &= \int_G \kappa^+(y, x) f(u^{-1}x)(x) d\mu_G(x) \end{aligned}$$

where the last equality follows from the induced action on functions (pull backs on  $f$ )  $(\pi_i(u) \cdot f)(x) = f(u^{-1}x)$ .

For the RHS, we first apply the induced action and then the integral transform,

$$\begin{aligned} \text{RHS} &= [\pi_{i+1}(u) \cdot (\phi(f))](y) = \phi(f)(u^{-1}y) \\ &= \int_G \kappa^+(u^{-1}y, x) f(x) d\mu_G(x) \end{aligned}$$

# Why Should We Expect This? (Proof Sketch)

Lecture 3:  
Equivariant  
Convolutional  
Neural  
Networks on  
Homogeneous  
Spaces

CNNs with  
respect to  
Groups

Equivariant  
Neural  
Networks

Example:  
3D Steerable  
CNNs

A glimpse of  
implementations

Now we have that

$$\int_G \kappa^+(y, x) f(u^{-1}x) d\mu_G(x) = \int_G \kappa^+(u^{-1}y, x) f(x) d\mu_G(x).$$

Consider a change of variables  $x \mapsto ux$  on the left hand side. This is okay since we are integrating over all of  $G$ , and we have that

$$\begin{aligned} \int_G \kappa^+(y, ux) f(u^{-1}ux) d\mu_G(x) &= \int_G \kappa^+(y, x) f(u^{-1}x) d\mu_G(x) \\ &= \int_G \kappa^+(u^{-1}y, x) f(x) d\mu_G(x) \end{aligned}$$

# Why Should We Expect This? (Proof Sketch)

Lecture 3:  
Equivariant  
Convolutional  
Neural  
Networks on  
Homogeneous  
Spaces

CNNs with  
respect to  
Groups

Equivariant  
Neural  
Networks

Example:  
3D Steerable  
CNNs

A glimpse of  
implementations

Thus, we have that

$$\int_G \kappa^+(y, ux) f(x) d\mu_G(x) = \int_G \kappa^+(u^{-1}y, x) f(x) d\mu_G(x).$$

Since this equality holds for all  $f$ 's, it implies that

$$\kappa^+(y, ux) = \kappa^+(u^{-1}y, x).$$

Doing another round of **variable substitutions**<sup>8</sup>, we have that

$$\kappa^+(uy, ux) = \kappa^+(y, x)$$

for all  $x, y, u \in G$ .

---

<sup>8</sup>Consider relabeling  $y \rightarrow uy$  in the equality  $\kappa^+(y, ux) = \kappa^+(u^{-1}y, x)$ .

# Why Should We Expect This? (Proof Sketch)

Now since we have the equality

$$\kappa^+(uy, ux) = \kappa^+(y, x) \quad (\dagger),$$

we observe that

$$\begin{aligned} \kappa^+(y, x) &= \kappa^+(y(e), (e)x) \\ &= \kappa^+(y(e), (yy^{-1})x) \\ &= \kappa^+(e, y^{-1}x) \quad \text{by } (\dagger)^9 \end{aligned}$$

Now we can define our **one-parameter convolutional kernel** as

$$\kappa(y^{-1}x) := \kappa^+(e, y^{-1}x).$$

---

<sup>9</sup>Here we take the symbol  $y$  in this step to play the role of  $u$  in the LHS of  $(\dagger)$ .



# Why Should We Expect This? (Proof Sketch)

Lecture 3:  
Equivariant  
Convolutional  
Neural  
Networks on  
Homogeneous  
Spaces

$$\kappa(y^{-1}x) := \kappa^+(e, y^{-1}x) = \kappa^+(y, x).$$

Now for the map  $\phi : \Gamma(E_i) \cong \mathcal{I}_G^i \rightarrow \Gamma(E_{i+1}) \cong \mathcal{I}_G^{i+1}$ , we have that

$$\begin{aligned}(\phi f)(y) &= \int_G \kappa^+(y, x) f(x) d\mu_G(x) \\ &= \int_G \kappa(y^{-1}x) f(x) d\mu_G(x) \\ &= (\kappa \star f)(y).\end{aligned}$$

Thus,  $\phi$  can be written as a convolution!

CNNs with  
respect to  
Groups

Equivariant  
Neural  
Networks

Example:  
3D Steerable  
CNNs

A glimpse of  
implementations

## Question:

When can we represent  $\phi$  as an integral transform with  $\kappa^+$ ?

Representations of this sorts are usually linked to **Dunford-Pettis-like results** (recall Lecture 1), such as:

**Theorem [See Theorem 1.3 of [Arendt, 1994]]:**

Let  $\mathcal{K} : L^p(X) \rightarrow L^\infty(Y)$  be **linear, bounded operator** with  $1 \leq p < \infty$  then  $\mathcal{K}$  admits an integral representation.

The requirement for  $L^\infty$  in the codomain indicates that something might go wrong when  $X = Y$  if the domain is not **compact**.

- In practice, many ML researchers treat the ability to represent  $\phi$  as an **integral transform** as a **given assumption** (or what [Weiler et al., 2025] calls **ansatz**).
- We do ultimately want to optimize the parameters in a neural network layer. The integral representation gives a **more descriptive parameter** in terms of certain **matrix-valued functions**.
- For equivariant linear maps that do not admit the integral representation description, it is sometimes difficult to write down such linear maps.

From now on, we also adopt this **ansatz**.

# Bi-Equivariant Kernels

Lecture 3:  
Equivariant  
Convolutional  
Neural  
Networks on  
Homogeneous  
Spaces

CNNs with  
respect to  
Groups

Equivariant  
Neural  
Networks

Example:  
3D Steerable  
CNNs

A glimpse of  
implementations

Let  $\mathcal{H}$  be the collection of **equivariant linear maps**  $\mathcal{I}_G^i \rightarrow \mathcal{I}_G^{i+1}$ .  
There is an alternative characterization of  $\mathcal{H}$ :

**Theorem ([Cohen et al., 2019])**

$\mathcal{H}$  is isomorphic to the space of **bi-equivariant kernels**  
 $\kappa : G \rightarrow \text{Hom}(V_i, V_{i+1})$ , satisfying the condition

$$\kappa(h_2 g h_1) = \rho_{i+1}(h_2) \circ \kappa(g) \circ \rho_i(h_1).$$

Here recall  $\rho_{i+1}$  and  $\rho_i$  are the representations  
 $H_{i+1} \rightarrow \text{GL}(V_{i+1})$  and  $H_i \rightarrow \text{GL}(V_i)$  respectively.

# Left-Equivariant Kernels

Lecture 3:  
Equivariant  
Convolutional  
Neural  
Networks on  
Homogeneous  
Spaces

CNNs with  
respect to  
Groups

Equivariant  
Neural  
Networks

Example:  
3D Steerable  
CNNs

A glimpse of  
implementations

Although Mackey functions and sections are equivalent in this context,

- A **Mackey function** specifies a map  $f : G \rightarrow V$ .
- Whereas a **section** admits a map with domain being  $G/H$ .

In this perspective, a Mackey function contains **redundant information** and sections save more memories. We would therefore like a characterization of  $\mathcal{H}$  with  $G/H$ .

## Theorem ([Cohen et al., 2019])

$\mathcal{H}$  is isomorphic to the space of **left equivariant kernels**  $\overleftarrow{\kappa} : G/H_i \rightarrow \text{Hom}(V_i, V_{i+1})$  satisfying

$$\overleftarrow{\kappa}(h_2 x) := \rho_{i+1}(h_2) \circ \overleftarrow{\kappa}(x) \circ \rho_i(h_1(x, h_2)^{-1})$$

for all  $h_2 \in H_{i+1}, x \in G/H_i$ . Here  $h_1(x, g) := s(gx)^{-1}g$ , where  $s$  is a choice of local section.

# Kernels on the Double Coset

Lecture 3:  
Equivariant  
Convolutional  
Neural  
Networks on  
Homogeneous  
Spaces

CNNs with  
respect to  
Groups

Equivariant  
Neural  
Networks

Example:  
3D Steerable  
CNNs

A glimpse of  
implementations

Since there are two groups involved, we might as well consider the **double coset**:

$$H_{i+1} \backslash G / H_i := \text{right cosets of } H_{i+1} \text{ acting on } G / H_i.$$

**Theorem ([Cohen et al., 2019])**

$\mathcal{H}$  is isomorphic to the space  $\mathcal{K}_D$  comprising of functions

$$\bar{\kappa} : H_{i+1} \backslash G / H_i \rightarrow \text{Hom}(V_i, V_{i+1}),$$

$$\bar{\kappa}(x) = \rho_{i+1}(h) \bar{\kappa}(x) \rho_i^x(h)^{-1}, \forall x \in H_{i+1} \backslash G / H_i, h \in H_{i+1}^{\gamma(x)H_i}$$

Here  $\gamma : H_{i+1} \backslash G / H_i \rightarrow G$  is a choice of **coset representatives**:

- $H_{i+1}^{\gamma(x)H_i} = \{h \in H_{i+1} \mid h\gamma(x)H_i = \gamma(x)H_i\} \leq H_i.$
- $\rho_i^x(h) := \rho_i(\gamma(x)^{-1}h\gamma(x)).$

The **takeaway** is that this theorem is the most **memory-efficient way** to represent the data.

Lecture 3:  
Equivariant  
Convolutional  
Neural  
Networks on  
Homogeneous  
Spaces

CNNs with  
respect to  
Groups

Equivariant  
Neural  
Networks

Example:  
3D Steerable  
CNNs

A glimpse of  
implementations

- 1 CNNs with respect to Groups
- 2 Equivariant Neural Networks
- 3 Example: 3D Steerable CNNs**
- 4 A glimpse of implementations

# Example: 3D Steerable CNNs

Lecture 3:  
Equivariant  
Convolutional  
Neural  
Networks on  
Homogeneous  
Spaces

CNNs with  
respect to  
Groups

Equivariant  
Neural  
Networks

Example:  
3D Steerable  
CNNs

A glimpse of  
implementations

The work of **3D Steerable CNNs** [Weiler et al., 2018] is concerned with the group  $\text{SE}(3)$  of **orientation preserving rigid body motions** (ie. rotations and translations) in  $\mathbb{R}^3$ . This is called the **special Euclidean group** for  $\mathbb{R}^3$ .

In their work, the layer is built with the choice

$$G = \text{SE}(3) \text{ and } H = H_1 = H_2 = \text{SO}(3).$$

and representations  $\rho_1$  and  $\rho_2$ .


Note that  $G$  can be written as a certain **twisted product**<sup>10</sup> of  $\text{SO}(3)$  and  $\mathbb{R}^3$ .

---

<sup>10</sup>To be more precise, a semi-direct product.



# Example: 3D Steerable CNNs



Picture from [Cohen et al., 2019].

Note in this diagram  $G/H_1 \cong \mathbb{R}^3$  and  $H_2 \backslash G/H_1 \cong [0, \infty)$ .

## Theorem

The equivariant linear maps  $\mathcal{H}$  can be identified with left-equivariant kernels  $\overleftarrow{\kappa}$  such that

$$\overleftarrow{\kappa}(rx) = \rho_2(r) \circ \overleftarrow{\kappa}(x) \circ \rho_1(r^{-1}).$$

# Example: 3D Steerable CNNs

## Theorem

The equivariant linear maps  $\mathcal{H}$  can be identified with **left-equivariant kernels**  $\overleftarrow{\kappa}$  such that

$$\overleftarrow{\kappa}(rx) = \rho_2(r) \circ \overleftarrow{\kappa}(x) \circ \rho_1(r^{-1}), r \in \text{SO}(3), x \in \mathbb{R}^3.$$

By the general characterization in the last section, we have that

$$\overleftarrow{\kappa}(rx) = \rho_2(r) \circ \overleftarrow{\kappa}(x) \circ \rho_1(h_1(x, r)^{-1}),$$

and  $h_1(x, r) := s(rx)^{-1}r$  for  $s$  some local section. In this case, since  $\text{SE}(3)$  is a twisted product of  $\mathbb{R}^3$  and  $\text{SO}(3)$ , it is actually a **trivial principal  $H$ -bundle**, so we can actually choose  $s$  to be the identity section! Thus

$$h_1(x, r)^{-1} = r^{-1}.$$

Lecture 3:  
Equivariant  
Convolutional  
Neural  
Networks on  
Homogeneous  
Spaces

CNNs with  
respect to  
Groups

Equivariant  
Neural  
Networks

Example:  
3D Steerable  
CNNs

A glimpse of  
implementations

- 1 CNNs with respect to Groups
- 2 Equivariant Neural Networks
- 3 Example: 3D Steerable CNNs
- 4 A glimpse of implementations

# What About Practical Implementations?

Lecture 3:  
Equivariant  
Convolutional  
Neural  
Networks on  
Homogeneous  
Spaces

CNNs with  
respect to  
Groups

Equivariant  
Neural  
Networks

Example:  
3D Steerable  
CNNs

A glimpse of  
implementations

Up to now we have developed the theory; now we will discuss some implementations.

Practical implementations ask: How do we discretize a (possibly continuous) group, parameterize kernels, and execute the resulting tensor operations efficiently?

Now we will briefly survey some works that have already translated  $G$ -convolution theory into code.

[e2cnn](#) [Weiler and Cesa, 2019] is a PyTorch add-on that builds  $E(2)$ -equivariant CNN layers whose kernels are automatically parameterized to rotations and reflections.

[Gupta et al., 2021] gives a [rotation equivariant siamese network for tracker](#) that estimates in-plane pose and lifts the SOTA on the Rotating-Object Benchmark.

# Examples

**e3nn** [Geiger and Smidt, 2022] is also a PyTorch library providing  $E(3)$ -equivariant convolutional operators via steerable filters built from irreducible  $SO(3)$  representations.

**escnn** [Cesa et al., 2022] is a successor to e2cnn that proposes a general procedure to build arbitrary CNNs with respect to any compact group  $G$ .

- In practice, they considered a way to construct  $G$ -steerable (equivariant) kernels with any  $G \leq O(3)$ .
- This gave a way to build CNNs with respect to symmetries of **platonic solids** or choosing  $G = SO(2)$  in 3D to only have **azimuthal symmetries**.
- Achieve SOTA on volumetric datasets ModelNet10 [Wu et al., 2015], a rotated version of it, and LBA [Townshend et al., 2022].



Arendt, Wolfgang, B. A. V. (1994).

Integral representations of resolvents and semigroups.  
*Forum mathematicum*, 6(1):111–136.



Cesa, G., Lang, L., and Weiler, M. (2022).

A program to build  $e(N)$ -equivariant steerable cnns.  
International Conference on Learning Representations  
(ICLR) 2022.



Cohen, T. S., Geiger, M., and Weiler, M. (2019).

*A general theory of equivariant CNNs on homogeneous spaces.*

Curran Associates Inc., Red Hook, NY, USA.



Geiger, M. and Smidt, T. E. (2022).

e3nn: Euclidean neural networks.



Gupta, D. K., Arya, D., and Gavves, E. (2021).  
Rotation-equivariant siamese networks for tracking.



Townshend, R. J. L., Vögele, M., Suriana, P., Derry, A.,  
Powers, A., Laloudakis, Y., Balachandar, S., Jing, B.,  
Anderson, B., Eismann, S., Kondor, R., Altman, R. B., and  
Dror, R. O. (2022).  
Atom3d: Tasks on molecules in three dimensions.



Weiler, M. and Cesa, G. (2019).  
General  $e(2)$ -equivariant steerable cnns.



Weiler, M., Forré, P., Verlinde, E., and Welling, M. (2025).  
*Equivariant and coordinate independent convolutional  
networks.*  
WORLD SCIENTIFIC.





Weiler, M., Geiger, M., Welling, M., Boomsma, W., and Cohen, T. S. (2018).

3d steerable cnns: Learning rotationally equivariant features in volumetric data.

In Bengio, S., Wallach, H., Larochelle, H., Grauman, K., Cesa-Bianchi, N., and Garnett, R., editors, *Advances in Neural Information Processing Systems*, volume 31. Curran Associates, Inc.



Wu, Z., Song, S., Khosla, A., Yu, F., Zhang, L., Tang, X., and Xiao, J. (2015).

3d shapenets: A deep representation for volumetric shapes.