

Assignment 2.

Jithin D. George, No. 1622555

Due Monday, Jan. 22.

1. (Inverse matrix and Green's functions)

- (a) Write out the 5×5 matrix A from (2.43) for the boundary value problem $u''(x) = f(x)$ with $u(0) = u(1) = 0$ for $h = 0.25$.

Solution:

```
def f(x):
    return x
```

$$A = \begin{bmatrix} 1. & 0. & 0. & 0. & 0. \\ 0.25 & -0.5 & 0.25 & 0. & 0. \\ 0. & 0.25 & -0.5 & 0.25 & 0. \\ 0. & 0. & 0.25 & -0.5 & 0.25 \\ 0. & 0. & 0. & 0. & 1. \end{bmatrix}$$

- (b) Write out the 5×5 inverse matrix A^{-1} explicitly for this problem.

Solution:

$$A^{-1} = \begin{bmatrix} 1. & -0. & -0. & -0. & 0. \\ 0.75 & -0.046875 & -0.03125 & -0.015625 & 0.25 \\ 0.5 & -0.03125 & -0.0625 & -0.03125 & 0.5 \\ 0.25 & -0.015625 & -0.03125 & -0.046875 & 0.75 \\ 0. & 0. & 0. & 0. & 1. \end{bmatrix}$$

- (c) If $f(x) = x$, determine the discrete approximation to the solution of the boundary value problem on this grid and sketch this solution and the five Green's functions whose sum gives this solution.

Solution:

$$u(x) = \int_0^1 \bar{x} G(x; \bar{x}) d\bar{x} = 0.25hG_1 + 0.5hG_2 + 0.75hG_3$$




Figure 1: Interior Green's functions, Boundary Green's functions and Solution

2. (Another way of analyzing the error using Green's functions) The *composite trapezoid rule* for integration approximates the integral from a to b of a function g by dividing the interval into segments of length h and approximating the integral over each segment by the integral of the linear function that matches g at the endpoints of the segment. (For $g > 0$, this is the area of the trapezoid with height $g(x_j)$ at the left endpoint x_j and height $g(x_{j+1})$ at the right endpoint x_{j+1} .) Letting $h = (b - a)/(m + 1)$ and $x_j = a + jh$, $j = 0, 1, \dots, m, m + 1$:

$$\int_a^b g(x) dx \approx h \sum_{j=0}^m \frac{g(x_j) + g(x_{j+1})}{2} = h \left[\frac{g(x_0)}{2} + \sum_{j=1}^m g(x_j) + \frac{g(x_{m+1})}{2} \right].$$

- (a) Assuming that g is sufficiently smooth, show that the error in the composite trapezoid rule approximation to the integral is $O(h^2)$. [Hint: Show that the error on each subinterval is $O(h^3)$.]

Solution:

$$\begin{aligned} g(x) &= g(x_i) + g'(x_i)(x - x_i) + \frac{1}{2}g''(x_i)(x - x_i)^2 + O((x - x_i)^3) \\ \int_{x_i}^{x_{i+1}} g(x) dx &= \int_{x_i}^{x_{i+1}} g(x_i) dx + \int_{x_i}^{x_{i+1}} g'(x_i)(x - x_i) dx + O((x_{i+1} - x_i)^3) \\ &= g(x_i)(x_{i+1} - x_i) + g'(x_i)(x_{i+1} - x_i)^2 + O((x_{i+1} - x_i)^3) \\ \int_{x_i}^{x_{i+1}} g(x) dx &= g(x_i)h + g'(x_i)\frac{h^2}{2} + O(h^3) \end{aligned} \quad (1)$$

$$\begin{aligned} g(x) &= g(x_{i+1}) + g'(x_{i+1})(x - x_{i+1}) + \frac{1}{2}g''(x_{i+1})(x - x_{i+1})^2 + O((x - x_{i+1})^3) \\ g(x) &= g(x_{i+1}) + g'(x_{i+1})(x - x_{i+1}) + g''(x_{i+1})h(x - x_{i+1}) + \frac{1}{2}g''(x_{i+1})(x - x_{i+1})^2 + O((x - x_{i+1})^3) \\ \int_{x_i}^{x_{i+1}} g(x) dx &= \int_{x_i}^{x_i} g(x_{i+1}) dx + \int_{x_i}^{x_{i+1}} g'(x_{i+1})(x - x_{i+1}) + O((x_{i+1} - x_i)^3) \\ \int_{x_i}^{x_{i+1}} g(x) dx &= g(x_{i+1})h - g'(x_{i+1})\frac{h^2}{2} + O(h^3) \end{aligned} \quad (2)$$

Adding (1) and (2) together,

$$\int_{x_i}^{x_{i+1}} g(x) dx = \frac{h}{2}(g(x_i) + g(x_{i+1})) + O(h^3)$$

- (b) Recall that the true solution of the boundary value problem $u''(x) = f(x)$, $u(0) = u(1) = 0$ can be written as

$$u(x) = \int_0^1 f(\bar{x})G(x; \bar{x}) d\bar{x}, \quad (3)$$

where $G(x; \bar{x})$ is the Green's function corresponding to \bar{x} . The finite difference approximation u_i to $u(x_i)$, using the centered finite difference scheme in (2.43), is

$$u_i = h \sum_{j=1}^m f(x_j)G(x_i; x_j), \quad i = 1, \dots, m. \quad (4)$$

Show that formula (4) is the trapezoid rule approximation to the integral in (3) when $x = x_i$, and conclude from this that the error in the finite difference approximation is $O(h^2)$ at each node x_i . [Recall: The Green's function $G(x; x_j)$ has a *discontinuous* derivative at $x = x_j$. Why does this not degrade the accuracy of the composite trapezoid rule?]

Solution:

$$u(x) = \int_0^1 f(\bar{x})G(x; \bar{x}) d\bar{x}$$

Setting $x = x_i$ and using the Trapezoidal approximation,

$$u_i = h \frac{f(0)G(x_i; 0)}{2} + h \sum_{j=1}^m f(x_j)G(x_i; x_j) + h \frac{f(1)G(x_i; 1)}{2}$$

Since the Green's function is the solution to the ODE and since at the boundaries, u is 0, $G(x,0)=G(x,1)=0$. So, we get

$$u_i = h \sum_{j=1}^m f(x_j)G(x_i; x_j)$$

3. (Green's function with Neumann boundary conditions)

- (a) Determine the Green's functions for the two-point boundary value problem $u''(x) = f(x)$ on $0 < x < 1$ with a Neumann boundary condition at $x = 0$ and a Dirichlet condition at $x = 1$, i.e, find the function $G(x, \bar{x})$ solving

$$u''(x) = \delta(x - \bar{x}), \quad u'(0) = 0, \quad u(1) = 0$$

and the functions $G_0(x)$ solving

$$u''(x) = 0, \quad u'(0) = 1, \quad u(1) = 0$$

and $G_1(x)$ solving

$$u''(x) = 0, \quad u'(0) = 0, \quad u(1) = 1.$$

Solution:

We need

$$u'(0) = 0, u(1) = 0$$

and

$$u(\bar{x} + \epsilon) - u(\bar{x} - \epsilon) = 1$$

So, the Green's function satisfying that would be given by

$$G(x, \bar{x}) = \begin{cases} \bar{x} - 1, & \text{for } 0 \leq x \leq \bar{x} \\ x - 1, & \text{for } \bar{x} \leq x \leq 1 \end{cases}$$

Solving the odes for the boundaries, we have

$$G_0(x) = x - 1$$

$$G_1(x) = 1$$

- (b) Using this as guidance, find the general formulas for the elements of the inverse of the matrix in equation (2.54). Write out the 5×5 matrices A and A^{-1} for the case $h = 0.25$.

Solution:

$$A = \begin{bmatrix} 3. & -4. & 1. & 0. & 0. \\ 0.25 & -0.5 & 0.25 & 0. & 0. \\ 0. & 0.25 & -0.5 & 0.25 & 0. \\ 0. & 0. & 0.25 & -0.5 & 0.25 \\ 0. & 0. & 0. & 0. & 1. \end{bmatrix}$$

$$A^{-1} = \begin{bmatrix} -1. & -0.75 & -0.5 & -0.25 & 1. \\ -0.75 & -0.75 & -0.5 & -0.25 & 1. \\ -0.5 & -0.5 & -0.5 & -0.25 & 1. \\ -0.25 & -0.25 & -0.25 & -0.25 & 1. \\ 0. & 0. & 0. & 0. & 1. \end{bmatrix}$$

4. (Solvability condition for Neumann problem) Determine the null space of the matrix A^T , where A is given in equation (2.58), and verify that the condition (2.62) must hold for the linear system to have solutions.

Solution:

$$A = \begin{pmatrix} -h & h & & & \\ 1 & -2 & 1 & & \\ & 1 & \ddots & \ddots & \\ & & \ddots & \ddots & 1 \\ & & & 1 & -2 & 1 \\ & & & & h & -h \end{pmatrix}$$

$$A^T = \begin{pmatrix} -h & 1 & & & \\ h & -2 & 1 & & \\ & 1 & \ddots & \ddots & \\ & & \ddots & \ddots & 1 \\ & & & 1 & -2 & h \\ & & & & 1 & -h \end{pmatrix}$$

The nullspace of this matrix is the vector $[1, h, h, h, h, \dots, h, 1]$ because we can see multiplying them yields 0. So, a solution only exists if the f-vector is orthogonal to the nullspace. Setting the dot-product equal to 0, we obtain the following criteria

$$\sigma_0 + (h/2)f(x_0) + hf(x_1) + \dots + hf(x_{m-1}) + hf(x_m) - \sigma_1 + \frac{h}{2}f(x_{m+1}) = 0$$

which is the same as 2.62.

5. (Symmetric tridiagonal matrices)

- (a) Consider the **Second approach** described on p. 31 for dealing with a Neumann boundary condition. If we use this technique to approximate the solution to the boundary value problem $u''(x) = f(x)$, $0 \leq x \leq 1$, $u'(0) = \sigma$, $u(1) = \beta$, then the resulting linear system $A\mathbf{u} = \mathbf{f}$ has the following form:

$$\frac{1}{h^2} \begin{pmatrix} -h & h & & & \\ 1 & -2 & 1 & & \\ & 1 & \ddots & \ddots & \\ & & \ddots & \ddots & 1 \\ & & & 1 & -2 \end{pmatrix} \begin{pmatrix} u_0 \\ u_1 \\ \vdots \\ u_{m-1} \\ u_m \end{pmatrix} = \begin{pmatrix} \sigma + (h/2)f(x_0) \\ f(x_1) \\ \vdots \\ f(x_{m-1}) \\ f(x_m) - \beta/h^2 \end{pmatrix}.$$

Show that the above matrix is similar to a symmetric tridiagonal matrix via a *diagonal* similarity transformation; that is, there is a diagonal matrix D such that DAD^{-1} is symmetric.

Solution: Experimenting with a smaller version,

$$\begin{pmatrix} d_0 & 0 & 0 \\ 0 & d_1 & 0 \\ 0 & 0 & d_2 \end{pmatrix} \begin{pmatrix} -h & h & 0 \\ 1 & -2 & 1 \\ 0 & 1 & -2 \end{pmatrix} \begin{pmatrix} \frac{1}{d_0} & 0 & 0 \\ 0 & \frac{1}{d_1} & 0 \\ 0 & 0 & \frac{1}{d_2} \end{pmatrix} = \begin{pmatrix} -h & h\frac{d_0}{d_1} & 0 \\ \frac{d_1}{d_0} & -2 & \frac{d_1}{d_2} \\ 0 & \frac{d_2}{d_1} & -2 \end{pmatrix}$$

So, for the resulting matrix to be symmetric, we choose the diagonal vector given by $[1 \ \sqrt{h} \ \sqrt{h} \ \sqrt{h} \ \sqrt{h}]$

$$\begin{pmatrix} 1 & & & & \\ & \sqrt{h} & & & \\ & & \ddots & & \\ & & & \ddots & \\ & & & & \sqrt{h} \end{pmatrix} \begin{pmatrix} -h & h & & & \\ 1 & -2 & 1 & & \\ & 1 & \ddots & \ddots & \\ & & \ddots & \ddots & 1 \\ & & & 1 & -2 \end{pmatrix} \begin{pmatrix} 1 & & & & \\ & \frac{1}{\sqrt{h}} & & & \\ & & \ddots & & \\ & & & \ddots & \\ & & & & \frac{1}{\sqrt{h}} \end{pmatrix} \\ = \begin{pmatrix} -h & \sqrt{h} & & & \\ \sqrt{h} & -2 & 1 & & \\ & 1 & \ddots & \ddots & \\ & & \ddots & \ddots & 1 \\ & & & 1 & -2 \end{pmatrix}$$

This is a symmetric matrix.

- (b) Consider the **Third approach** described on pp. 31-32 for dealing with a Neumann boundary condition. [**Note:** If you have an older edition of the text, there is a typo in the matrix (2.57) on p. 32. There should be a row above what is written there that has entries $\frac{3}{2}h$, $-2h$, and $\frac{1}{2}h$ in columns 1 through 3 and 0's elsewhere. I believe this was corrected in newer editions.] Show that if we use that first equation (given at the bottom of p. 31) to eliminate u_0 and we also eliminate u_{m+1} from the equations by setting it equal to β and modifying the right-hand side vector accordingly, then we obtain an m by m linear system $A\mathbf{u} = \mathbf{f}$, where A is similar to a symmetric tridiagonal matrix via a diagonal similarity transformation.

Solution:

$$\frac{1}{h^2} \begin{pmatrix} \frac{3}{2}h & -2h & \frac{1}{2}h & & \\ 1 & -2 & 1 & & \\ & 1 & \ddots & \ddots & \\ & & \ddots & \ddots & 1 \\ & & & 1 & -2 \end{pmatrix} \begin{pmatrix} u_0 \\ u_1 \\ \vdots \\ u_{m-1} \\ u_m \end{pmatrix} = \begin{pmatrix} \sigma \\ f(x_1) \\ \vdots \\ f(x_{m-1}) \\ f(x_m) - \beta/h^2 \end{pmatrix}.$$

$$\frac{1}{h^2} \begin{pmatrix} 3 & -4 & 1 & & \\ 3 & -6 & 3 & & \\ & 1 & \ddots & \ddots & \\ & & \ddots & \ddots & 1 \\ & & & 1 & -2 \end{pmatrix} \begin{pmatrix} u_0 \\ u_1 \\ \vdots \\ u_{m-1} \\ u_m \end{pmatrix} = \begin{pmatrix} \frac{2\sigma}{h} \\ 3f(x_1) \\ \vdots \\ f(x_{m-1}) \\ f(x_m) - \beta/h^2 \end{pmatrix}.$$

Eliminating the boundary element,

$$\frac{1}{h^2} \begin{pmatrix} -1 & 1 & & & \\ 1 & \ddots & \ddots & & \\ & \ddots & \ddots & 1 & \\ & & & 1 & -2 \end{pmatrix} \begin{pmatrix} u_1 \\ \vdots \\ u_{m-1} \\ u_m \end{pmatrix} = \begin{pmatrix} \frac{3}{2}f(x_1) - \frac{\sigma}{h} \\ \vdots \\ f(x_{m-1}) \\ f(x_m) - \beta/h^2 \end{pmatrix}.$$

This is a symmetric matrix.