

Summary

Dis·count

August 28, 2020

Introduction

At first, we will introduce some preliminary knowledge about cooperative game theory as follows. A pair (V, c) is usually used to represent a cooperative game with transferable utilities, among which $V = \{1, 2, \dots, v\}$ denotes a set of v players and $c : 2^V \rightarrow \mathbb{R}$ indicates the characteristic function of the game. A coalition is defined as a non-empty subset of players; while V is referred to as a coalition, $S = 2^V \setminus \{\emptyset\}$ denotes the set of all feasible coalitions. The characteristic function of the game, $c(s)$, represents the minimum coalitional cost the players in s have to pay in order to cooperate together. A cost allocation vector $\alpha = [\alpha_1, \alpha_2, \dots, \alpha_v] \in \mathbb{R}^v$ is required by the game to maintain cooperation in the grand coalition, and α_k is the cost assigned to players $k \in V$ to assure no individual or group of players has the incentive to deviate. For the convenience of abbreviation, we use $\alpha(s) = \sum_{k \in s} \alpha_k$ to denote the total cost assigned to the players in coalition s . One of the most important concepts in cooperative game theory is core, which is a cost allocation satisfying two kinds of constraints, one is the budget balance constraint $\alpha(V) = c(V)$ and the other is the coalitional stability constraints $\alpha(s) \leq c(s)$. In other words, core can be expressed as

$$\text{Core}(V, c) = \{\alpha : \alpha(V) = c(V), \alpha(s) \leq c(s) \text{ for all } s \in S \setminus \{V\}, \alpha \in \mathbb{R}^v\}.$$

When the cost allocation exists, core is called non-empty. And if and only if the core is non-empty, the grand coalition of the associating cooperative game (V, c) will be stable or balanced.

However, cooperative games can be unbalanced in many cases owing to the joint restrictions of the above-mentioned two kinds of constraints. To stabilize the grand coalitions in unbalanced cooperative games, researchers have already developed several effective instruments, such as subsidization, penalization and simultaneous subsidization and penalization. The similarity of these instruments is that there exists an outside party who will take measures to stabilize the grand coalition. But the penalization will always arouse the discontent of players in coalitions, it's promising to find a new instrument to replace the penalization, we call it taxation. The significant idea of this instrument is that we can erase the role of the third party by collecting taxation as subsidy of the corresponding coalitions. That being said, the players of the games can stabilize themselves without the third party.

To make the project concrete, we will apply the cooperative theory on the machine schedule problem to

show our ideas.

Definition 0.

In an Identical Variable Parallel machine scheduling of Unweighted jobs (IVPU) game, each player k in $V = \{1, 2, \dots, v\}$ has a job k that needs to be processed on one of identical machines in $M = \{1, 2, \dots, m_v\}$, where m_v is a given positive integer no more than v . Meanwhile, each machine has a setup cost T . Each job $k \in V$ has a processing time denoted by t_k . Each coalition $s \in S$, where $S = 2^V \setminus \{\emptyset\}$, aims to schedule the jobs in s on all machines in M so that the total completion time of the jobs in s is minimized, i.e., to minimize $\sum_{k \in s} C_k + m_s T$, where C_k is the completion time of job $k \in s$.

Then we will illustrate the taxation instrument with a simple example of an IAPU game as follows.

Problem Description

There is an IPU game which contains four players, whose processing times on the identical parallel machine are $t_1 = 2, t_2 = 3, t_3 = 4, t_4 = 5$ respectively. Each machine setup cost is $t_0 = 9.5$, and $c(s)$ is the minimum total completion time of jobs in coalition s plus the machine setup cost.

In this example, the grand coalition has a minimum cost, which is $c(V) = \sum_{k \in v} C_k + m_v T = 38$, and the optimum number of machine is $m_v = 2$. It's easy to check that this example is unbalanced and calculate the subsidy which equals $c(V) - \alpha(V) = 0.75$.

When the setup cost increases from 9.5 to 13, that is, the taxation increases from 0 to 3.5, the number of machine will decrease from 2 to 1, accordingly.

Although this case is easily understood, it demonstrate the concept of taxation we have strong interests in. Then we will define the corresponding model to further elaborate on this concept.

Definition 1.

A cooperative TU game (V, c) is called an IVPU game if it satisfies the following formulations.

$$\begin{aligned}
c(s, P) = & \min \sum_{k \in V} \sum_{j \in O} c_{kj} x_{kj} + P \sum_{k \in s} x_{k1} \\
s.t. \quad & \sum_{j \in O} x_{kj} - y_k^s = 0, \forall k \in V, \\
& \sum_{k \in V} x_{kj} \leq m, \forall j \in O, \\
& x_{kj} \in \{0, 1\}, \forall k \in V, \forall j \in O, \\
& y_k^s = 1, k \in s, y_k^s = 0, k \notin s.
\end{aligned}$$

Among these formulations, P and m denotes the price and the number of using machines, respectively. Now we've already extended the number of players to a more complex case, that is, the number of players is set to

n . Let $N = \{1, 2, \dots, n\}$ be a set of n players. The number of machines is m and the setup cost is P . For convenience of expression, we set the processing times $t_i (i \in N)$ satisfy $t_1 < t_2 < \dots < t_n$.

Then we need to define two functions, practical subsidy-price function (PSPF) and theoretical subsidy-price function (TSPF) denoted by $\omega(P)$ and $\hat{\omega}(P)$, respectively:

$$\omega(P) = \min_{\alpha} \{c(V, m(V, P)) - \alpha(V) : \alpha(s) \leq c(s, m(s, P)), \forall s \in S, \alpha \in \mathbb{R}^v\},$$

$$\hat{\omega}(P) = \min_{\alpha} \{c(V, m(V, P)) - \alpha(V) : \alpha(s) \leq c(s, m(s, P)), \forall s \in S \setminus \{V\}, \alpha \in \mathbb{R}^v\}.$$

By dividing the PSPF function into two parts like $c(s) = c_0(s) + m(s)P, s \in S$, as a common skill, we can obtain its dual:

$$\omega^*(P) = \max_{\rho} \{c_0(V) + m(V)P + \sum_{s \in S \setminus \{V\}} -\rho_s [c_0(s) + m(s)P] : \sum_{s \in S \setminus \{V\} : k \in s} \rho_s = 1, \forall k \in V, \rho_s \geq 0, \forall s \in S \setminus V\} \quad (1)$$

Then we need to define an interval $[P_L(m, s), P_H(m, s)]$ to denote the value range of price when the number of using machines m and the coalition s are given.

Remark 1. It's easy to know that $P_L(m, s) = P_H(m + 1, s)$, for each $s \in S$ and $m < |s|$.

Remark 2. For the grand coalition V , we define $P_L(v, V) = 0, P_H(1, V) = P^*$. So the practical domain of the price is $[0, P^*]$ which is divided into v parts by the number of using machines.

Remark 3. Note that $P_L(m, s) = P_H(m + 1, s)$, for the sake of brevity and readability, we use P_{m+1} to substitute for $P_L(m, s)$ or $P_H(m + 1, s)$ later on.

We can establish Theorem 1 below, showing the property of the function.

Theorem 1. $\omega(P)$ is piecewise linear, and convex in price P at each subinterval $[P_L(m, V), P_H(m, V)]$ in $[0, P^*]$.

In this situation, the interval size of setup cost can be calculated when t_i is known. And we can obtain that under what circumstances the machine number will change. We set the different intervals where the number of machines remain unchanged as I_i respectively. The extreme points of these intervals are recorded as P_i respectively, which represents the price when the number of using machine changes from i to $i - 1$. Specially, P_1 denotes the extreme point of the whole interval, which is the value of price or setup cost when using one machine and subsidy equals zero.

Note that the costs of all the exponential coalitions can be easily calculated by the SPT rules, we have Lemma 1.

Lemma 1. The value at the extreme points of the sub-intervals I_i can be calculated with processing times t_i by comparing the costs of the grand coalitions where all the players use two adjacent numbers of machines.

According to the above lemma 1, we can obtain all the extreme points of the sub-intervals, i.e., the number of using machines decreases by one when the price or setup cost equals P_i .

Theorem 2. According to the foregoing description, we have the equation $P_1 = P_2 + \dots + P_n = \sum_{i=2}^n P_i$.

With Theorem 2, we obtain all the breakpoints during the interval of the price where the machine number changes and the subsidy is 0. Then we'll focus on the specific property of subsidy.

Theorem 3. $\omega(P)$ is hard to calculate, however, $\omega(P)$ can be bounded by zero when the number of using machines, m , is larger than $\frac{n}{2}$.

In other words, when the numbers of machine is larger than half of players, the grand coalition doesn't need any subsidy from the externality.

Theorem 4. The sum of absolute values of slopes at both sides of S_i is 1. When the number of machines used is 1, the range of slopes of the line segments in the interval is $\left(-1, -\frac{1}{n-1}\right]$, and the number of breakpoints is $O(v^2)$.

Until here, we described the main property of the whole figure. And a diagram of the number of machines and subsidy on setup cost, with its essential features being showed below.




Figure 1: The number of machines and subsidy on setup cost.

The processing times of all players with the arrangement from small to large are listed on the top of this figure. The red and blue lines stand for the machine number and subsidy, respectively. The horizontal ordinate represents the price or setup cost. The left ordinate represents the value of machine number while the right represents the value of subsidy.

As the coalitional stability constraints showed above, the exponential inequality constraints are so tricky that we must figure out a method to simplify these constraints to obtain the optimal results. Fortunately for us, we find the Theorem 5.

Theorem 5. *Define that*

$$\omega_1(P) = \min_{\alpha} \{c(V, m(V, P)) - \alpha(V) : \alpha(s) \leq c(s) + P, \forall s \in S, \alpha \in \mathbb{R}^v\},$$

then the original problem $\omega(P)$ is equivalent to $\omega_1(P)$ which means that all sub-coalitions only use one machine.

By establishing the Theorem 5, we can design the specific algorithm to eliminate the redundant inequality, which will save us from the trouble of solving $c(s)$ and speed up the solution.

At first, we need to construct the PSPF function, which is piecewise linear and convex at every subinterval. If we can obtain the specific subsidy value $\omega(P)$ given any P , then we can calculate the breakpoints during the interval by IPC algorithm.

At first, we need to divide the interval into n subintervals according to the number of using machine of the grand coalition, at each subinterval we use the IPC algorithm to get the breakpoints. After applying the algorithm to all subintervals, we can obtain the PSPF function.

Algorithm 1 The Intersection Points Computation(IPC) Algorithm to Construct the PSPF Function.

Step 1. Initially, set $I^* = \{P_L, P_H\}$ and $\mathbb{I} = \{[P_L, P_H]\}$.

Step 2. If \mathbb{I} is not empty, update I^* and \mathbb{I} by the following steps:

Step 3. Sort values in I^* by $P_0 < P_1 < \dots < P_q$, where $P_0 = P_L, P_q = P_H$ and $q = |I^*| - 1$.

Step 4. Select any interval from \mathbb{I} , denoted by $[P_{k-1}, P_k]$ with $1 \leq k \leq q$

Step 5. Construct two linear function $R_{k-1}(P)$ and $L_k(P)$ so that $R_{k-1}(P)$ passes $(P_{k-1}, \omega(P_{k-1}))$ with a slope equal to a right derivative $K_r^{P_{k-1}}$ of $\omega(P)$ at P_{k-1} , and that $L_k(z)$ passes $(P_k, \omega(P_k))$ with a slope equal to a left derivative $K_l^{P_k}$ of $\omega(P)$ at P_k .

Step 6. If $R_{k-1}(P)$ passes $(P_k, \omega(P_k))$ or $L_k(P)$ passes $(P_{k-1}, \omega(P_{k-1}))$, then update \mathbb{I} by removing $[P_{k-1}, P_k]$. Otherwise, $R_{k-1}(P)$ and $L_k(P)$ must have a unique intersection point at $P = P'$ for some $P' \in (P_{k-1}, P_k)$. Update I^* by adding P' , and update \mathbb{I} by removing $[P_{k-1}, P_k]$, adding $[P_L, P']$ and $[P', P_r]$.

Step 7. Go to step 2.

Step 8. Return a piecewise linear function by connecting points $(P, \omega(P))$ for all $P \in I^*$.

Then we need to calculate the $\omega(P)$ when given an arbitrary P . We can follow the basic Cutting Plane(CP) algorithm, the specific process are represented in algorithm 2.

As we all know, the cost arised from the partial players in the grand coalition, that is $c(s)$, can be calculated handily by the SPT rule (The corresponding conclusion see Lemma 1). Meanwhile, inspired by the paper (Please

Algorithm 2 The Cutting Plane(CP) Algorithm to compute $\omega(P)$ for a given P .

Step 1. Let $\mathbb{S}' \subseteq \mathbb{S} \setminus \{N\}$ indicates a restricted coalition set, which includes some initial coalitions, e.g., $\{1\}, \{2\}, \dots, \{v\}$.

Step 2. Find an optimal solution $\bar{\alpha}(\cdot, P)$ to LP $\tau(P)$:

$$\max_{\alpha \in \mathbb{R}^n} \{ \alpha(N, P) : \alpha(s, P) \leq c(s) + P, \text{ for all } s \in \mathbb{S}' \}.$$

Step 3. Find an optimal solution s^* to the separation problem:

$$\delta = \min \{ c(s) + P - \bar{\alpha}(s, z) : \forall s \in \mathbb{S} \setminus \{N\} \}.$$

Step 4. If $\delta < 0$, then add s^* to \mathbb{S}' , and go to step 2; otherwise, return $\omega(P) = c(N) - \bar{\alpha}(N, P)$ and the pair of derivatives $(K_l^{\bar{\beta}}, K_r^{\bar{\beta}})$.

refer to Liu et.al.2018), we have the following approach to solve this game.

There is an effective domain $[0, P^*]$, where the grand coalition may need a subsidy from the external to maintain the balance. (The corresponding conclusion see Theorem 1 and 2 where $P^* = P_L(V, 1)$).

For the effective domain, we just need to divide this interval into several parts and the breakpoints are denoted as $P_L(V, i), i = 1, 2, \dots, n$. At first, we don't need to calculate the initial part because at this part the corresponding subsidy is 0 always. (The corresponding conclusion see Theorem 3) Then we just need to focus on the latter part which shows some interesting properties we present above.

As we've already known that $P_L(V, i), i = 1, 2, \dots, n$ can be obtained by Lemma 1 and Theorem 2, we just need to follow the CP approach (Algorithm 3 in Liu et.al.2018) to calculate the weak derivatives at each sub-interval $[P_L(m, V), P_H(m, V)]$ where the corresponding derivative is $m_V - \sum_{s \in \mathbb{S} \setminus \{V\}} \rho_s$.

Then use IPC Algorithm which will return all the breakpoints during the $[0, P^*]$ to obtain the subsidy $\omega(P)$.

Notice that we can calculate the characteristic function $c(s)$ easily according to Theorem 5, we can formulate Theorem 6.

Lemma 2. *For the IVPU game, the corresponding separation problem can be solved in $O(v^2)$ time, which could be shown in the following DP algorithm.*

Now we know the separation problem can be solved in $O(v^2)$ time and the breakpoints during the whole interval are polynomial, so for the IVPU game the PSPF function can be constructed in polynomial time.

Theorem 6. *The IVPU game can be solved in polynomial time.*

Until here, we've solved the IVPU game with alternative machines.

Then, we will try to extend the core concept to a more general case. So we need to define a new game as follow.

Algorithm 3 The Dynamic Programming(DP) Algorithm to Solve the separation problem.

Step 1. Initially, let $D(k, u)$ indicate the minimum objective value of the restricted problem of separation problem, where $k \in \{1, 2, \dots, v\}$ and $u \in \{0, 1, \dots, v\}$.

Step 2. Given the initial conditions $D(1, 0) = P$ and $D(1, 1) = t_1 - \beta_1 + P$. The boundary conditions are $D(k, u) = \text{if } u > k, \text{ for all } k \in V$.

Step 3. Given the recursion:

$$D(k, u) = \min \begin{cases} D(k-1, u), \text{ for the case when } s^* \text{ does not contain } k, \\ D(k-1, u-1) + ut_k - \alpha_k, \text{ for the case when } s^* \text{ contains } k. \end{cases}$$

Step 4. Obtain the optimal objective value of separation problem by $\delta_{IVPU} = \min\{D(v, u) : u \in \{1, 2, \dots, v-1\}\}$. return δ_{IVPU} .

Definition 2

A cooperative TU game (V, c) is a PIM(Price Integer Minimization) game if there exist:

- positive integers e, v, t, P and m ;
- integer vectors $x \in \mathbb{Z}^{t \times 1}$ and $\tilde{\alpha} \in \mathbb{Z}^{1 \times t}$;
- left hand side matrix $A \in \mathbb{R}^{e \times t}$;
- right hand side matrix $B \in \mathbb{R}^{e \times v}$;
- a right hand side vector $D \in \mathbb{Z}^e$;
- an objective function vector $c \in \mathbb{Z}^{1 \times t}$;
- an incidence vector $y^s \in \{0, 1\}^v$ with $y_k^s = 1$ if $k \in s$ and $y_k^s = 0$ otherwise, $\forall k \in V$,

such that the characteristic function $c(s)$ equals the optimal objective value of the following integer linear program:

$$c(s, P) = \min_x \{cx + Pm(x) : Ax \geq By^s + D, \tilde{\alpha}x \leq m, x \in \mathbb{Z}^{t \times 1}\}$$

Note that the object function is not easy to analyze due to the joint effect of cx and Pm , we need to divide the function into two parts for a further discussion.

With this idea, we divide $c(s, m(s, P))$ as $c_0(s, m(s)) + Pm$. Via analysis and study, we find some properties about $c_0(s, m(s))$ which we call primary function later.

The primary function must be supermodular, then we can get the positive price or setup cost when the number of facilities changes. This corresponds to the following two lemmas.

Lemma 3.

$$c_0(V, m) - c_0(V, m-1) > 0 \Leftrightarrow P_m > 0, m = 2, \dots, n.$$

Lemma 4.

$$\begin{aligned} c_0(V, m) - c_0(V, m+1) &< c_0(V, m-1) - c_0(V, m) \\ \Leftrightarrow P_m &< P_{m+1}, m = 2, 3, \dots, n. \end{aligned}$$

Once the primary function must be supermodular, we can specify this property in the following two aspects. One is the concavity about the number of facilities, the other is the nature about the number of players ($s_1 \subset s_2$). They correspond to the following two formulas:

$$c_0(s_1, m-1) - c_0(s_1, m) \geq c_0(s_1, m) - c_0(s_1, m+1) \quad m = 2, \dots, n. \quad (2)$$

and

$$c_0(s_1, m-1) - c_0(s_1, m) \leq c_0(s_2, m-1) - c_0(s_2, m) \quad m = 2, \dots, n. \quad (3)$$

, respectively.

If the characteristic function satisfies the above property. Then P_1 can be calculated by the following procedure: When $\alpha(V) = c_0(V) + P_1$, which means the $\omega(P_1) = 0$. And the active inequalities must be the coalition contains $(n-1)$ players. Then plus all the $(n+1)$ equality together, we can obtain $P_1 = (n-1)c(V) - \sum_{|s|=n-1, s \in S} c(s)$. Because other inequalities will be satisfied according to the supermodular.

Then we can develop the Theorem 7.

Theorem 7. When two coalitions s_1, s_2 satisfy $s_1 \subset s_2$, the corresponding number of using facilities m_{s_1}, m_{s_2} have $m_{s_1} \leq m_{s_2}$, if the primary function satisfies (2) and (3).

Lemma 5. When $c_0(s)$ satisfy the supermodular and $P = P_1$, where $m_V = 1$, if $\alpha(s) = c(s) + P_1, |s| = n-1$ and $\alpha(V) = c_0(V) + P_1$ the budget balance constraint holds, the other coalitional stability constraints $\alpha(s) \leq c(s) + P_1, |s| < n-1$ will hold.

Proof 1 (Lemma 1). In fact, according to the SPT rule, a fixed number of using machines for the given grand coalition corresponds to the only deterministic sort order working on the machines, i.e. the deterministic cost. Therefore, by comparing the cost of the number of adjacent machines $(k, k+1)$ used by the grand coalition, the corresponding setup cost or the price P_i can be obtained. \square

Proof 2 (Theorem 1). By deriving from $\omega^*(P)$ (1), it can be seen that the SPF $\omega(P)$ is in fact the point-wise maximum of a set of straight lines, $c_0(V) + m(V)P + \sum_{s \in S \setminus \{V\}} -\rho_s [c_0(s) + m(s)P]$, each with a slope of $m(V) + \sum_{s \in S \setminus \{V\}} \rho_s m(s)$, for ρ in $\{\rho : \sum_{s \in S \setminus \{V\}: k \in s} \rho_s = 1 \text{ for all } k \in V, \text{ and } \rho_s \geq 0 \text{ for all } s \in S \setminus \{V\}\}$. Meanwhile, it's easy to see that $m(s)$ does not increase as P increases. Thus, we can claim that the SPF $\omega(P)$ is a convex function in P when $m(V)$ is a fixed integer.

Notice that the existence of $m(V)$ does not affect the property of the numbers of the breakpoints. To show that the SPF $\omega^*(P)$ is a piecewise linear function with a finite number of breakpoints, consider the feasible region of LP (1) for $\omega^*(P)$ without the part of $m(V)$ that is written as $\hat{\omega}(P)$, which is a convex polyhedron, denoted by \hat{R} . It can be seen that \hat{R} has a finite number of extreme points, and is independent of P . For any

given $P \in [0, P^*]$, by LP (1) we know that there must exist an extreme point ρ of \hat{R} such that $\dot{\omega}(P)$ equals $c_0(V) + m(V)P + \sum_{s \in S \setminus \{V\}} -\rho_s [c_0(s) + m(s)P]$ and that the derivative of $\dot{\omega}(P)$ at P equals $\sum_{s \in S \setminus \{V\}} \rho_s m(s)$. Thus, the derivative of $\dot{\omega}(P)$ for $P \in [0, P^*]$ can have only a finite number of possible values. Moreover, due to the convexity of $\dot{\omega}(P)$, the derivative of $\dot{\omega}(P)$ is non-decreasing in P . Hence, the derivative of $\dot{\omega}(P)$ can change for only a finite number of times when P increases from 0 to P^* . Therefore, $\omega^*(P)$ is a piecewise linear function with a finite number of breakpoints. \square

Proof 3 (Theorem 2). For convenience of expression, we set the setup cost as S_1, S_2, \dots, S_n at interval points while the number of machine changes. And S_i denotes the setup cost when the machine number changes from i to $i - 1$, especially, S_1 denotes the least setup cost when machine number is 1 and the corresponding subsidy is 0. We just need to prove the following equality

$$S_1 = S_2 + \dots + S_n = \sum_{i=2}^n S_i.$$

Notice that

$$(n-1) \sum_{s \in S \setminus \{V\}} \rho_s \geq \sum_{k \in V} \sum_{s \in S \setminus \{V\} : k \in s} \rho_s = n.$$

The left side of the inequality means for every ρ_s can appear at most $(n-1)$ times, so we should know that if and only if for every $\rho_s > 0$ appears $n-1$ times the quality holds. That is to say, the coalitions which contains $n-1$ players are all maximally unsatisfied coalitions. Then we have $\binom{n}{n-1}$ equalities.

$$\begin{cases} \alpha_1 + \alpha_2 + \dots + \alpha_{n-1} &= x_1 \\ \alpha_1 + \alpha_3 + \dots + \alpha_n &= x_2 \\ \vdots &\vdots \\ \alpha_2 + \alpha_3 + \dots + \alpha_n &= x_n. \end{cases}$$

Add these n equations together, and we can get

$$(n-1)(\alpha_1 + \alpha_2 + \dots + \alpha_n) = \sum_{i=1}^n x_i$$

As we know, x_1, x_2, \dots, x_n can be expressed as follows:

$$\begin{cases} x_1 = S_0 + (n-1)t_1 + (n-2)t_2 + \dots + t_{n-1} \\ x_2 = S_0 + (n-1)t_1 + (n-2)t_3 + \dots + t_{n-1} \\ \vdots \\ x_n = S_0 + (n-1)t_2 + (n-2)t_3 + \dots + t_n \end{cases}$$

According to SPT rule, we can obtain the equality $c(V) = \alpha_1 + \alpha_2 + \dots + \alpha_n = S_0 + nt_1 + (n-1)t_2 + \dots + t_n$.

By replacing x_1, x_2, \dots, x_n together with the expression of $c(V)$, we can get a equality only with $S_0, x_1, x_2, \dots, x_n$.

Finally, we can obtain $S_0 = \sum_{k=1}^n (n-k)t_k$. \square

Proof 4 (Theorem 3). For the IPU game, when using m machines is optimal, the setup cost S_0 must satisfies $t_m < S_0 < t_{m+1}$, which can be obtained from the process of calculating S_k , $\frac{n}{2} < k \leq n$, where $S_k = t_{n-k+1}$.

When $m > \frac{n}{2}$, the optimal sequence which means the minimum cost is each of the $n-m$ machines has to process two jobs, while the rest $2m-n$ machines each with one job. At present, there exists the following allocation which satisfies $\alpha(s) \leq c(s), s \in S$:

$$\begin{aligned}\alpha(1) &= 2t_1, \alpha(2) = 2t_2, \dots, \alpha(n-m) = 2t_{n-m}, \dots, \\ \alpha(n-m+1) &= S_0 + t_{n-m+1}, \dots, \alpha(m) = S_0 + t_m, \dots, \\ \alpha(m+1) &= S_0 + t_{m+1}, \dots, \alpha(n) = S_0 + t_n. \\ \text{where } t_m &< S_0 < t_{m+1}.\end{aligned}$$

Right now, $\alpha(1), \dots, \alpha(n)$ can not be bigger any more. That is, $\max \alpha(V) = c(V)$, then subsidy equals $c(V) - \alpha(V) = 0$. \square

Proof 5 (Theorem 4). From (1), we know that the slope is $m(V) - \sum_{s \in S \setminus \{V\}} \rho_s \cdot m(s)$ when setup cost is S_i . At the left side of S_i , the number of using machines is $m(V) = i$ and the slope equals $i - \sum_{s \in S \setminus \{V\}} \rho_s \cdot m(s)$, which is positive, while at the right side of S_i , the number of using machines is $m(V) = i-1$ and the slope equals $\sum_{s \in S \setminus \{V\}} \rho_s \cdot m(s) - (i-1)$, which is negative. So the sum of absolute values of slope on the left and right sides at S_i is 1. When $m(V)$ is 1, the expression of slope is $1 - \sum_{s \in S \setminus \{V\}} \rho_s \cdot m(s)$. Meanwhile, $m(s) \leq m(V)$, so $m(s) = 1$, then the slope is $1 - \sum_{s \in S \setminus \{V\}} \rho_s$. Because we know that the sum of absolute values of slope at both sides of S_2 is 1, so the absolute value of the right side of slope less than 1.

From Theorem 1, we have:

$$(n-1) \sum_{s \in S \setminus \{V\}} \rho_s \geq \sum_{k \in V} \sum_{s \in S \setminus \{V\}: k \in s} \rho_s = n$$

$\sum_{s \in S \setminus \{V\}} \rho_s$ minimum is $\frac{n}{n-1}$ the corresponding maximum slope is $1 - \sum_{s \in S \setminus \{V\}} \rho_s = -\frac{1}{n-1}$. \square

Proof 6 (Theorem 5). When the value of P is between (P_2, P_1) , where the grand coalition only uses one machine, all the other sub-coalitions also use one machine. Now assume that the theorem is not tenable. When decreasing the value of P , for example, from P_2 to P_3 . At some time, there must be such a situation that a sub-coalition denoted as s' ($|s'| \geq 2$) uses two machines which is optimal. Then we have $\alpha(s') = c(s') + 2P$, there exist two coalitions s_1, s_2 which satisfy $s_1 \cup s_2 = s', s_1 \cap s_2 = \emptyset$. Meanwhile, s_1, s_2 also satisfy the two constraints $\alpha(s_1) \leq c(s_1) + P, \alpha(s_2) \leq c(s_2) + P$. Then we can obtain that $c(s_1) + c(s_2) \geq c(s')$. At the same time, we know that the optimal sequence result shows $c(s_1) + c(s_2) = c(s')$, which means the bigger coalition can

be substituted by two smaller ones. Consequently, the coalitions s only use one machine if the corresponding constraints are valid. \square

Proof 7 (Theorem 6). The complexity of the whole problem can be divided into several stages, the first one is how to obtain the value of P_i . For the IVPU game, the SPT rule can be used to calculate the $c(s)$. By solving the last n equalities, we can obtain P_1 . Then we need to know the number of breakpoints, by IPC algorithm, we can construct the whole graph. Then the left problem is how to solve the slope in IPC algorithm, and how is the essential coalition set generated. The CP algorithm is used to obtain the subsidy and conclude a separation problem, if the separation problem can be solved easily, then the subsidy can be obtained easily. Then we use DP algorithm to solve the separation problem and with the separation problem we can obtain the essential set. If the set is polynomial, we can calculate the slope by solving the LP. \square

Proof 8 (Lemma 2&3). As we all know, when $P = 0$, it's optimal that every player in the coalition s use one machine each. For the coalition s , when the price equals P_m , where the costs produced by the grand coalition are equal in value, the following formulas hold $c(V, m) = c_0(s, m) + Pm$, $c(V, m - 1) = c_0(s, m - 1) + P(m - 1)$ and $c(V, m) = c(V, m - 1)$. Then we can obtain $P_m = c_0(V, m) - c_0(V, m - 1)$, $m = 2, \dots, n$. Therefore, if $c_0(V, m) - c_0(V, m - 1) > 0$, we can get $P_m > 0$.

Similarly, by substituting m with $m + 1$, we can obtain that $P_{m+1} = c_0(V, m + 1) - c_0(V, m)$, $m = 2, 3, \dots, n$. If $c_0(V, m) - c_0(V, m + 1) < c_0(V, m - 1) - c_0(V, m)$, then $P_m < P_{m+1}$, $m = 2, 3, \dots, n$. \square

Proof 9 (Theorem 7). As we described before, the characteristic function $c(s, m(s, P))$ can be expressed as $c_0(s, m(s)) + Pm$. For the two coalitions s_1, s_2 , we use $c_0(s_2, m(s_1))$ to represent the cost produced by coalition s_2 using $m(s_1)$ machines, where $m(s_1)$ is the number of machines used by coalition s_1 when the cost is minimum.

According to this definition and Lemma 2, obviously the following relation holds, $c(s_1, m(s_1)) < c(s_1, m(s_1) - 1) \Rightarrow c_0(s_1, m(s_1) - 1) - c_0(s_1, m(s_1)) > 0$, because $c_0(s_1, m(s_1))$ is the minimum cost for s_1 using $m(s_1)$ machines.

For the two coalitions s_1 , when the coalition s_1 uses $m(s_1)$ machines which is optimal, the following relation is established: $m(s_1)P + c_0(s_1, m) < (m(s_1) - 1)P + c_0(s_1, m - 1) \Rightarrow P < c_0(s_1, m - 1) - c_0(s_1, m)$.

When coalition s_1 uses $m(s_1)$, we assume that coalition s_2 uses $m(s_2) = m(s_1) - 1 < m(s_1)$ machines, which means $c(s_2, m(s_1) - 1) < c(s_2, m(s_1))$. Then expand the formula, we obtain $P > c_0(s_2, m(s_1) - 1) - c_0(s_2, m(s_1))$. However, because using $m(s_1)$ by coalition s_1 assures that $c(s_1, m(s_1)) < c(s_1, m(s_1) - 1) \Rightarrow P < c_0(s_1, m(s_1) - 1) - c_0(s_1, m(s_1))$. So we can obtain that $c_0(s_2, m(s_1) - 1) - c_0(s_2, m(s_1)) < P < c_0(s_1, m(s_1) - 1) - c_0(s_1, m(s_1))$, which contradicts the condition (2). With the result of that, we know $m(s_2) \neq m(s_1) - 1$.

Meanwhile, according to the condition (3) which tells us about the concavity of the number of machines. The larger the number of machines, the smaller the difference between these two items $c_0(s_1, m(s_1) - 1)$ and $c_0(s_1, m(s_1))$. Take $m(s_2) = m(s_1) - 2$ as an example, similar to the above, we can get $c(s_2, m(s_1) - 2) < c(s_2, m(s_1) - 1) \Rightarrow P > c_0(s_2, m(s_1) - 2) - c_0(s_2, m(s_1) - 1)$. According to the conditions (2), $c(s_1, m(s_1)) < c(s_1, m(s_1) - 1) \Rightarrow P < c_0(s_1, m(s_1) - 1) - c_0(s_1, m(s_1)) < c_0(s_1, m(s_1) - 2) - c_0(s_1, m(s_1) - 1) < c_0(s_2, m(s_1) -$

$2) - c_0(s_2, m(s_1) - 1)$ contradicts $P > c_0(s_2, m(s_1) - 2) - c_0(s_2, m(s_1) - 1)$ we just obtained above. Consequently, we know that $m(s_2)$ cannot be less than $m(s_1) - 1$. In conclusion, $m(s_1) \leq m(s_2)$. \square

Proof 10 (Lemma 4). With the n equations for $|s| = n - 1$ and $\alpha(V) = c_0(V) + P_1$, we can obtain the values of $\alpha(i)$. For each $|s'| < n - 1$, suppose that $s' = \{i, \dots, j\}$, we just need to prove:

$$\alpha(i, \dots, j) \leq c_0(i, \dots, j) + P_1 \quad (4)$$

Substitute $P_1 = \alpha(V) - c(V)$ and $\alpha(i) = c(V) - c(-i)$ into the equation (4), denote $c(-i) = c(\{1, \dots, i - 1, i + 1, \dots, n\})$, we just need to prove the following relation:

$$\sum_i^{N \setminus s'} c(-i) \leq (n - 1 - |s'|)c(V) + c(s') \quad (5)$$

\Rightarrow

$$c(-k) - c(s') \leq (n - 1 - |s'|)c(V) - \sum_i^{N \setminus s' \setminus k} c(-i)$$

Because $c(s)$ satisfies the supermodular, we have the following set of inequalities.

$$\begin{cases} c(V) - c(-t) \geq & c(-k) - c(-k - t) \\ c(V) - c(-m) \geq & c(-k - t) - c(-k - t - m) \\ \vdots & \vdots \\ c(V) - c(-s) \geq & c(s' \cup s) - c(s') \end{cases}$$

Add these $(n - 1 - |s'|)$ inequalities together, we can obtain $(n - 1 - |s'|)c(V) - \sum_i^{N \setminus s'} c(-i) + c(s') \geq 0$.

That is, $\alpha(s') \leq c(s') + P_1, \forall s', |s'| < n - 1$. \square

The IPU game is very special so that we can use SPT rule to compute the $c_{IPU}(s)$, which means jobs in coalition s with the shortest processing time should be processed first and the longest ones last. Then we can use dynamic programming to solve the separation problem $\delta_{IPU} = \min \{c_{IPU}(s) + P - \bar{\alpha}(s, P) : \forall s \in \mathbb{S} \setminus \{N\}\}$