

Homework 1

Li Zikang

2-5

(a) Write down a recursion for $f^{(n)}(x)$ in terms of $f^{(n-1)}(\cdot)$ and the parameters of the problem.

For any participant, with n more spins, he/she can choose to spin or stop. When choosing **stopping**, he/she will take away the current accumulation of x . We can obtain $f^{(n)}(x) = x$. While choosing **spinning**, it is also easy to know that

$$f^{(n)}(x) = \sum_i f^{(n-1)}(x + r_i) \cdot p_i, \text{ where } i = 1, \dots, m.$$

$$f^{(n)}(x) = \max\{\sum_i f^{(n-1)}(x + r_i) \cdot p_i, x\}$$

(b) Solve the problem numerically for the case in which $m = 3, p_i = 1/4$, for $i = 1, 2, 3, r_1 = 1, r_2 = 5, r_3 = 10$, and there are one or two more spins available.

At the beginning, the winning is 0, so must choose to spin.

$$f^{(2)}(0) = \begin{cases} f^{(1)}(1), \text{ probability } 1/4 \\ f^{(1)}(5), \text{ probability } 1/4 \\ f^{(1)}(10), \text{ probability } 1/4 \\ 0 \quad \text{probability } 1/4 \end{cases}$$

$$f^{(1)}(x) = \begin{cases} x + 1, \text{probability } 1/4 \\ x + 5, \text{probability } 1/4 \\ x + 10, \text{probability } 1/4 \\ 0 \quad \text{probability } 1/4 \\ x \quad \text{choose to stop} \end{cases}$$

So when spinning once, the expected winnings are $1/4 \times (1 + 5 + 10) = 4$. And when spinning twice, $f^{(1)}(x) = 1/4 \times [(x + 1) + (x + 5) + (x + 10)]$. $f^{(2)}(0) = 1/4 \times (f^{(1)}(1) + f^{(1)}(5) + f^{(1)}(10)) = 1/4 \times (4 + 3/4 + 4 + 15/4 + 4 + 30/4) = 6$

In particular, $f^{(0)}(x) = x, f^{(1)}(x) = \max\{3/4x + 4, x\}$. Let $3/4x + 4 = x, x = 16$. When $x > 16$, the participant will take away the winnings; when $x < 16$, the participant will spin again. Consequently, the expected total winnings are 16 for this case.

2-6

(a) Write down a recursion for $v(x)$ and use it to find an explicit expression for $v(x)$ in terms of $x, \Phi(x), I(x)$, and α , where $I(x) := E(X - x)^+ = \int_x^\infty (\xi - x)\phi(\xi)d\xi$.

$$v(x, n) = v(x, n - 1) \cdot \alpha \cdot \Phi(x) + \int_x^\infty \xi \phi(\xi) d\xi$$

$$\text{Let } V = \int_x^\infty \xi \phi(\xi) d\xi.$$

$$v(x, n) = v(x, 1) \cdot [\alpha \cdot \Phi(x)]^{(n-1)} + V[1 + [\alpha \cdot \Phi(x)] + [\alpha \cdot \Phi(x)]^2 + \dots + [\alpha \cdot \Phi(x)]^{n-2}]$$

$$v(x, 1) = \int_x^\infty \xi \phi(\xi) d\xi = V$$

Then we can obtain that $v(x) = V \frac{1-(\alpha \cdot \Phi(x))^n}{1-\alpha \cdot \Phi(x)}$, $\alpha \in (0, 1)$. Because this is an infinite period, let $n \rightarrow \infty$, so $v(x) = \frac{V}{1-\alpha \cdot \Phi(x)}$.

$$I(x) = \int_x^\infty (\xi - x) \phi(\xi) d\xi = \int_x^\infty \xi \phi(\xi) d\xi - x(1 - \Phi(x))$$

So $V = I(x) + x(1 - \Phi(x))$, substitute it in the $v(x)$, we get $v(x) = \frac{I(x) + x(1 - \Phi(x))}{1 - \alpha \cdot \Phi(x)}$

(b) $\alpha = 0.99$ bids $\sim N(1000, 200)$. $x = 1200$. So $v(1200) = \frac{I(1200) + 1200(1 - \Phi(1200))}{1 - \alpha \Phi(1200)}$, $I(1200) = 200I_N(1)$, $\Phi(1200) = \Phi_N(1) = 0.841$, $I_N(1) = \phi_N(1) - 1 + \Phi_N(1) = 0.083$.

By calculating, $v(1200) = 1238.9$

2-7

(a) Display Ω .

$$\Omega = \{(3, 3, 1), (2, 3, 2), (2, 2, 2), (1, 3, 2), (2, 3, 1), (0, 3, 2), (1, 3, 1), (1, 1, 2), (2, 2, 1), (2, 0, 2), (3, 0, 1), (1, 0, 2), (2, 0, 1), (1, 1, 1), (0, 0, 2)\}$$

(b) $S_0 = \{(3, 3, 1)\}$, $S_1 = \{(2, 3, 2), (2, 2, 2), (1, 3, 2)\}$, $S_2 = \{(2, 3, 1)\}$, $S_3 = \{(0, 3, 2)\}$, $S_4 = \{(1, 3, 1)\}$, $S_5 = \{(1, 1, 2)\}$, $S_6 = \{(2, 2, 1)\}$, $S_7 = \{(2, 0, 2)\}$, $S_8 = \{(3, 0, 1)\}$, $S_9 = \{(1, 0, 2)\}$, $S_{10} = \{(2, 0, 1), (1, 1, 1)\}$, $S_{11} = \{(0, 0, 2)\}$.

$$m = 11.$$

3-4

(a) Write down a recursion for $f^n(x|S)$.

$$f^{(n)}(x|S) = \sum_i f^{(n-1)}(x + r_i|S) \cdot p_i.$$

$$f^{(n)}(x|S) = \begin{cases} \sum_i f^{(n-1)}(x + r_i|S) \cdot p_i, & \text{spin or } x < S. \\ x & , \text{stop.} \end{cases}$$

(b) Being indifferent means $\sum_i f^{(n-1)}(S + r_i|S) \cdot p_i = S = f^{(n)}(S|S) \Rightarrow \sum_i^m (S^* + r_i)p_i = S^* \Rightarrow S^* = \frac{\mu}{p}$.

3-5

(a) Write a recursion for f_n in terms of f_{n-1}

$$f_n(x) = f_{n-1}(x) \cdot \alpha + I(\alpha x)$$

(b) Calculate f_n . $f_2 =$

3-6

test1.

4-4

test1.

4-5

导数小于1, 所以f(x) 线性函数且系数为1.