

# 1 Binary Logistic Regression

Binary Logistic regression is used to determine the impact of multiple independent variables presented simultaneously to predict membership of one or other of the two dependent variable categories.

## 1.1 The Hosmer-Lemeshow Test

The Hosmer-Lemeshow test of goodness of fit is not automatically a part of the SPSS logistic regression output. To get this output, we need to go into 'options and tick the box marked Hosmer-Lemeshow test of goodness of fit. In our example, this gives us the following output:

Step	Chi-square	df	Sig.
1	142.032	6	.000

Therefore, our model is significant, suggesting it does not fit the data. However, as we have a sample size of over 13,000, even very small divergencies of the model from the data would be flagged up and cause significance. Therefore, with samples of this size it is hard to find models that are parsimonious (i.e. that use the minimum amount of independent variables to explain the dependent variable) and fit the data. Therefore, other fit indices might be more appropriate.

## 2 Logistic Regression

Logistic regression, also called a logit model, is used to model **dichotomous outcome** variables. In the logit model the **log odds** of the outcome is modeled as a linear combination of the predictor variables.

In logistic regression theory, the predicted dependent variable is a function of the probability that a particular subject will be in one of the categories (for example, the probability that a patient has the disease, given his or her set of scores on the predictor variables).

## 3 Model Diagnostics for Logistic Regression

### 4 The Wald Test

The Wald test is a way of testing the significance of particular explanatory variables in a statistical model.

In logistic regression we have a binary outcome variable and one or more explanatory variables. For each explanatory variable in the model there will be an associated parameter. The Wald test is one of a number of ways of testing whether the parameters associated with a group of explanatory variables are zero.

If for a particular explanatory variable, or group of explanatory variables, the Wald test is significant, then we would conclude that the parameters associated with these variables are not zero, so that the variables should be included in the model. If the Wald test is not significant then these explanatory variables can be omitted from the model.

When considering a single explanatory variable, Altman (1991) uses a t-test to check whether the parameter is significant. For a single parameter the Wald statistic is just the square of the

t-statistic and so will give exactly equivalent results.

An alternative and widely used approach to testing the significance of a number of explanatory variables is to use the likelihood ratio test. This is appropriate for a variety of types of statistical models.

Agresti (1990) argues that the likelihood ratio test is better, particularly if the sample size is small or the parameters are large.

## 4.1 Logistic Regression

$$\pi(x) = \frac{e^{(\beta_0 + \beta_1 x)}}{e^{(\beta_0 + \beta_1 x)} + 1} = \frac{1}{e^{-(\beta_0 + \beta_1 x)} + 1},$$

and

$$g(x) = \ln \frac{\pi(x)}{1 - \pi(x)} = \beta_0 + \beta_1 x,$$

and

$$\frac{\pi(x)}{1 - \pi(x)} = e^{(\beta_0 + \beta_1 x)}.$$

## 4.2 Wald statistic

Alternatively, when assessing the contribution of individual predictors in a given model, one may examine the significance of the Wald statistic. The Wald statistic, analogous to the t-test in linear regression, is used to assess the significance of coefficients. The Wald statistic is the ratio of the square of the regression coefficient to the square of the standard error of the coefficient and is asymptotically distributed as a chi-square distribution.

## 4.3 Wald statistic

The Wald statistic is commonly used to test the significance of individual logistic regression coefficients for each independent variable (that is, to test the null hypothesis in logistic regression that a particular logit (effect) coefficient is zero).

The Wald Statistic is the ratio of the unstandardized logit coefficient to its standard error. The Wald statistic and its corresponding p probability level is part of SPSS output in the section ***Variables in the Equation***. This corresponds to significance testing of b coefficients in OLS regression. The researcher may well want to drop independents from the model when their effect is not significant by the Wald statistic.

## 4.4 Logistic Regression: Decision Rule

Our decision rule will take the following form: If the probability of the event is greater than or equal to some threshold, we shall predict that the event will take place. By default, SPSS sets this threshold to .5. While that seems reasonable, in many cases we may want to set it higher or lower than .5.

## 5 Model Diagnostics for Logistic Regression

### 5.1 Omnibus Test for Model Coefficients

The overall significance is tested using what SPSS calls the *Model Chi-square*, which is derived from the likelihood of observing the actual data under the assumption that the model that has been fitted is accurate. There are two hypotheses to test in relation to the overall fit of the model:

$H_0$  The model is a good fitting model.

$H_1$  The model is not a good fitting model (i.e. the predictors have a significant effect).

- For the example provided here, the model chi square has 2 degrees of freedom, a value of 24.096 and a probability of  $p < 0.000$ . (*see Next Page for SPSS Output*).
- Thus, the indication is that the model has a poor fit, with the model containing only the constant indicating that the predictors do have a significant effect and create essentially a different model.
- So we need to look closely at the predictors and from later tables determine if one or both are significant predictors.

### Remark upon “Steps”

- This table has 1 step. This is because we are entering both variables and at the same time providing only one model to compare with the constant model.
- In stepwise logistic regression there are a number of steps listed in the table as each variable is added or removed, creating different models.
- The step is a measure of the improvement in the predictive power of the model since the previous step.
- I will revert to this next class.

## 5.2 Omnibus Test for Model Coefficients

The overall significance is tested using what SPSS calls the ***Model Chi-square***, which is derived from the likelihood of observing the actual data under the assumption that the model that has been fitted is accurate. There are two hypotheses to test in relation to the overall fit of the model:

$H_0$  The model is a good fitting model.

$H_1$  The model is not a good fitting model (i.e. the predictors have a significant effect).

- For the example provided here, the model chi square has 2 degrees of freedom, a value of 24.096 and a probability of  $p < 0.000$ . (see Next Page for SPSS Output).
- Thus, the indication is that the model has a poor fit, with the model containing only the constant indicating that the predictors do have a significant effect and create essentially a different model.
- So we need to look closely at the predictors and from later tables determine if one or both are significant predictors.

### Remark upon “Steps”

- This table has 1 step. This is because we are entering both variables and at the same time providing only one model to compare with the constant model.
- In stepwise logistic regression there are a number of steps listed in the table as each variable is added or removed, creating different models.
- The step is a measure of the improvement in the predictive power of the model since the previous step.
- I will revert to this next class.

## 5.3 Omnibus Test for Model Coefficients

The overall significance is tested using what SPSS calls the ***Model Chi-square***, which is derived from the likelihood of observing the actual data under the assumption that the model that has been fitted is accurate. There are two hypotheses to test in relation to the overall fit of the model:

$H_0$  The model is a good fitting model.

$H_1$  The model is not a good fitting model (i.e. the predictors have a significant effect).

- For the example provided here, the model chi square has 2 degrees of freedom, a value of 24.096 and a probability of  $p < 0.000$ . (*see Next Page for SPSS Output*).
- Thus, the indication is that the model has a poor fit, with the model containing only the constant indicating that the predictors do have a significant effect and create essentially a different model.
- So we need to look closely at the predictors and from later tables determine if one or both are significant predictors.

### Remark upon “Steps”

- This table has 1 step. This is because we are entering both variables and at the same time providing only one model to compare with the constant model.
- In stepwise logistic regression there are a number of steps listed in the table as each variable is added or removed, creating different models.
- The step is a measure of the improvement in the predictive power of the model since the previous step.
- I will revert to this next class.

## 5.4 Omnibus Test for Model Coefficients

The overall significance is tested using what SPSS calls the **Model Chi-square**, which is derived from the likelihood of observing the actual data under the assumption that the model that has been fitted is accurate. There are two hypotheses to test in relation to the overall fit of the model:

$H_0$  The model is a good fitting model.

$H_1$  The model is not a good fitting model (i.e. the predictors have a significant effect).

- For the example provided here, the model chi square has 2 degrees of freedom, a value of 24.096 and a probability of  $p < 0.000$ . (*see Next Page for SPSS Output*).
- Thus, the indication is that the model has a poor fit, with the model containing only the constant indicating that the predictors do have a significant effect and create essentially a different model.
- So we need to look closely at the predictors and from later tables determine if one or both are significant predictors.

### Remark upon “Steps”

- This table has 1 step. This is because we are entering both variables and at the same time providing only one model to compare with the constant model.
- In stepwise logistic regression there are a number of steps listed in the table as each variable is added or removed, creating different models.
- The step is a measure of the improvement in the predictive power of the model since the previous step.
- I will revert to this next class.

## 5.5 Model Summary Table

- The likelihood function can be thought of as a measure of how well a candidate model fits the data (although that is a very simplistic definition).
- Later on in the course, we will meet the AIC Function. The AIC criterion is based on the Likelihood function.
- The likelihood function of a fitted model is commonly re-expressed as  $-2LL$  (i.e. The log of the likelihood times minus 2). The 2LL value from the Model Summary table below is 17.359.
- It is not immediately clear how to interpret  $-2LL$  yet. However this measure is useful in comparing various candidate models, and is therefore used in **Variable Selection Procedure**, such as backward and forward selection.

### Block 1: Method = Enter

**Omnibus Tests of Model Coefficients**

		Chi-square	df	Sig.
Step 1	Step	24.096	2	.000
	Block	24.096	2	.000
	Model	24.096	2	.000

**Model Summary**

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	17.359 <sup>a</sup>	.552	.737

a. Estimation terminated at iteration number 8 because parameter estimates changed by less than .001.

**Hosmer and Lemeshow Test**

Step	Chi-square	df	Sig.
1	6.378	8	.605

## 5.6 Model Summary Table

- The likelihood function can be thought of as a measure of how well a candidate model fits the data (although that is a very simplistic definition).
- Later on in the course, we will meet the AIC Function. The AIC criterion is based on the Likelihood function.
- The likelihood function of a fitted model is commonly re-expressed as  $-2LL$  (i.e. The log of the likelihood times minus 2). The 2LL value from the Model Summary table below is 17.359.
- It is not immediately clear how to interpret  $-2LL$  yet. However this measure is useful in comparing various candidate models, and is therefore used in **Variable Selection Procedure**, such as backward and forward selection.

### Block 1: Method = Enter

**Omnibus Tests of Model Coefficients**

		Chi-square	df	Sig.
Step 1	Step	24.096	2	.000
	Block	24.096	2	.000
	Model	24.096	2	.000

**Model Summary**

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	17.359 <sup>a</sup>	.552	.737

a. Estimation terminated at iteration number 8 because parameter estimates changed by less than .001.

**Hosmer and Lemeshow Test**

Step	Chi-square	df	Sig.
1	6.378	8	.605



## 5.7 Model Summary Table

- The likelihood function can be thought of as a measure of how well a candidate model fits the data (although that is a very simplistic definition).
- Later on in the course, we will meet the AIC Function. The AIC criterion is based on the Likelihood function.
- The likelihood function of a fitted model is commonly re-expressed as  $-2LL$  (i.e. The log of the likelihood times minus 2). The 2LL value from the Model Summary table below is 17.359.
- It is not immediately clear how to interpret  $-2LL$  yet. However this measure is useful in comparing various candidate models, and is therefore used in **Variable Selection Procedure**, such as backward and forward selection.

### Block 1: Method = Enter

**Omnibus Tests of Model Coefficients**

		Chi-square	df	Sig.
Step 1	Step	24.096	2	.000
	Block	24.096	2	.000
	Model	24.096	2	.000

**Model Summary**

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	17.359 <sup>a</sup>	.552	.737

a. Estimation terminated at iteration number 8 because parameter estimates changed by less than .001.

**Hosmer and Lemeshow Test**

Step	Chi-square	df	Sig.
1	6.378	8	.605

## 5.8 Hosmer-Lemeshow Goodness-of-Fit

The Hosmer-Lemeshow Goodness-of-Fit test tells us whether we have constructed a valid overall model or not. If the model is a good fit to the data then the Hosmer-Lemeshow Goodness-of-Fit test should have an associated p-value greater than 0.05.

## 5.9 Hosmer-Lemeshow Goodness-of-Fit Test

- The Hosmer-Lemeshow Goodness-of-Fit test tells us whether we have constructed a valid overall model or not, and is an alternative to model chi square procedure.
- If the model is a good fit to the data then the Hosmer-Lemeshow Goodness-of-Fit test should have an associated p-value greater than 0.05.

If the H-L goodness-of-fit test statistic is greater than .05, as we want for well-fitting models, we fail to reject the null hypothesis that there is no difference between observed and model-predicted values, implying that the models estimates fit the data at an acceptable level. That is, well-fitting models show non-significance on the H-L goodness-of-fit test. This desirable outcome of non-significance indicates that the model prediction does not significantly differ from the observed.

### Performing the Hosmer-Lemeshow Goodness-of-Fit Test

- The Hosmer-Lemeshow test of goodness of fit is not automatically a part of the SPSS logistic regression output. To get this output, we need to go into **options** and tick the box marked Hosmer-Lemeshow test of goodness of fit.

In our example, this gives us the following output:

Step	Chi-square	df	Sig.
1	142.032	6	.000

Therefore, our model is significant, suggesting it does not fit the data. However, as we have a sample size of over 13,000, even very small divergencies of the model from the data would be flagged up and cause significance. Therefore, with samples of this size it is hard to find models that are parsimonious (i.e. that use the minimum amount of independent variables to explain the dependent variable) and fit the data. Therefore, other fit indices might be more appropriate.

### Computing the Hosmer and Lemeshow Test Statistic

- The Hosmer and Lemeshow test divides subjects into 10 ordered groups of subjects and then compares the number actually in the each group (observed) to the number predicted by the logistic regression model (predicted).
- The 10 ordered groups are created based on their estimated probability; those with estimated probability below .1 form one group, and so on, up to those with probability .9 to 1.0.

- Each of these categories is further divided into two groups based on the actual observed outcome variable (success, failure). The expected frequencies for each of the cells are obtained from the model. A probability (p) value is computed from the chi-square distribution with 8 degrees of freedom to test the fit of the logistic model.

		take solar panel offer = decline offer		take solar panel offer = take offer		Total
		Observed	Expected	Observed	Expected	
Step 1	1	3	2.996	0	.004	3
	2	3	2.957	0	.043	3
	3	3	2.629	0	.371	3
	4	2	2.136	1	.864	3
	5	2	1.881	1	1.119	3
	6	0	.833	3	2.167	3
	7	0	.376	3	2.624	3
	8	1	.171	2	2.829	3
	9	0	.019	3	2.981	3
	10	0	.000	3	3.000	3

Figure 1: Hosmer and Lemeshow Table

### Sample Adequacy

The H-L statistic assumes sampling adequacy, with a rule of thumb being enough cases so that 95% of cells (typically, 10 decile groups times 2 outcome categories = 20 cells) have an expected frequency  $> 5$ . Our H-L statistic has a significance of .605 which means that it is not statistically significant and therefore our model is quite a good fit.

Hosmer and Lemeshow Test			
Step	Chi-square	df	Sig.
1	6.378	8	.605

Figure 2: Hosmer and Lemeshow Statistic

## 5.10 Hosmer-Lemeshow Goodness-of-Fit Test

- The Hosmer-Lemeshow Goodness-of-Fit test tells us whether we have constructed a valid overall model or not, and is an alternative to model chi square procedure.
- If the model is a good fit to the data then the Hosmer-Lemeshow Goodness-of-Fit test should have an associated p-value greater than 0.05.

If the H-L goodness-of-fit test statistic is greater than .05, as we want for well-fitting models, we fail to reject the null hypothesis that there is no difference between observed and model-predicted values, implying that the models estimates fit the data at an acceptable level. That is, well-fitting models show non-significance on the H-L goodness-of-fit test. This desirable outcome of non-significance indicates that the model prediction does not significantly differ from the observed.

### Performing the Hosmer-Lemeshow Goodness-of-Fit Test

- The Hosmer-Lemeshow test of goodness of fit is not automatically a part of the SPSS logistic regression output. To get this output, we need to go into **options** and tick the box marked Hosmer-Lemeshow test of goodness of fit.

In our example, this gives us the following output:

Step	Chi-square	df	Sig.
1	142.032	6	.000

Therefore, our model is significant, suggesting it does not fit the data. However, as we have a sample size of over 13,000, even very small divergencies of the model from the data would be flagged up and cause significance. Therefore, with samples of this size it is hard to find models that are parsimonious (i.e. that use the minimum amount of independent variables to explain the dependent variable) and fit the data. Therefore, other fit indices might be more appropriate.

### Computing the Hosmer and Lemeshow Test Statistic

- The Hosmer and Lemeshow test divides subjects into 10 ordered groups of subjects and then compares the number actually in the each group (observed) to the number predicted by the logistic regression model (predicted).
- The 10 ordered groups are created based on their estimated probability; those with estimated probability below .1 form one group, and so on, up to those with probability .9 to 1.0.
- Each of these categories is further divided into two groups based on the actual observed outcome variable (success, failure). The expected frequencies for each of the cells are obtained from the model. A probability (p) value is computed from the chi-square distribution with 8 degrees of freedom to test the fit of the logistic model.

		take solar panel offer = decline offer		take solar panel offer = take offer		Total
		Observed	Expected	Observed	Expected	
Step 1	1	3	2.996	0	.004	3
	2	3	2.957	0	.043	3
	3	3	2.629	0	.371	3
	4	2	2.136	1	.864	3
	5	2	1.881	1	1.119	3
	6	0	.833	3	2.167	3
	7	0	.376	3	2.624	3
	8	1	.171	2	2.829	3
	9	0	.019	3	2.981	3
	10	0	.000	3	3.000	3

Figure 3: Hosmer and Lemeshow Table

### Sample Adequacy

The H-L statistic assumes sampling adequacy, with a rule of thumb being enough cases so that 95% of cells (typically, 10 decile groups times 2 outcome categories = 20 cells) have an expected frequency  $> 5$ . Our H-L statistic has a significance of .605 which means that it is not statistically significant and therefore our model is quite a good fit.

Hosmer and Lemeshow Test			
Step	Chi-square	df	Sig.
1	6.378	8	.605

Figure 4: Hosmer and Lemeshow Statistic

### 5.11 Hosmer-Lemeshow Goodness-of-Fit Test

- The Hosmer-Lemeshow Goodness-of-Fit test tells us whether we have constructed a valid overall model or not, and is an alternative to model chi square procedure.
- If the model is a good fit to the data then the Hosmer-Lemeshow Goodness-of-Fit test should have an associated p-value greater than 0.05.

If the H-L goodness-of-fit test statistic is greater than .05, as we want for well-fitting models, we fail to reject the null hypothesis that there is no difference between observed and model-predicted values, implying that the models estimates fit the data at an acceptable level. That

is, well-fitting models show non-significance on the H-L goodness-of-fit test. This desirable outcome of non-significance indicates that the model prediction does not significantly differ from the observed.

### Performing the Hosmer-Lemeshow Goodness-of-Fit Test

- The Hosmer-Lemeshow test of goodness of fit is not automatically a part of the SPSS logistic regression output. To get this output, we need to go into **options** and tick the box marked Hosmer-Lemeshow test of goodness of fit.

In our example, this gives us the following output:

Step	Chi-square	df	Sig.
1	142.032	6	.000

Therefore, our model is significant, suggesting it does not fit the data. However, as we have a sample size of over 13,000, even very small divergencies of the model from the data would be flagged up and cause significance. Therefore, with samples of this size it is hard to find models that are parsimonious (i.e. that use the minimum amount of independent variables to explain the dependent variable) and fit the data. Therefore, other fit indices might be more appropriate.

### Computing the Hosmer and Lemeshow Test Statistic

- The Hosmer and Lemeshow test divides subjects into 10 ordered groups of subjects and then compares the number actually in the each group (observed) to the number predicted by the logistic regression model (predicted).
- The 10 ordered groups are created based on their estimated probability; those with estimated probability below .1 form one group, and so on, up to those with probability .9 to 1.0.
- Each of these categories is further divided into two groups based on the actual observed outcome variable (success, failure). The expected frequencies for each of the cells are obtained from the model. A probability (p) value is computed from the chi-square distribution with 8 degrees of freedom to test the fit of the logistic model.

#### 5.11.1 Sample Adequacy

The H-L statistic assumes sampling adequacy, with a rule of thumb being enough cases so that 95% of cells (typically, 10 decile groups times 2 outcome categories = 20 cells) have an expected frequency  $> 5$ . Our H-L statistic has a significance of .605 which means that it is not statistically significant and therefore our model is quite a good fit.

Contingency Table for Hosmer and Lemeshow Test						
		take solar panel offer = decline offer		take solar panel offer = take offer		Total
		Observed	Expected	Observed	Expected	
Step 1	1	3	2.996	0	.004	3
	2	3	2.957	0	.043	3
	3	3	2.629	0	.371	3
	4	2	2.136	1	.864	3
	5	2	1.881	1	1.119	3
	6	0	.833	3	2.167	3
	7	0	.376	3	2.624	3
	8	1	.171	2	2.829	3
	9	0	.019	3	2.981	3
	10	0	.000	3	3.000	3

Figure 5: Hosmer and Lemeshow Table

Hosmer and Lemeshow Test			
Step	Chi-square	df	Sig.
1	6.378	8	.605

Figure 6: Hosmer and Lemeshow Statistic