

Supplementary Results Table 3. All-cause deaths, death FAPs, DALYs, DALY FAPs, and percentage change for each risk factor and outcome for both sexes combined in 2000 and 2016 for China.

	2000 Deaths (in thousands)	2000 Deaths FAP	2016 Deaths (in thousands)	2016 Deaths FAP	Percentage change Deaths (2000-2016)	Percentage change Deaths FAP (2000-2016)	Percentage change DALYs (2000-2016)	2000 DALYs (in thousands)	2000 DALYs FAP	2016 DALYs (in thousands)	2016 DALYs FAP	Percentage change DALYs (2000-2016)	Percentage change DALYs FAP (2000-2016)	Percentage change DALYs per 100,000 pop (2000-2016)
Stroke, bronch, and lung cancer	48.4 (17.0 to 81.2)	2.86 (1.26 to 4.46)	48.7 (17.2 to 80.2)	2.87 (1.26 to 4.47)	-0.3% (-1.2% to 0.7%)	0.0% (-1.2% to 1.2%)	1.22% (0.2% to 2.2%)	122.8 (49.2 to 196.4)	6.06 (2.56 to 9.56)	129.3 (49.2 to 209.4)	6.09 (2.56 to 9.62)	0.6% (-0.5% to 1.7%)	1.1% (-0.1% to 2.3%)	1.1% (-0.1% to 2.3%)
Tuberculosis	18.0 (6.4 to 30.0)	1.00 (0.40 to 1.60)	18.0 (6.4 to 30.0)	1.00 (0.40 to 1.60)	0.0% (-0.8% to 0.8%)	-0.2% (-1.0% to 0.6%)	-0.2% (-1.0% to 0.6%)	27.0 (10.0 to 44.0)	1.50 (0.60 to 2.40)	27.0 (10.0 to 44.0)	1.50 (0.60 to 2.40)	-0.2% (-1.0% to 0.6%)	-0.2% (-1.0% to 0.6%)	-0.2% (-1.0% to 0.6%)
Occupational exposure to sulphuric acid: All causes	1.0 (0.0 to 1.0)	0.00 (0.00 to 0.00)	5.3 (0.0 to 10.6)	0.29 (0.00 to 0.58)	437% (0.0% to 974%)	14.7% (0.0% to 29.4%)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	102.0 (0.0 to 204.0)	5.82 (0.00 to 11.6)	102.0 (0.0 to 204.0)	5.82 (0.00 to 11.6)	102.0 (0.0 to 204.0)
Larynx cancer	3.0 (1.0 to 5.0)	0.16 (0.06 to 0.26)	3.0 (1.0 to 5.0)	0.16 (0.06 to 0.26)	0.0% (-0.8% to 0.8%)	0.0% (-0.8% to 0.8%)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	102.0 (0.0 to 204.0)	5.82 (0.00 to 11.6)	102.0 (0.0 to 204.0)	5.82 (0.00 to 11.6)	102.0 (0.0 to 204.0)
Occupational exposure to trichloroethylene: All causes	0.0 (0.0 to 0.0)	0.00 (0.00 to 0.00)	0.0 (0.0 to 0.0)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)
Kidney cancer	0.0 (0.0 to 0.0)	0.00 (0.00 to 0.00)	0.0 (0.0 to 0.0)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)
Occupational asbestos: All causes	36.8 (20.0 to 53.6)	0.16 (0.06 to 0.26)	37.0 (20.0 to 54.0)	0.16 (0.06 to 0.26)	0.5% (-0.5% to 1.5%)	0.0% (-0.8% to 0.8%)	2.12% (0.2% to 4.0%)	212.0 (100.0 to 324.0)	11.7 (5.8 to 17.6)	212.0 (100.0 to 324.0)	11.7 (5.8 to 17.6)	0.5% (-0.5% to 1.5%)	0.0% (-0.8% to 0.8%)	0.5% (-0.5% to 1.5%)
Arthritis	56.8 (20.0 to 93.6)	0.26 (0.06 to 0.46)	57.0 (20.0 to 94.0)	0.26 (0.06 to 0.46)	0.3% (-0.7% to 1.3%)	0.0% (-0.8% to 0.8%)	2.12% (0.2% to 4.0%)	212.0 (100.0 to 324.0)	11.7 (5.8 to 17.6)	212.0 (100.0 to 324.0)	11.7 (5.8 to 17.6)	0.3% (-0.7% to 1.3%)	0.0% (-0.8% to 0.8%)	0.3% (-0.7% to 1.3%)
Occupational particulate matter, gases, and fumes: All causes	40.7 (18.0 to 63.6)	0.00 (0.00 to 0.00)	424.3 (180.0 to 668.6)	0.00 (0.00 to 0.00)	942% (780% to 1100%)	0.0% (-0.8% to 0.8%)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)
Chronic obstructive pulmonary disease	331.0 (231.0 to 431.0)	16.0 (11.0 to 21.0)	331.0 (231.0 to 431.0)	16.0 (11.0 to 21.0)	0.0% (-0.8% to 0.8%)	0.0% (-0.8% to 0.8%)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)
Cutaneous melanoma	0.0 (0.0 to 0.0)	0.00 (0.00 to 0.00)	0.0 (0.0 to 0.0)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)
Other pneumonias	4.0 (1.0 to 7.0)	0.00 (0.00 to 0.00)	4.0 (1.0 to 7.0)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)
Occupational noise: All causes	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Aggregated and other hearing loss	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Occupational isotope: All causes	33.0 (34.0 to 32.0)	0.76 (0.00 to 1.52)	33.0 (34.0 to 32.0)	0.76 (0.00 to 1.52)	0.0% (-0.8% to 0.8%)	-0.5% (-1.3% to 0.3%)	-0.7% (-1.5% to 0.1%)	21.0 (21.0 to 21.0)	1.16 (0.00 to 2.32)	21.0 (21.0 to 21.0)	1.16 (0.00 to 2.32)	0.0% (-0.8% to 0.8%)	-0.5% (-1.3% to 0.3%)	-0.7% (-1.5% to 0.1%)
Pediatric meningitis	0.0 (0.0 to 0.0)	0.00 (0.00 to 0.00)	0.0 (0.0 to 0.0)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)
Cyctic oral tipitis	0.0 (0.0 to 0.0)	0.00 (0.00 to 0.00)	0.0 (0.0 to 0.0)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)
Meningococcal meningitis	0.0 (0.0 to 0.0)	0.00 (0.00 to 0.00)	0.0 (0.0 to 0.0)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)
Measles virus oral tipitis	0.0 (0.0 to 0.0)	0.00 (0.00 to 0.00)	0.0 (0.0 to 0.0)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)
Other oral tipitis	0.0 (0.0 to 0.0)	0.00 (0.00 to 0.00)	0.0 (0.0 to 0.0)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)
Other respiratory tipitis	0.0 (0.0 to 0.0)	0.00 (0.00 to 0.00)	0.0 (0.0 to 0.0)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)
Falk	0.0 (0.0 to 0.0)	0.00 (0.00 to 0.00)	0.0 (0.0 to 0.0)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)
Drowning	20.0 (10.0 to 30.0)	0.00 (0.00 to 0.00)	20.0 (10.0 to 30.0)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)
Fire, hot, and hot substances	0.0 (0.0 to 0.0)	0.00 (0.00 to 0.00)	0.0 (0.0 to 0.0)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)
Pneumonia	0.0 (0.0 to 0.0)	0.00 (0.00 to 0.00)	0.0 (0.0 to 0.0)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)
Unintentional firearm injuries	4.0 (1.0 to 7.0)	0.00 (0.00 to 0.00)	4.0 (1.0 to 7.0)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)
Unintentional self-harm	0.0 (0.0 to 0.0)	0.00 (0.00 to 0.00)	0.0 (0.0 to 0.0)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)
Other exposure to mechanical forces	0.0 (0.0 to 0.0)	0.00 (0.00 to 0.00)	0.0 (0.0 to 0.0)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)
Verminous animal contact	0.0 (0.0 to 0.0)	0.00 (0.00 to 0.00)	0.0 (0.0 to 0.0)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)
Non-verminous animal contact	0.0 (0.0 to 0.0)	0.00 (0.00 to 0.00)	0.0 (0.0 to 0.0)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)
Pneumonia and foreign body in airway	0.0 (0.0 to 0.0)	0.00 (0.00 to 0.00)	0.0 (0.0 to 0.0)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)
Foreign body in other body part	0.0 (0.0 to 0.0)	0.00 (0.00 to 0.00)	0.0 (0.0 to 0.0)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)
Other anatomical injuries	0.0 (0.0 to 0.0)	0.00 (0.00 to 0.00)	0.0 (0.0 to 0.0)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)
Occupational exposure to force: All causes	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Low back pain	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Behavioral risks: All causes	23,912 (22,222 to 25,602)	42.00 (36.00 to 48.00)	23,912 (22,222 to 25,602)	42.00 (36.00 to 48.00)	0.0% (-0.8% to 0.8%)	-0.5% (-1.3% to 0.3%)	-0.5% (-1.3% to 0.3%)	1,000,000 (900,000 to 1,100,000)	6.00 (5.00 to 7.00)	1,000,000 (900,000 to 1,100,000)	6.00 (5.00 to 7.00)	0.0% (-0.8% to 0.8%)	-0.5% (-1.3% to 0.3%)	-0.5% (-1.3% to 0.3%)
Child and maternal malnutrition: All causes	4,301 (4,107 to 4,495)	8.26 (7.06 to 9.46)	4,301 (4,107 to 4,495)	8.26 (7.06 to 9.46)	0.0% (-0.8% to 0.8%)	-0.5% (-1.3% to 0.3%)	-0.5% (-1.3% to 0.3%)	400,000 (360,000 to 440,000)	1.00 (0.80 to 1.20)	400,000 (360,000 to 440,000)	1.00 (0.80 to 1.20)	0.0% (-0.8% to 0.8%)	-0.5% (-1.3% to 0.3%)	-0.5% (-1.3% to 0.3%)
Isolated breastfeeding: All causes	23.0 (22.0 to 24.0)	0.00 (0.00 to 0.00)	23.0 (22.0 to 24.0)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)
Non-exclusive breastfeeding: All causes	242.0 (230.0 to 254.0)	0.00 (0.00 to 0.00)	242.0 (230.0 to 254.0)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.00 (0.00 to 0.00)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)	0.0% (0.0% to 0.0%)
Dietary diversity	0.0 (0.0 to 0.0)	0.00 (0.00 to 0.00)	0.0 (0.0 to 0.0)	0.00 (0.00 to 0.00)										

Supplementary Results Table 3. AE-age deaths, death PAFs, DALYs, DALY PAFs, and percentage change for each risk factor and outcome for both sexes combined in 2006 and 2016 for Global.

[illegible]

Supplementary Results Table 3. All-cause deaths, death FAPs, DALYs, DALY FAPs, and percentage change for each risk factor and outcome for both sexes combined in 2006 and 2016 for China.

		2006 Deaths (in thousands)	2006 Deaths FAP	2016 Deaths (in thousands)	2016 Deaths FAP	Percentage change Deaths (2006-2016)	Percentage change Deaths FAP (2006-2016)	Percentage change age-standardized Deaths (2006-2016)	2006 DALYs (in thousands)	2006 DALYs FAP	2016 DALYs (in thousands)	2016 DALYs FAP	Percentage change DALYs (2006-2016)	Percentage change DALYs FAP (2006-2016)	Percentage change DALYs (2006-2016)
Breast cancer	52.7	11.7%	9.2	12.7	10.7%	-12.7%	-12.7%	-12.7%	1.613	1.613	1.613	1.613	-2.7%	-2.7%	-14.4%
	(41.6 to 63.8)	(7.7 to 15.7)	(4.7 to 14.7)	(8.7 to 16.7)	(5.7 to 17.7)	(-1.7 to 11.7)	(-1.7 to 11.7)	(-1.7 to 11.7)	(0.9 to 1.9)	(0.9 to 1.9)	(0.9 to 1.9)	(0.9 to 1.9)	(-3.7 to 3.7)	(-3.7 to 3.7)	(-18.4 to -8.4)
Ischemic heart disease	58.8	-4.7%	-24.2	-4.7%	-4.7%	-4.7%	-4.7%	-4.7%	92.4	92.4	92.4	92.4	-4.7%	-4.7%	-10.7%
	(21.1 to 106.5)	(-2.7 to 18.7)	(-10.7 to 42.7)	(-2.7 to 18.7)	(-10.7 to 42.7)	(-12.7 to 3.7)	(-12.7 to 3.7)	(-12.7 to 3.7)	(32.4 to 152.4)	(32.4 to 152.4)	(32.4 to 152.4)	(32.4 to 152.4)	(-12.7 to 3.7)	(-12.7 to 3.7)	(-14.7 to -6.7)
Ischemic stroke	176.0	4.7%	13.2	6.7%	16.7%	6.7%	-11.7%	-11.7%	2.987	2.987	2.987	2.987	16.7%	16.7%	-10.1%
	(141.7 to 210.7)	(3.7 to 5.7)	(1.7 to 15.7)	(3.7 to 5.7)	(1.7 to 15.7)	(-1.7 to 11.7)	(-1.7 to 11.7)	(-1.7 to 11.7)	(2.487 to 3.487)	(2.487 to 3.487)	(2.487 to 3.487)	(2.487 to 3.487)	(16.7 to 16.7)	(16.7 to 16.7)	(-10.1 to -0.1)
Hypertensive stroke	497.7	10.7%	497.7	16.7%	16.7%	7.6%	16.7%	16.7%	16.7%	16.7%	16.7%	16.7%	16.7%	16.7%	16.7%
	(317.7 to 677.7)	(11.7 to 19.7)	(317.7 to 677.7)	(12.7 to 20.7)	(12.7 to 20.7)	(1.7 to 15.7)	(1.7 to 15.7)	(1.7 to 15.7)	(16.7 to 16.7)	(16.7 to 16.7)	(16.7 to 16.7)	(16.7 to 16.7)	(16.7 to 16.7)	(16.7 to 16.7)	(16.7 to 16.7)
Hypertensive heart disease	96.1	14.7%	14.7%	14.7%	14.7%	14.7%	14.7%	14.7%	14.7%	14.7%	14.7%	14.7%	14.7%	14.7%	14.7%
	(66.1 to 136.1)	(10.7 to 18.7)	(66.1 to 136.1)	(10.7 to 18.7)	(10.7 to 18.7)	(1.7 to 15.7)	(1.7 to 15.7)	(1.7 to 15.7)	(14.7 to 14.7)	(14.7 to 14.7)	(14.7 to 14.7)	(14.7 to 14.7)	(14.7 to 14.7)	(14.7 to 14.7)	(14.7 to 14.7)
Alcohol cirrhosis and fatty liver	96.1	14.7%	14.7%	14.7%	14.7%	14.7%	14.7%	14.7%	14.7%	14.7%	14.7%	14.7%	14.7%	14.7%	14.7%
	(66.1 to 136.1)	(10.7 to 18.7)	(66.1 to 136.1)	(10.7 to 18.7)	(10.7 to 18.7)	(1.7 to 15.7)	(1.7 to 15.7)	(1.7 to 15.7)	(14.7 to 14.7)	(14.7 to 14.7)	(14.7 to 14.7)	(14.7 to 14.7)	(14.7 to 14.7)	(14.7 to 14.7)	(14.7 to 14.7)
Atrial fibrillation and flutter	17.6	10.7%	20.7	10.7%	10.7%	-4.7%	-4.7%	-4.7%	92.4	92.4	92.4	92.4	10.7%	10.7%	6.7%
	(11.6 to 23.6)	(7.7 to 13.7)	(11.6 to 23.6)	(7.7 to 13.7)	(7.7 to 13.7)	(-1.7 to 11.7)	(-1.7 to 11.7)	(-1.7 to 11.7)	(32.4 to 152.4)	(32.4 to 152.4)	(32.4 to 152.4)	(32.4 to 152.4)	(10.7 to 10.7)	(10.7 to 10.7)	(6.7 to 16.7)
Coronary and other chronic heart diseases due to atherosclerosis	284.7	10.7%	51.7	10.7%	10.7%	10.7%	10.7%	10.7%	6.054	6.054	6.054	6.054	10.7%	10.7%	-11.8%
	(214.7 to 354.7)	(10.7 to 10.7)	(214.7 to 354.7)	(10.7 to 10.7)	(10.7 to 10.7)	(1.7 to 15.7)	(1.7 to 15.7)	(1.7 to 15.7)	(6.054 to 6.054)	(6.054 to 6.054)	(6.054 to 6.054)	(6.054 to 6.054)	(10.7 to 10.7)	(10.7 to 10.7)	(-11.8 to -11.8)
Pancreatic	32.0	10.7%	32.0	10.7%	10.7%	10.7%	10.7%	10.7%	10.7%	10.7%	10.7%	10.7%	10.7%	10.7%	10.7%
	(25.0 to 39.0)	(10.7 to 10.7)	(25.0 to 39.0)	(10.7 to 10.7)	(10.7 to 10.7)	(1.7 to 15.7)	(1.7 to 15.7)	(1.7 to 15.7)	(10.7 to 10.7)	(10.7 to 10.7)	(10.7 to 10.7)	(10.7 to 10.7)	(10.7 to 10.7)	(10.7 to 10.7)	(10.7 to 10.7)
Esophy	289.7	17.6%	17.6%	17.6%	17.6%	17.6%	17.6%	17.6%	17.6%	17.6%	17.6%	17.6%	17.6%	17.6%	17.6%
	(114.7 to 464.7)	(12.7 to 22.7)	(114.7 to 464.7)	(12.7 to 22.7)	(12.7 to 22.7)	(1.7 to 15.7)	(1.7 to 15.7)	(1.7 to 15.7)	(17.6 to 17.6)	(17.6 to 17.6)	(17.6 to 17.6)	(17.6 to 17.6)	(17.6 to 17.6)	(17.6 to 17.6)	(17.6 to 17.6)
Alcohol use disorders	171.6	10.7%	171.6	10.7%	10.7%	10.7%	10.7%	10.7%	171.6	171.6	171.6	171.6	10.7%	10.7%	10.7%
	(107.7 to 235.7)	(10.7 to 10.7)	(107.7 to 235.7)	(10.7 to 10.7)	(10.7 to 10.7)	(1.7 to 15.7)	(1.7 to 15.7)	(1.7 to 15.7)	(171.6 to 171.6)	(171.6 to 171.6)	(171.6 to 171.6)	(171.6 to 171.6)	(10.7 to 10.7)	(10.7 to 10.7)	(10.7 to 10.7)
Diabetes mellitus	74.0	10.7%	74.0	10.7%	10.7%	10.7%	10.7%	10.7%	74.0	74.0	74.0	74.0	10.7%	10.7%	10.7%
	(49.0 to 99.0)	(10.7 to 10.7)	(49.0 to 99.0)	(10.7 to 10.7)	(10.7 to 10.7)	(1.7 to 15.7)	(1.7 to 15.7)	(1.7 to 15.7)	(74.0 to 74.0)	(74.0 to 74.0)	(74.0 to 74.0)	(74.0 to 74.0)	(10.7 to 10.7)	(10.7 to 10.7)	(10.7 to 10.7)
Polycystic ovary syndrome	65.0	12.7%	65.0	12.7%	12.7%	12.7%	12.7%	12.7%	65.0	65.0	65.0	65.0	12.7%	12.7%	12.7%
	(37.0 to 93.0)	(11.7 to 13.7)	(37.0 to 93.0)	(11.7 to 13.7)	(11.7 to 13.7)	(1.7 to 15.7)	(1.7 to 15.7)	(1.7 to 15.7)	(65.0 to 65.0)	(65.0 to 65.0)	(65.0 to 65.0)	(65.0 to 65.0)	(12.7 to 12.7)	(12.7 to 12.7)	(12.7 to 12.7)
Cystic nodules	17.6	10.7%	17.6	10.7%	10.7%	10.7%	10.7%	10.7%	17.6	17.6	17.6	17.6	10.7%	10.7%	10.7%
	(11.6 to 23.6)	(7.7 to 13.7)	(11.6 to 23.6)	(7.7 to 13.7)	(7.7 to 13.7)	(-1.7 to 11.7)	(-1.7 to 11.7)	(-1.7 to 11.7)	(17.6 to 17.6)	(17.6 to 17.6)	(17.6 to 17.6)	(17.6 to 17.6)	(10.7 to 10.7)	(10.7 to 10.7)	(10.7 to 10.7)
Menstrual cycle disorders	32.0	10.7%	32.0	10.7%	10.7%	10.7%	10.7%	10.7%	32.0	32.0	32.0	32.0	10.7%	10.7%	10.7%
	(18.0 to 46.0)	(10.7 to 10.7)	(18.0 to 46.0)	(10.7 to 10.7)	(10.7 to 10.7)	(1.7 to 15.7)	(1.7 to 15.7)	(1.7 to 15.7)	(32.0 to 32.0)	(32.0 to 32.0)	(32.0 to 32.0)	(32.0 to 32.0)	(10.7 to 10.7)	(10.7 to 10.7)	(10.7 to 10.7)
Motor vehicle road injuries	45.0	12.7%	45.0	12.7%	12.7%	12.7%	12.7%	12.7%	45.0	45.0	45.0	45.0	12.7%	12.7%	12.7%
	(30.0 to 60.0)	(10.7 to 10.7)	(30.0 to 60.0)	(10.7 to 10.7)	(10.7 to 10.7)	(1.7 to 15.7)	(1.7 to 15.7)	(1.7 to 15.7)	(45.0 to 45.0)	(45.0 to 45.0)	(45.0 to 45.0)	(45.0 to 45.0)	(12.7 to 12.7)	(12.7 to 12.7)	(12.7 to 12.7)
Other road injuries	17.6	10.7%	17.6	10.7%	10.7%	10.7%	10.7%	10.7%	17.6	17.6	17.6	17.6	10.7%	10.7%	10.7%
	(11.6 to 23.6)	(7.7 to 13.7)	(11.6 to 23.6)	(7.7 to 13.7)	(7.7 to 13.7)	(-1.7 to 11.7)	(-1.7 to 11.7)	(-1.7 to 11.7)	(17.6 to 17.6)	(17.6 to 17.6)	(17.6 to 17.6)	(17.6 to 17.6)	(10.7 to 10.7)	(10.7 to 10.7)	(10.7 to 10.7)
Other transport injuries	17.6	10.7%	17.6	10.7%	10.7%	10.7%	10.7%	10.7%	17.6	17.6	17.6	17.6	10.7%	10.7%	10.7%
	(11.6 to 23.6)	(7.7 to 13.7)	(11.6 to 23.6)	(7.7 to 13.7)	(7.7 to 13.7)	(-1.7 to 11.7)	(-1.7 to 11.7)	(-1.7 to 11.7)	(17.6 to 17.6)	(17.6 to 17.6)	(17.6 to 17.6)	(17.6 to 17.6)	(10.7 to 10.7)	(10.7 to 10.7)	(10.7 to 10.7)
Drowning	16.7	10.7%	16.7	10.7%	10.7%	10.7%	10.7%	10.7%	16.7	16.7	16.7	16.7	10.7%	10.7%	10.7%
	(11.7 to 21.7)	(7.7 to 13.7)	(11.7 to 21.7)	(7.7 to 13.7)	(7.7 to 13.7)	(-1.7 to 11.7)	(-1.7 to 11.7)	(-1.7 to 11.7)	(16.7 to 16.7)	(16.7 to 16.7)	(16.7 to 16.7)	(16.7 to 16.7)	(10.7 to 10.7)	(10.7 to 10.7)	(10.7 to 10.7)
Fire, heat, and hot substances	7.0	5.7%	7.0	5.7%	5.7%	5.7%	5.7%	5.7%	7.0	7.0	7.0	7.0	5.7%	5.7%	5.7%
	(5.0 to 9.0)	(4.7 to 6.7)	(5.0 to 9.0)	(4.7 to 6.7)	(4.7 to 6.7)	(1.7 to 15.7)	(1.7 to 15.7)	(1.7 to 15.7)	(7.0 to 7.0)	(7.0 to 7.0)	(7.0 to 7.0)	(7.0 to 7.0)	(5.7 to 5.7)	(5.7 to 5.7)	(5.7 to 5.7)
Poisoning	3.0	5.7%	3.0	5.7%	5.7%	5.7%	5.7%	5.7%	3.0	3.0	3.0	3.0	5.7%	5.7%	5.7%
	(1.0 to 5.0)	(4.7 to 6.7)	(1.0 to 5.0)	(4.7 to 6.7)	(4.7 to 6.7)	(1.7 to 15.7)	(1.7 to 15.7)	(1.7 to 15.7)	(3.0 to 3.0)	(3.0 to 3.0)	(3.0 to 3.0)	(3.0 to 3.0)	(5.7 to 5.7)	(5.7 to 5.7)	(5.7 to 5.7)
Unintentional firearm injuries	1.0	5.7%	1.0	5.7%	5.7%	5.7%	5.7%	5.7%	1.0	1.0	1.0	1.0	5.7%	5.7%	5.7%
	(0.0 to 2.0)	(4.7 to 6.7)	(0.0 to 2.0)	(4.7 to 6.7)	(4.7 to 6.7)	(1.7 to 15.7)	(1.7 to 15.7)	(1.7 to 15.7)	(1.0 to 1.0)	(1.0 to 1.0)	(1.0 to 1.0)	(1.0 to 1.0)	(5.7 to 5.7)	(5.7 to 5.7)	(5.7 to 5.7)
Other unintentional injuries	1.0	5.7%	1.0	5.7%	5.7%	5.7%	5.7%	5.7%	1.0	1.0	1.0	1.0	5.7%	5.7%	5.7%
	(0.0 to 2.0)	(4.7 to 6.7)	(0.0 to 2.0)	(4.7 to 6.7)	(4.7 to 6.7)	(1.7 to 15.7)	(1.7 to 15.7)	(1.7 to 15.7)	(1.0 to 1.0)	(1.0 to 1.0)	(1.0 to 1.0)	(1.0 to 1.0)	(5.7 to 5.7)	(5.7 to 5.7)	(5.7 to 5.7)
Self-harm by firearm	1.0	5.7%	1.0	5.7%	5.7%	5.7%	5.7%	5.7%	1.0	1.0	1.0	1.0	5.7%	5.7%	5.7%
	(0.0 to 2.0)	(4.7 to 6.7)	(0.0 to 2.0)	(4.7 to 6.7)	(4.7 to 6.7)	(1.7 to 15.7)	(1.7 to 15.7)	(1.7 to 15.7)	(1.0 to 1.0)	(1.0 to 1.0)	(1.0 to 1.0)	(1.0 to 1.0)	(5.7 to 5.7)	(5.7 to 5.7)	(5.7 to 5.7)
Self-harm by other specified means	1.0	5.7%	1.0	5.7%	5.7%	5.7%	5.7%	5.7%	1.0	1.0	1.0	1.0	5.7%	5.7%	5.7%
	(0.0 to 2.0)	(4.7 to 6.7)	(0.0 to 2.0)	(4.7 to 6.7)	(4.7 to 6.7)	(1.7 to 15.7)	(1.7 to 15.7)	(1.7 to 15.7)	(1.0 to 1.0)	(1.0 to 1.0)	(1.0 to 1.0)	(1.0 to 1.0)	(5.7 to 5.7)	(5.7 to 5.7)	(5.7 to 5.7)
Suicide by other specified means	1.0	5.7%	1.0	5.7%	5.7%	5.7%	5.7%	5.7%	1.0	1.0	1.0	1.0	5.7%	5.7%	5.7%
	(0.0 to 2.0)	(4.7 to 6.7)	(0.0 to 2.0)	(4.7 to 6.7)	(4.7 to 6.7)	(1.7 to 15.7)	(1.7 to 15.7)	(1.7 to 15.7)	(1.0 to 1.0)	(1.0 to 1.0)	(1.0 to 1.0)	(1.0 to 1.0)	(5.7 to 5.7)	(5.7 to 5.7)	(5.7 to 5.7)
Drug use, all causes	45.0	12.7%	45.0	12.7%	12.7%	12.7%	12.7%	12.7%	45.0	45.0	45.0	45.0	12.7%	12.7%	12.7%
	(30.0 to 60.0)	(10.7 to 10.7)	(30.0 to 60.0)	(10.7 to 10.7)	(10.7 to 10.7)	(1.7 to 15.7)	(1.7 to 15.7)	(1.7 to 15.7)	(45.0 to 45.0)	(45.0 to 45.0)	(45.0 to 45.0)	(45.0 to 45.0)	(12.7 to 12.7)	(12.7 to 12.7)	(12.7 to 12.7)
Drug consumption: HIV/AIDS - Tuberculosis	17.6	10.7%	17.6	10.7%	10.7%	10.7%	10.7%	10.							

Supplementary Results Table 3. All-cause deaths, death FAFs, DALYs, DALY FAFs, and percentage change for each risk factor and outcome for both sexes combined in 2000 and 2016 for China.

	2000 Deaths (in thousands)	2000 DALYs (in thousands)	2000 Deaths (in thousands)	2000 DALYs (in thousands)	Percentage change Deaths (2000-2016)	Percentage change DALYs (2000-2016)	Percentage change Deaths (2000-2016)	2000 DALYs (in thousands)	2000 DALYs (in thousands)	2000 DALYs (in thousands)	2000 DALYs (in thousands)	Percentage change Deaths (2000-2016)	Percentage change DALYs (2000-2016)	Percentage change Deaths (2000-2016)	Percentage change DALYs (2000-2016)
Trachea, bronchus, and lung cancer	101.1 (22.7 to 172.6)	6.7 (1.6 to 12.7)	117.1 (25.1 to 189.1)	7.2 (1.7 to 12.7)	16.7 (1.5 to 31.1)	-1.2 (-2.9 to 0.5)	-1.2 (-2.9 to 0.5)	2,027.1 (1,126.1 to 2,928.1)	2,027.1 (1,126.1 to 2,928.1)	2,027.1 (1,126.1 to 2,928.1)	2,027.1 (1,126.1 to 2,928.1)	1.6 (-0.4 to 3.6)	1.6 (-0.4 to 3.6)	-12.7 (-15.8 to -9.6)	-12.7 (-15.8 to -9.6)
Breast cancer	28.1 (14.5 to 41.7)	5.6 (3.0 to 8.2)	31.1 (16.5 to 45.7)	6.7 (3.5 to 10.0)	5.7 (1.9 to 9.5)	-0.6 (-1.4 to 0.2)	-0.6 (-1.4 to 0.2)	67.4 (35.3 to 99.5)	67.4 (35.3 to 99.5)	67.4 (35.3 to 99.5)	67.4 (35.3 to 99.5)	1.6 (-0.4 to 3.6)	1.6 (-0.4 to 3.6)	2.6 (-0.4 to 5.6)	2.6 (-0.4 to 5.6)
Ovarian cancer	6.1 (1.1 to 11.1)	1.0 (0.2 to 1.8)	10.0 (3.0 to 17.0)	1.7 (0.4 to 3.0)	6.7 (1.7 to 11.7)	1.1 (-0.1 to 2.3)	1.1 (-0.1 to 2.3)	10.0 (3.0 to 17.0)	10.0 (3.0 to 17.0)	10.0 (3.0 to 17.0)	10.0 (3.0 to 17.0)	2.0 (-0.1 to 4.1)	2.0 (-0.1 to 4.1)	2.0 (-0.1 to 4.1)	2.0 (-0.1 to 4.1)
Bladder cancer	10.8 (2.0 to 19.6)	1.5 (0.5 to 2.5)	11.5 (2.1 to 20.9)	1.6 (0.6 to 2.6)	0.7 (-0.3 to 1.7)	0.2 (-0.8 to 1.2)	0.2 (-0.8 to 1.2)	22.4 (8.0 to 36.8)	22.4 (8.0 to 36.8)	22.4 (8.0 to 36.8)	22.4 (8.0 to 36.8)	0.6 (-0.4 to 1.6)	0.6 (-0.4 to 1.6)	0.6 (-0.4 to 1.6)	0.6 (-0.4 to 1.6)
Infectious heart disease	1,876.2 (1,479.2 to 2,273.2)	1,481.2 (1,151.2 to 1,811.2)	1,481.2 (1,151.2 to 1,811.2)	1,481.2 (1,151.2 to 1,811.2)	0.0 (-0.1 to 0.1)	0.0 (-0.1 to 0.1)	0.0 (-0.1 to 0.1)	1,481.2 (1,151.2 to 1,811.2)	1,481.2 (1,151.2 to 1,811.2)	1,481.2 (1,151.2 to 1,811.2)	1,481.2 (1,151.2 to 1,811.2)	0.0 (-0.1 to 0.1)	0.0 (-0.1 to 0.1)	0.0 (-0.1 to 0.1)	0.0 (-0.1 to 0.1)
Infectious stroke	1,481.2 (1,151.2 to 1,811.2)	1,481.2 (1,151.2 to 1,811.2)	1,481.2 (1,151.2 to 1,811.2)	1,481.2 (1,151.2 to 1,811.2)	0.0 (-0.1 to 0.1)	0.0 (-0.1 to 0.1)	0.0 (-0.1 to 0.1)	1,481.2 (1,151.2 to 1,811.2)	1,481.2 (1,151.2 to 1,811.2)	1,481.2 (1,151.2 to 1,811.2)	1,481.2 (1,151.2 to 1,811.2)	0.0 (-0.1 to 0.1)	0.0 (-0.1 to 0.1)	0.0 (-0.1 to 0.1)	0.0 (-0.1 to 0.1)
Ischemic heart disease	464.1 (369.1 to 559.1)	17.0 (13.0 to 21.0)	471.1 (376.1 to 566.1)	17.7 (13.7 to 21.7)	1.5 (-0.5 to 3.5)	-0.2 (-1.2 to 0.8)	-0.2 (-1.2 to 0.8)	10,703.1 (8,583.1 to 12,823.1)	10,703.1 (8,583.1 to 12,823.1)	10,703.1 (8,583.1 to 12,823.1)	10,703.1 (8,583.1 to 12,823.1)	0.6 (-0.4 to 1.6)	0.6 (-0.4 to 1.6)	-2.0 (-3.0 to -1.0)	-2.0 (-3.0 to -1.0)
Peripheral vascular disease	12.0 (6.0 to 18.0)	1.0 (0.5 to 1.5)	11.7 (5.7 to 17.7)	0.9 (0.4 to 1.4)	-0.3 (-1.3 to 0.7)	-0.1 (-1.1 to 0.9)	-0.1 (-1.1 to 0.9)	23.0 (13.0 to 33.0)	23.0 (13.0 to 33.0)	23.0 (13.0 to 33.0)	23.0 (13.0 to 33.0)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	-0.4 (-1.4 to 0.6)	-0.4 (-1.4 to 0.6)
Alzheimer's disease and other dementias	8.0 (4.0 to 12.0)	0.7 (0.3 to 1.1)	7.7 (3.7 to 11.7)	0.6 (0.2 to 1.0)	-0.3 (-1.3 to 0.7)	-0.1 (-1.1 to 0.9)	-0.1 (-1.1 to 0.9)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	-0.4 (-1.4 to 0.6)	-0.4 (-1.4 to 0.6)
Dementia	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)
Chronic kidney disease due to diabetes mellitus	364.1 (169.1 to 559.1)	10.0 (5.0 to 15.0)	364.1 (169.1 to 559.1)	10.0 (5.0 to 15.0)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	1,151.2 (651.2 to 1,651.2)	1,151.2 (651.2 to 1,651.2)	1,151.2 (651.2 to 1,651.2)	1,151.2 (651.2 to 1,651.2)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)
Chronic kidney disease due to hypertension	101.2 (51.2 to 151.2)	4.0 (2.0 to 6.0)	101.2 (51.2 to 151.2)	4.0 (2.0 to 6.0)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	2,123.1 (1,323.1 to 2,923.1)	2,123.1 (1,323.1 to 2,923.1)	2,123.1 (1,323.1 to 2,923.1)	2,123.1 (1,323.1 to 2,923.1)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)
Chronic kidney disease due to glomerulonephritis	41.9 (21.9 to 61.9)	1.0 (0.5 to 1.5)	41.9 (21.9 to 61.9)	1.0 (0.5 to 1.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	1,151.2 (651.2 to 1,651.2)	1,151.2 (651.2 to 1,651.2)	1,151.2 (651.2 to 1,651.2)	1,151.2 (651.2 to 1,651.2)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)
Chronic kidney disease due to other causes	12.0 (6.0 to 18.0)	1.0 (0.5 to 1.5)	11.7 (5.7 to 17.7)	0.9 (0.4 to 1.4)	-0.3 (-1.3 to 0.7)	-0.1 (-1.1 to 0.9)	-0.1 (-1.1 to 0.9)	23.0 (13.0 to 33.0)	23.0 (13.0 to 33.0)	23.0 (13.0 to 33.0)	23.0 (13.0 to 33.0)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	-0.4 (-1.4 to 0.6)	-0.4 (-1.4 to 0.6)
Glaucoma	10.8 (5.8 to 15.8)	1.5 (0.5 to 2.5)	11.5 (6.5 to 16.5)	1.6 (0.6 to 2.6)	0.7 (-0.3 to 1.7)	0.2 (-0.8 to 1.2)	0.2 (-0.8 to 1.2)	22.4 (8.0 to 36.8)	22.4 (8.0 to 36.8)	22.4 (8.0 to 36.8)	22.4 (8.0 to 36.8)	0.6 (-0.4 to 1.6)	0.6 (-0.4 to 1.6)	0.6 (-0.4 to 1.6)	0.6 (-0.4 to 1.6)
Cataract	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)
High total cholesterol: All causes	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)
Ischemic heart disease	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)
Ischemic stroke	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)
High systolic blood pressure: All causes	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)
Ischemic heart disease	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)
Ischemic stroke	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)
Ischemic heart disease due to hypertension	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)
Ischemic heart disease due to glomerulonephritis	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)
Ischemic heart disease due to other causes	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)
Ischemic stroke due to hypertension	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)
Ischemic stroke due to glomerulonephritis	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)
Ischemic stroke due to other causes	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)
Ischemic stroke due to other causes	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)
Ischemic stroke due to other causes	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)
Ischemic stroke due to other causes	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)	0.0 (-0.5 to 0.5)
Ischemic stroke due to other causes	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to 2,669.1)	1,869.1 (1,069.1 to											

Supplementary Results Table 3. All-cause deaths, deaths PAFs, DALYs, DALY PAFs, and percentage change for each risk factor and outcome for both years combined in 2000 and 2016 for China.														
	2000 Deaths (in thousands)	2000 Deaths PAF	2016 Deaths (in thousands)	2016 Deaths PAF	Percentage change deaths (2000-2016)	Percentage change deaths PAF (2000-2016)	Percentage change age-standardized death rate (2000-2016)	2000 DALYs (in thousands)	2000 DALYs PAF	2016 DALYs (in thousands)	2016 DALYs PAF	Percentage change DALYs (2000-2016)	Percentage change DALYs PAF (2000-2016)	Percentage change age-standardized DALYs rate (2000-2016)
Grid	-	-	-	-	-	-	-	802.1 (703.9 to 915.3)	52.0%	517.8 (463.9 to 579.0)	52.9%	54.0%	6.0%	2.0%